

APPENDIX A

USFWS Species List

Yuma County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
California Brown pelican	<i>Pelecanus occidentalis californicus</i>	Endangered	Large dark gray-brown water bird with a pouch underneath long bill and webbed feet. Adults have a white head and neck, brownish black breast, and silver gray upper parts.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	Varies	Coastal land and islands; species found around many Arizona lakes and rivers.	Subspecies is found on Pacific Coast and is endangered due to pesticides. It is an uncommon transient in Arizona on many Arizona lakes and rivers. Individuals wander up from Mexico in summer and fall. No breeding records in Arizona.
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Species is also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam; Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizon/New Mexico border to Coodidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.
Sonoran pronghorn	<i>Antilocapra americana sonoriensis</i>	Endangered	Buff on back and white below, hooped with slightly curved black horns having a single prong. Smallest and palest of the pronghorn subspecies	Maricopa, Pima, Yuma	500 - 2,000 ft	Broad intermountain alluvial valleys with creosote-bursage and palo verde-mixed cacti associations.	Typically, bajadas are used as fawning areas and sandy dune areas provide food seasonally. Historical range was probably larger than exists today. This subspecies also occurs in Mexico.
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	<8500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886) and can be viewed at http://arizonaes.fws.gov . In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Yuma clapper rail	<i>Rallus longirostris yumanensis</i>	Endangered	Water bird with long legs and short tail. Long, slender decurved bill. Mottled brown or gray on its rump. Flanks and undersides are dark gray with narrow vertical stripes producing a barring effect.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	< 4,500 ft	Fresh water and brackish marshes.	Species is associated with dense emergent riparian vegetation. Requires wet substrate (mudflat, sandbar) with dense herbaceous or woody vegetation for nesting and foraging. Channelization and marsh destruction are primary sources of habitat loss.
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill, which is blue-black with yellow on the lower half of the bill. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Listing was found warranted, but precluded as a distinct vertebrate population segment in the western U.S. on July 25, 2001. This finding indicates that the Service has sufficient information to list the bird, but other, higher priority listing actions prevent the Service from addressing the listing of the cuckoo at this time.
Flat-tailed horned lizard	<i>Phrynosoma mcallii</i>	Conservation Agreement	Typical flattened body shape of horned lizards; dark vertebral stripe; lacks external ear openings; color is cryptic ranging from pale gray to light rust brown; has two rows of fringed scales on each side of body.	Yuma	500 ft	Sandy flats or areas with fine, windblown sand; creosote-white bursage series of Sonoran Desert.	Conservation Agreement finalized in May 1997. Species also found in portions of San Diego County, central Riverside County, and Imperial County, California; also Sonora and Baja California, Mexico.