



Figure 4. Known key use area within the Menan Buttes bald eagle breeding area, South Fork Snake River. Intensive monitoring has not occurred at this breeding area, and the information portrayed is preliminary only. The circle encloses the Principal Management Area (PMA). The star indicates the current nest site. Dots and numbers indicate prominent foraging perches/locations determined in 1998 observations. (Two historic alternate nests in an old heron rookery approximately 3/4 mile upriver of current nest are not shown within the PMA. The area of the historic alternates should also be managed as a part of the PMA.)



Figure 5. Known key use area within the Cartier Slough bald eagle breeding area, Henry's Fork Snake River. Intensive monitoring has not occurred at this breeding area, and the information portrayed is preliminary only. The circle encloses the Principal Management Area. The three stars indicate the known nest sites, the highest and lowest of which are still intact. (The 1998 nest was the lowest star.) The numbers refer to primary foraging perches and locations. Perch 1 is 1.5 miles southeast of the photo; perch 2 is 2.2 miles to the northeast.



Figure 6. Known key use area within the Annis Slough bald eagle breeding area, Henry's Fork Snake River. Intensive monitoring has not occurred at this breeding area, and the information portrayed is preliminary only. The circle encloses the Principal Management Area. The star indicates the only known nest site. The numbers indicate primary foraging perches used by the adults during the nesting season. One primary perch not shown in this figure is located downstream below the Menan Bridge over the Snake River in a small cottonwood stand on the south side of the river.

Figure 7




Figure 8

