1989.10

Review of Stream Flow Analyses in the USDI Bureau of Land Management (BLM) Draft Environmental Impact Statement (DEIS) for its Western Oregon Plan Revisions (WOPR)

December 28, 2007

Jonathan J. Rhodes
Conservation Hydrologist, Planeto Azul Hydrology
P.O. Box 15286 •• Portland, OR 97293-5286

I. INTRODUCTION

Management effects on peakflows are a significant issue within the analysis area of the DEIS for several reasons. Scientific assessments have repeatedly concluded that management effects on watershed-scale hydrology and peakflows affect aquatic conditions that strongly affect salmonid populations (USFS et al., 1993; Murphy, 1995; Spence et al., 1996). Studies have repeatedly demonstrated that logging and roads cumulatively elevate peakflows, especially in smaller watersheds.

Elevated peakflows have numerous negative impacts on stream conditions and processes, including increased sediment transport, bank erosion, channel scour, and sedimentation of downstream salmonid habitats. Elevated peakflows also contribute to channel widening, which contributes to increased summer water temperatures. High summer water temperatures are already a widespread problem for salmonid populations within the DEIS analysis area.

Proposed logging levels vary considerably among the alternatives analyzed in the DEIS. Hence, effects of the alternatives on peakflows will also vary among the alternatives, because logging and associated activities elevate peakflows. However, the DEIS failed to reasonably analyze and disclose the impacts of the alternatives on peakflows due to several defects in the analysis. These deficiencies include the following:

- The DEIS failed to use the results of Grant et al. (2007) which found that forest canopy removal of more than about 20% of watershed area elevated peakflows generated by rain-on-snow.

- The DEIS did not analyze the impacts of the existing conditions and the alternatives at scales where peakflow impacts are most pronounced and ecologically significant.

- The DEIS's analysis narrowly focused on the effects of forest canopy removal on peakflows and ignored other important causes of peakflow elevation, including cumulative soil compaction from roads, logging, and grazing, and the acceleration of runoff routing by roads.

- The DEIS's analysis is fraught with potential error, yet, the DEIS failed to assess and disclose the likely magnitude and implications of potential individual errors, nor of combining or compounding of error in the analysis, although this has long been standard scientific practice.

Each of these defects contributes to underestimation of the magnitude, extent, and significance of existing peakflow elevation within the analysis area under existing conditions. This is significant because impacts to already damaged systems can be considerably different and more ecologically serious than those in systems that have not been impaired (Reid, 1993; Dunne et al., 2001).

Each of the aforementioned deficiencies also contributes to underestimation of peakflow impacts under the alternatives. These defects also have a combined effect that contributes to underestimation of the magnitude, extent, and significance of peakflow impacts under the

alternatives. Therefore, the DEIS fails to adequately differentiate among the alternatives in terms of their effects on peakflows. Based on available information, it is highly likely that the action alternatives will elevate peakflows to a significantly greater degree than forecast in the DEIS. Because peakflows influence a host of aquatic conditions, the DEIS's failure to reasonably analyze and disclose peakflow impacts also causes the DEIS to fail to reasonably differentiate among the alternatives with respect to their impacts on aquatic habitat conditions and salmonids.

The foregoing defects in the DEIS's analysis of peakflows need to be rectified. The FEIS must analyze and disclose:

- All cumulative sources of peakflow elevation under existing conditions and the alternatives, including their extent and severity within the analysis area at scales where impacts are likely to be most pronounced;

- The uncertainties and other limitations inherent in the analysis approach, and their implications for accuracy and ecological consequences;

- The potential accuracy of the analysis, including its expected error, and their ramifications.

Low flows can also be reduced by the cumulative effects of management activities on BLM lands throughout the analysis area. Reductions in low flows negatively impact aquatic habitat conditions and salmonids. However, the DEIS is without any reasonable analysis of the cumulative management-induced impacts on low flows under existing conditions or the action alternatives. This significant defect must be rectified by taking a hard look at all sources of impacts to low flows and their cascading effects on aquatic conditions and salmonids.

II. THE DEIS'S ANALYSIS OF PEAKFLOW IMPACTS FAILED TO REASONABLY INCORPORATE AVAILABLE INFORMATION ON THE LEVELS OF FOREST CANOPY REMOVAL THAT INCREASE PEAKFLOWS GENERATED BY RAIN-ON-SNOW

The DEIS notes that "For basins within the rain-on-snow hydroregion, the detection threshold is 20% of a harvested watershed area (Grant et al. 2007)." However, the DEIS's completely fails to use this information to analyze the impacts of management activities on peakflows under existing conditions and the action alternatives. Instead, the DEIS relies on a relatively arcane method of estimating effects on peakflows from rain-on-snow, which has not been empirically verified, and based on assumptions and approximations that are fraught with potential error.

The failure to apply the results of Grant et al. (2007) with respect to logging impacts on peakflows is a significant flaw for several reasons. The results of Grant et al. (2007) are plainly applicable, because the DEIS used the results of Grant et al. (2007) in the DEIS's analysis of peakflow impacts in rain-dominated watersheds (DEIS, pp. 382-388, 1095-1105).

Although the DEIS discloses the findings in Grant et al. (2007) regarding the effects of logged areas on peakflows generated by rain-on-snow, the DEIS provides absolutely no rationale for

why these findings were not used in its analysis of peakflow impacts. Therefore, the failure to use the findings of Grant et al. (2007) with respect peakflow elevation in watersheds subject to rain-on-snow is plainly arbitrary.

Second, it is highly likely that many watersheds analyzed in the DEIS have had more than 20% of the watershed area logged, which Grant et al. (2007) indicates would detectably elevate peakflows in watersheds affected by rain-on-snow. Therefore, the failure to apply this information causes the DEIS to underestimate the magnitude and extent of increased peakflow under existing conditions and the action alternatives.

Third, the use of the threshold of detectable peakflow increase in Grant et al. (2007) is eminently tractable analytically. Such an analysis is much less onerous than the approach employed in the DEIS. It could also be easily applied at ecologically relevant scales, including watersheds smaller than those analyzed in the DEIS.

Fourth, peakflow from rain-on-snow events is environmentally significant. Some of the highest peakflows within the analysis area occur in response to rain-on-snow events. These events often act as catalyst for high levels of management-induced erosion and sediment delivery within the analysis area.

The DEIS has plainly ignored relevant scientific information by failing to apply the results of Grant et al. (2007) with respect to the effects of logging on peakflows generated by rain-on-snow. It is highly likely that the failure to incorporate these findings causes the DEIS to greatly underestimate the magnitude and extent of existing peakflow elevation. This failure also causes the DEIS to underestimate the differences among the alternatives with respect to their effects on peakflows generated by rain-on-snow.

III. THE ANALYTICAL SCALE OF THE DEIS'S ANALYIS OF PEAKFLOWS IS INADEQUATE FOR DISCLOSING PEAKFLOW EFFECTS UNDER EXISTING CONDITIONS AND THE ALTERNATIVES

There are several reasons why the DEIS analysis of peakflows at the scale of sixth-field watersheds (DEIS, pp. pp. 382-388, 1095-1105) is inadequate to reasonably analyze and disclose impacts on peakflows and their effects on important aquatic conditions. First, peakflows in smaller watersheds are more prone to elevation by logging (MacDonald and Coe, 2007). Studies have consistently demonstrated that logging activities elevate peakflows in smaller watersheds (MacDonald and Ritland, 1989; Bowling et al., 2000).

Second, smaller watersheds often have a greater percentage of their watershed area that has been recently logged. This higher percentage of watershed disturbance causes proportionately greater peakflow increases in these smaller watersheds. Due to the patchiness in the distribution of logged areas, analysis at larger scales does not capture the more intensive disturbance levels existing in smaller watershed systems that significantly elevate peakflows. Thus, the DEIS's scale of analysis obscures, rather than discloses, the effects of logging on peakflows in headwater systems in watersheds smaller than the sixth-field scale analyzed in the DEIS.

Third, increases in peakflows in smaller watersheds have significant impacts. Some channel types in headwaters are highly vulnerable to increased channel erosion caused by peakflow elevation (Rosgen, 1996). Once degraded, many headwater streams in have very poor prospects for recovery, even after the causes of degradation have been eliminated (Rosgen, 1996). Due to their position in the channel network, elevated erosion in headwater channels increases downstream sediment transport and sedimentation in downstream fish habitats (Montgomery and Buffington, 1998).

For these reasons, the DEIS's analysis of peakflows effects at the scale of sixth-field watersheds is inadequate for assessing and disclosing the impacts on peakflows under existing conditions and the action alternatives. Due to these defects, the DEIS has failed to reasonably differentiate among the alternatives with respect to effects on peakflows and, instead, has underestimated the extent and magnitude of peakflow elevation under existing conditions and the action alternatives.

These scale defects must be rectified in the FEIS by analysis at scales smaller than the 6th field. Such an analysis is tractable, especially using the results of Grant et al. (2007).

IV. THE DEIS UNDERESTIMATES IMPACTS ON PEAKFLOWS BECAUSE IT DOES NOT REASONABLY EVALUATE AND DISCLOSE THE EFFECTS OF ALL SOURCES OF INCREASED PEAKFLOWS DUE TO MANAGEMENT ACTIVITIES

The DEIS does not reasonably examine and disclose the effects of all management activities on peakflows due to several defects which synergistically cause the DEIS to underestimate the extent and magnitude of increases in peakflows under existing conditions and the action alternatives. The DEIS's analysis only narrowly considers some effects of forest vegetation conditions on peakflows, while ignoring other factors that have long been known to affect peakflows. For instance, soil compaction contributes to increased peakflow (Wissmar et al., 1994; USFS et al., 1993; Booth et al., 2002; Booth et al., 2004). Roads alter peakflow through additional mechanisms. However, the DEIS analysis does not reasonably analyze and disclose these impacts on peakflow alteration under existing conditions and the action alternatives. As a result, the DEIS's analysis of peakflow underestimates the extent and intensity of peakflow alteration under existing conditions and the alternatives.

Peakflow elevation by cumulative soil compaction

The DEIS fails to reasonably assess and disclose the effects of soil compaction on peakflows. The methods used in the DEIS to provide some nominal analysis of the extent of peakflow alteration within the analysis area do not explicitly incorporate the effects of soil compaction (DEIS, pp. 382-388, 1095-1105). Instead, the method only narrowly focuses on some impacts of forest canopy conditions (DEIS, pp. 382-388, 1095-1105).

This is a major defect. Soil compaction is an enduring cumulative impact of management activities. Soil compaction typically persists for many decades before full recovery (Beschta et al., 2004). There are many pervasive sources of soil compaction within the analysis area, including historic and on-going livestock grazing, previous logging, landings, and roads.

Soil compaction contributes to elevated runoff by decreasing infiltration rates and reducing the ability of the soil to store water (Maidment, 1993). The latter increases the duration and extent of saturated soils, increasing the magnitude and duration of surface runoff that contribute to elevated peakflow (O'Laughlin, 1986). Reductions in infiltration rates also contribute to increases in surface runoff that contribute to elevated peakflows.

Peakflow elevation by soil compaction from grazing

Livestock grazing significantly compacts soils (Reid, 1993; CWWR, 1996; Kauffman et al., 2004). The USBLM has estimated that the hooves of a 1,000 pound cow exert more than five times the pressure per square inch on soils and streambanks than that from a bulldozer (Cowley, 2002). Although the DEIS does not reasonably assess and disclose the cumulative impacts of grazing on soil compaction and peakflows, the DEIS concedes that compaction from livestock grazing decreases the ability of soils to absorb water and increases surface runoff (DEIS, p. 796-798).

The impacts of grazing on the hydrologic properties of soils that affect peakflows are far from trivial. Kauffman et al. (2004) documented that cattle grazing persistently reduced infiltration rates by about an average of 85% relative to areas that had not been grazed for more than a decade.

Kauffman et al. (2004) that noted that soil compaction from grazing profoundly reduced the ability of soils to absorb and store water:

> "Based upon the results of this study we calculated that saturated soils of the surface 10 cm of a single hectare of exclosed dry meadow would contain 61000 L more water than an equivalent grazed hectare. Under saturated conditions, a hectare of wet meadows with the pore space measured in the exclosed communities of this study would contain 121000 L more water than those with the pore space of the grazed wet-meadow communities. Our results suggest that if the entire area was excluded from livestock the surface 10 cm of soil in the meadows alone...could potentially store 16.6 X 10^6 L more of water than if the area were grazed by cattle. And, this estimate does not include the entire soil profile. This increase in soil water likely influences ecosystem productivity, soil temperature, biogeochemistry, and streamflows."

Kauffman et al. (2004) concluded that the measured impacts of grazing on the hydrologic properties of soils at the landscape scale had likely had significant effects on stream channel structure, water quality, and the aquatic biota.

CWWR (1996) noted that compaction is pervasive in areas that have been subjected to grazing. In some systems, grazing has rendered more than 80% of soils in compacted state (CWWR, 1996).

These hydrologic impacts of grazing are especially significant because livestock impacts are typically greatest in riparian areas near streams. Due to their proximity, this alteration of soil

hydrologic processes in riparian areas causes elevated surface runoff to streams, contributing to elevated peakflows. Grazing has been documented to increase peakflows at the watershed scale (Reid, 1993).

A significant amount of the analysis area has been grazed. Currently, about 560,000 acres, or roughly 22% of the BLM lands in the planning area are subjected to grazing (DEIS, p. 428). However, previous grazing has been more extensive (DEIS, pp. 428-429). Therefore, soils compacted by grazing likely occur over more than 22% of BLM lands covered by the DEIS, because soil compaction is highly persistent (Beschta et al., 2004).

The DEIS fails to reasonably disclose the extent of soils that have been compacted by grazing. This is a significant defect, due to the known impacts of grazing on soil compaction. This defect must be rectified in the FEIS by reasonably estimating the extent and distribution of soil compaction caused by on-going grazing and by grazing over the past several decades.

The DEIS also fails to adequately analyze and disclose how soil compaction from grazing contributes to peakflow increases. The DEIS compounds these problems by failing to clearly disclose that the DEIS's analysis of peakflows does not consider impacts from existing and future soil impacts caused by grazing. The DEIS also fails to reasonably disclose that this defect in the analysis of peakflow impacts in the DEIS results in the underestimation of the extent and intensity of cumulative peakflow alteration within the analysis area under existing conditions and the action alternatives.

Notably, much of the soil compaction caused by grazing occurs in watersheds that have been subjected to past logging and that would be affected by significantly accelerated logging under the action alternatives (DEIS, p. 802). Therefore, soil compaction from grazing is adding to peakflow elevation caused by logging in many watersheds. It will continue to do so, due to the persistence of soil compaction and on-going grazing. Therefore, the DEIS failed to reasonably assess the cumulative impacts on peakflows under the alternatives, because it failed to reasonably assess the contribution of grazing impacts to peakflows in conjunction with the other impacts of action alternatives on peakflows from canopy removal by logging. This defect causes the DEIS to underestimate the cumulative effects of the alternatives on peakflows.

These foregoing defects with respect to grazing, soil compaction, and peakflows, must be rectified in the FEIS. The FEIS must be revamped to ensure that the cumulative impacts of existing conditions and proposed actions on peakflows are analyzed and disclosed, including those from soil compaction caused by grazing.

<u>Peakflow elevation by soil compaction from logging, roads, and landings</u>

Because the DEIS's analysis of peakflows only narrowly focuses on forest canopy conditions, it fails to reasonably analyze and disclose the effects of soil compaction caused by roads, landings and logging. Roads persistently reduce infiltration rates by about 95-99% (Luce, 1997). Due to the extremely low infiltration rates on roads, they generate surface erosion and runoff in response to frequent, low-intensity rainfall and snowmelt events, for as long as the road exists, resulting in persistent and chronic elevation of surface runoff and peakflows. A copious amount of the

analysis area is occupied by roads. Although it has long been well-documented that the soil impacts of roads significantly alter runoff and peakflows, the DEIS's analysis of peakflows fails to reasonably disclose the existing effects of roads on peakflows.

Per unit area affected, landings have impacts on soil compaction and infiltration processes that are akin to those from roads in their severity and persistence, as other USFS cumulative methods have acknowledged (Menning et al., 1996). The DEIS analysis fails to reasonably incorporate the cumulative effects of the area affected by landings into the analysis of existing effects on peakflows.

The DEIS fails to disclose the amount of area occupied by existing landings, although the DEIS (p. 795) indicates that this affected area is known.[1] This is a considerable defect because a significant amount of the analysis area has been affected landings due to their association with logging.

Despite the lack of disclosure, the existing extent of compaction from landings can be estimated from the association of landings with logged areas. According to the DEIS, about 46% of the forested area on BLM lands within the planning area has a "management history." It can be reasonably assumed that this management history is from logging. Landings typically occupy about 1-2% of the area logged. Therefore, based on the foregoing, more than 15,000 acres of BLM lands within the analysis area have been severely compacted by landings. This estimated area of landings is roughly equivalent to the area of more than 4,100 miles of road with a mean width of 30 feet. This is clearly significant due to intensity of compaction in areas affected by landings and its effects on runoff and peakflows.

It is highly likely that a significant amount of the soils compacted by landings occur in relatively close proximity to streams. In the recent past, the flatter areas along streams were targeted for landing construction. Landings are usually proximate to ground-based logging operations. The DEIS (p. 345) concedes that many of the riparian areas within the planning area have been logged at least once, so it is likely that a significant amount of these riparian areas have compacted soils from landing construction and use, which contributes to elevated peakflows.

Based on the foregoing, it is obvious that compaction from landings has been significant and is contributing to increases in peakflows. Therefore, the DEIS's failure to reasonably analyze and disclose these effects on peakflows causes the DEIS to underestimate the magnitude and extent of peakflow elevation under existing conditions and the alternatives.

The DEIS (p. 426) concedes that compaction from past logging persists on BLM lands throughout the analysis area. However, the DEIS includes no estimate of the amount of this compaction. Although the DEIS (p. 426) asserts that the amount of compaction from past logging is "not known," it can be reasonably estimated from the information in the DEIS. The DEIS (p. 206) notes that about 42% of the 2.2 million acres of BLM forested lands within the

---

[1] It appears that the area occupied by landings is known because the DEIS (p. 795) states "The net effect of road building versus road decommissioning results in a less than 1% increase over current road and landing acreage in Alternatives 2 and 3 and a net decrease in acres in the No Action Alternative and Alternative 1." This statement requires knowledge of current landing acreage, although the total amount is not disclosed in the DEIS.

analysis consists of young stands. These young stands have been logged. It is likely that the majority of this previously logged area was logged by ground-based methods, with some limited amount of skyline. The DEIS (p. 795) estimates that ground-based and skyline logging methods, respectively, cause detrimental soil disturbance on about 15% and 3% of the area logged. Therefore, it is reasonable to assume that about 10%, or more than 92,000 acres, of the previously logged young stands on BLM lands within the analysis have compacted soils, which cumulatively contribute to increases in peakflows.

It is likely that a significant amount of riparian areas have soils compacted by previous logging. The DEIS (p. 345) concedes that many of the riparian areas within the planning area have been logged at least once. Compacted soils in riparian areas are especially likely to increase peakflows due to their proximity to streams, although this is not disclosed in the DEIS.

The foregoing clearly indicates that the DEIS failed to reasonably disclose the likely extent and distribution of existing cumulative soil compaction caused by roads, landings, and past logging. The DEIS also failed to reasonably evaluate and disclose that the soil compaction from these activities contributes to elevated peakflows. Notably, there are numerous tractable methods for estimating the changes in peakflows due the effects of soil compaction.

The DEIS exacerbates these deficiencies by failing to reasonably disclose that the methods used to provide its flawed estimates of management activities on peakflows do not adequately address the effects of soil compaction on peakflows. These manifold defects cause the DEIS to underestimate the extent and magnitude of peakflow alteration under existing conditions and under the activities proposed under the alternatives. In so doing, the DEIS has failed to reasonably disclose cumulative impacts on peakflows and to differentiate among the alternatives with respect to effects on peakflows.

Peakflow elevation due to accelerated topsoil loss from management activities

The DEIS also failed to factor the effects of topsoil loss into the analysis of existing impacts of management on peakflows. This is significant because there are numerous activities and conditions in the analysis area that have cumulatively accelerated topsoil erosion over many decades, including logging, landings, roads, and grazing.

The effects of topsoil loss on the hydrologic properties of soils and streamflows are not trivial. The loss of one inch of soil over one square mile results in the loss of more than 813,120 cubic feet of available water storage in the soil profile. This loss of the ability of soils to store water contributes to peakflow elevation. Topsoil loss also contributes to peakflow elevation by reducing infiltration rates, because the uppermost soil profiles typically have the highest infiltration rates (Maidment, 1993).

Peakflow elevation from the alteration of runoff routing by roads

The DEIS fails to credibly and reasonably analyze the multiple effects of roads on peakflows. This is significant because roads increase peakflows in several ways besides compaction. Roadcuts intercept subsurface flow (Megahan, 1972), converting it to surface flow during wetter

periods. The physics of water flow in soils make the interception of subsurface flows by roadcuts unavoidable (Kirkby, 1978).

Roads also contribute to elevating peakflows by concentrating runoff and rapidly shunting it to streams via integration with the stream network (Wemple et al., 1996; Jones and Grant, 1996; Bowling et al., 2000; USFS, 2000; Gucsinski et al., 2001; La Marche and Lettenmaier, 2001; MacDonald and Coe, 2007). The effects of roads alone on peakflows in the Pacific Northwest are estimated to be roughly equivalent to the effect of logging (La Marche and Lettenmaier, 2001). These impacts of roads on peakflows are in addition to those caused by loss of vegetation from logging (La Marche and Lettenmaier, 2001). USFS et al. (1993) noted that much of the massive road network in the area under the aegis of the Northwest Forest Plan, including the DEIS planning area, already had adversely affected peakflows.

It is also well established that a significant fraction of road networks is hydrologically connected to channel networks, elevating peakflows (Wemple et al., 1996; Jones and Grant, 1996; Bowling et al., 2000; La Marche and Lettenmaier, 2001; Gucinski et al., 2001). This is certainly the case for roads on BLM lands within the analysis area, although this is not adequately disclosed in the DEIS. A copious amount of roads occur within 200 feet of stream channels (DEIS, p. 377). Notably, it is likely that roads that are more than 200 feet from streams contribute accelerated runoff to streams. The Clearwater National Forest (2003) noted that roads within 300 feet of streams were likely to have some hydrologic connectivity with streams.

However, the DEIS's analysis of impacts on peakflows does not adequately incorporate and disclose these effects of roads on peakflow elevation. Notably, this failure in the DEIS exists despite the DEIS's (p. 388) acknowledgment that roads elevate peakflows by altering runoff pathways.

Estimating road impacts on peakflows is tractable because models for road impacts, including runoff routing, on peakflows have been developed (La Marche and Lettenmaier, 2001). The DEIS also fails to adequately disclose that its analysis omits these significant effects on peakflows, and therefore underestimates peakflow elevation under existing conditions and the alternatives. Because available scientific information indicates that the effects of roads on peakflows is equivalent and in addition to the impacts of vegetation removal by logging (e.g., Bowling et al., 2000; La Marche and Lettenmaier, 2000), it is likely that actual peakflow alteration within the DEIS analysis area is about double the level estimated by the DEIS's defective analysis. This is an extremely significant magnitude of underestimation.

<u>Peakflow elevation by wildfire</u>

Sporadic wildland fire occurs with some frequency within the analysis area (DEIS, Fig. 19, p. 195). Although patchy, wildland fire can affect a significant fraction of a watershed, and, especially, of smaller watersheds that exhibit the most pronounced increases in peakflows in response to vegetation removal. Depending on the fire severity and extent, the impacts on vegetation and soils can significantly elevate peakflows, as repeatedly documented in many studies.

Although wildfire occurrence at a given location cannot be predicted, the extent of future fire-affected areas over a larger analysis area, such as the DEIS analysis area, can be estimated based on past fire occurrence. However, the DEIS's analysis of effects on peakflows did not take into account the effects of wildfire occurrence. Therefore, the DEIS failed to reasonably analyze and disclose the cumulative effects of the alternatives' direct impacts on peakflows, together with those from wildfire processes.

V. THE DEIS'S ANALYSIS OF THE IMPACTS ON PEAKFLOWS ARE BASED ON ASSUMPTIONS THAT ARE NOT WARRANTED AND FRAUGHT WITH POTENTIAL ERROR, WHICH ARE NOT ADEQUATELY DISCLOSED

The methods used in the DEIS to screen for peakflow impacts rely on assumptions that are not warranted based on available scientific information. For instance, in rain-dominated watersheds, the DEIS arbitrarily assumes that only areas that have less than 30% canopy cover contribute to elevated peakflows. One of the mechanisms by which logging contributes to elevated peakflows is through the reduction of evapotranspiration (ET) and precipitation interception from tree removal (Reid, 1993; Rhodes et al., 1994). Reductions in canopy cover that retain above 30% canopy cover still contribute to elevation in peakflows by decreasing interception and ET. Therefore, the DEIS clearly fails to disclose the lack of scientific basis for its unfounded assumptions.

Similarly, the DEIS's analysis of peakflows assumes that only estimated increases in peakflows in excess of an arbitrary threshold are significant (DEIS, pp. 382-388, 1095-1105). The DEIS fails to examine the veracity of these assumed thresholds based on available scientific information. The DEIS also fails to reasonably disclose scientific information salient to the assumed thresholds of peakflow elevation in the DEIS. For instance, the DEIS does not disclose that Dunne et al. (2001) expressly noted that it cannot be reasonably assumed that relatively small increases in peakflows do not have significant adverse impacts on stream systems, aquatic habitats, and fish populations, because relatively small increases in peakflows exponentially increase sediment transport.

There are many more embedded assumptions in the DEIS analysis of peakflow elevation that cumulatively and synergistically influence the veracity of the results, but are not reasonably assessed for their veracity within the context of available scientific information. These include assumed differences in snowpack accumulation and snowmelt rates as a function of canopy conditions, assumed flow-frequency relationships in the watersheds analyzed, and the size of storms analyzed. Although these assumptions strongly influence the veracity of the results and are fraught with sources of cumulative error, the DEIS fails to reasonably assess and disclose how tenable the assumed approximations are, based on available scientific information.

The DEIS compounds these defects by failing to reasonably assess and disclose the likely magnitude of error inherent in these individual assumed approximations, in order to estimate the total potential error inherent in the DEIS's aggregate estimates of the magnitude and extent of peakflow elevation in the analysis area. Such an analysis is essential to reasonably disclosing the potential accuracy of the results and has long been a standard part of hydrologic analyses (Maidment, 1993). This is a very significant defect, because it is highly likely that the aggregate

potential for error in the DEIS's analysis of impacts on peakflow elevation is relatively large, which undermines the veracity of the DEIS's analysis.

The foregoing aspects of the DEIS's analysis of peakflow elevation are central to assessing its potential accuracy. Therefore, these existing defects must be rectified. The FEIS must reasonably examine the veracity of the assumptions in the analysis, based on available scientific information. The FEIS must also credibly assess and disclose the potential for error in the analysis.

VI. THE DEIS's USE OF ASSUMED THRESHOLDS OF DETECTABILITY IS NOT ADEQUATE TO REASONABLY DISCLOSE THE LIKELY IMPACTS OF MANAGEMENT-INDUCED PEAKFLOW ELEVATION

The DEIS's discussion of variability, detectability, and significance of peakflow elevation compounds the defects in its analysis of peakflow impacts. For instance, the DEIS asserts that variability in streamflow is likely greater than the level of elevation of peakflows caused by management activities. However, currently-increased peakflows caused by existing conditions are *in addition* to natural variation. This overlaying of management-induced impacts causes altered peakflows to be outside of the range of natural variability. Management-induced peakflow elevation increases not only the magnitude of peakflows, but also the frequency of peakflows of a given magnitude. Because sediment transport is strongly affected by peakflows, peakflow elevation inexorably increases channel erosion and sediment transport. This occurs whether or not peakflow increases are less than the detection thresholds assumed in the DEIS.

Similarly, the DEIS erroneously conflates detectability with ecological significance. Impacts that are manifest at levels below the threshold of detectability of peakflow increases can have nonetheless profound ecological impacts. Dunne et al. (2001) noted that peakflow elevation by logging and roads is an important concern because even minor changes in peakflow magnitude and frequency can have major effects on salmonids by triggering significant changes in channel erosion and sediment transport.

Further, the DEIS failed to reasonably assess the considerable error associated with its methods for analyzing peakflow impacts. It is highly likely that if the error in the methods were reasonably assessed, that it would more than bound the DEIS's assumed thresholds of detectability in peakflow increases. Thus, the magnitude of the expected error in the DEIS's peakflow analyses is likely so large that if expected errors are taken into account, peakflows may be elevated beyond the assumed detection limits, even where the DEIS's methods – which disclose only the point estimates – indicate that peakflows have not been elevated. This is undisclosed in the DEIS.

VII. THE DEIS'S FAILURE TO REASONABLY DISCLOSE IMPACTS ON PEAKFLOWS RENDERS THE ANALYSIS OF AQUATIC IMPACTS INADEQUATE.

These DEIS defects regarding peakflows are significant because peakflow strongly affects a host of aquatic resources and processes, including sediment delivery to streams, sediment transport, channel form, turbidity, fish habitat conditions, levels of fine sediment in streams, and

downstream flooding (USFS et al., 1993; Reid, 1993; Wissmar et al., 1994; Spence et al., 1996), although this is inadequately disclosed in the DEIS.

For instance, the DEIS fails to adequately disclose that increased peakflows inexorably widen stream channels as documented by Dose and Roper (1994) in logged watersheds in Oregon. The DEIS also inadequately considers and discloses that it is well-known that increases in stream channel width increase water temperature, even in the absence of shade loss (Rhodes et al., 1994; McCullough, 1999; Bartholow, 2000). This is a significant defect because elevated summer water temperatures are already a widespread aquatic problem within the analysis area.

Increased peakflows increase the downstream transport of sediment from headwater streams to the depositional reaches where salmonids spawn and rear. They can also trigger elevated channel erosion, adding to downstream sedimentation. Due to their characteristics, some headwater streams are extremely vulnerable to channel erosion caused by peakflow elevation (Rosgen, 1996); due to their position in the channel network, this increases downstream sedimentation in fish habitats (Montgomery and Buffington, 1998). Increases in peakflows can also increase bedload movement, reducing the survival of salmonid eggs and alevins (USFS et al., 1993). Peakflow elevation by logging and roads is an important concern because even minor changes in peakflow magnitude and frequency can have major effects on salmonids by triggering significant changes in channel erosion and sediment transport (Dunne et al., 2001).

Turbidity and temperature, which are both strongly affected by peakflow elevation, are set as state water quality standards. Because the DEIS failed to reasonably estimate cumulative elevation of peakflows and its resulting effects on turbidity and water temperature, the DEIS's analysis of compliance with water quality standards is defective.

These impacts all strongly affect the survival and production of salmonids that inhabit the streams that will be affected by action alternatives (Meehan, 1991; USFS et al., 1993). Therefore, it is clear that DEIS's analysis of the action alternatives' impacts on ESA listed fish, fish habitat condition, and fish populations is fatally flawed because it fails to adequately incorporate the likely cumulative impacts of the action alternatives on peakflows and resulting aquatic impacts.

VIII. THE DEIS FAILS TO REASONABLY ANALYZE AND DISCLOSE THE CUMULATIVE IMPACTS ON LOW FLOWS AND SUBSURFACE-SURFACE WATER INTERACTIONS

There are numerous activities and conditions in the analysis area that individually and cumulatively reduce low flows and negatively impact subsurface-surface water interactions. These include: soil compaction and loss caused by grazing, roads, landings, and logging; alterations in surface and subsurface runoff timing and routing caused by roads; and impacts on riparian vegetation and channel form caused by grazing. Reductions in low flows are a critically important factor for native aquatic species (e.g., salmonids, amphibians) survival and persistence. The DEIS fails to reasonably analyze and disclose these impacts.

Soil compaction contributes to reduced low flows by increasing surface runoff and reducing the amount of water stored in the soil profile that can supply streamflow during low flow periods. Grazing, roads, logging, and landings all increase soil compaction and surface runoff, while decreasing available water storage in the soil profile which can ultimately be supplied to streams during low flow periods. Kauffman et al. (2004) documented that grazing caused a significant loss of water storage capacity in soils due to compaction.

However, both grazing and roads have additional impacts on runoff that can reduce low flows. Grazing often causes channel incisement, which lowers near-stream water tables and xerifies adjacent riparian areas (Platts, 1991; Rhodes et al., 1994; CWWR, 1996; USFS, 2001). These impacts reduce low flows. Beschta et al. (1991) and Beschta et al. (1993) noted that channel incisement caused by grazing likely reduced baseflow contributions during the low flow period in damaged riparian systems. The cumulative effects of grazing in damaged systems have likely caused some perennial streams to become intermittent (Rhodes et al., 1994). Elimination of grazing in damaged riparian areas has increased low flows (Meehan, 1991), indicating that grazing had decreased low flows.

Seasonally-saturated and perennially-saturated riparian areas, wetlands and springs are vital to the maintenance of low flows and summer water temperatures. Many of these features are particularly vulnerable to significant damage from livestock grazing and, hence, are often pervasively degraded in areas subjected to livestock grazing (Rhodes et al., 1994; CWWR, 1996; USFS, 2001), contributing to the loss of instream flows derived from subsurface flows.

Reductions in subsurface flow caused by soil compaction, and subsurface flow interruption by roads, grazing, and logging negatively affect aquatic resources, because these impacts reduce hyporheic flows (Hancock, 2002). Cumulative soil compaction contributes to reductions in low flows (Booth et al., 2002; Booth et al., 2004).

Roadcuts inexorably interrupt subsurface flow. This interception by roadcuts is likely to reduce downslope soil moisture levels and subsurface flow contributions to affected streams, contributing to reduced baseflows (Tague and Band, 2001). Hancock (2002) noted that logging and roads reduced subsurface flows to hyporheic areas by reducing subsurface percolation and baseflow contributions to streams. Hicks et al. (1991) documented that watersheds in Oregon had statistically significant decreases in low flows after logging relative to an unlogged control watershed.

Subsurface flow interception by roads likely increases water temperatures via a two-pronged effect. Reductions in subsurface flows to streams reduce low flow volumes, which, alone, increase summer water temperatures (Beschta et al., 1987; Rhodes et al., 1994). However, subsurface flows entering streams are also typically far cooler than surface flows, aiding in the thermal regulation of streams during low flows (Beschta et al., 1987), so the loss of this cooler water also contributes to stream warming (Rhodes et al., 1994). Increases in summer water temperature have negative effects on native salmonids. Elevated summer water temperature is a widespread water-quality problem afflicting salmonids within the analysis area (USFS et al., 1993; DEIS, p. 359).

Despite this information, the DEIS is without any adequate analysis and disclosure of cumulative impacts of management activities on low flows and surface-subsurface hydrologic interactions. The

defects on this front must be rectified in the FEIS by fully analyzing and disclosing all of the impacts of existing conditions and proposed activities on low flows and subsurface-surface water interactions based on available scientific information.

IX. CONCLUSION

The DEIS's analysis of the cumulative impacts of management activities on the stability of streamflow is wholly inadequate due to several significant defects. The DEIS failed to reasonably examine and disclose the impacts of logging on peakflows generated by rain-on-snow based on information in Grant et al. (2007) on the level of forest removal that generates detectable increases in these peakflows.

The DEIS also failed to examine the peakflow impacts of existing conditions and the proposed activities in the action alternatives on smaller watersheds where peakflow increases are likely to be most commonly manifest and pronounced. The DEIS failed to analyze and disclose the impacts of roads, cumulative soil compaction, topsoil loss, and future wildfire on peakflows. These foregoing defects cause the DEIS to significantly underestimate the magnitude and prevalence of peakflow elevation across the analysis area under existing conditions and the action alternatives. They also cause the DEIS to underestimate the differences among the alternatives with respect to peakflows. The DEIS compounds the foregoing defects by failing to reasonably disclose the limitations of its analysis and its likely degree of error. These are serious failures, because the action alternatives will significantly increase impacts on peakflows with negative consequences for aquatic conditions and salmonid populations. For these combined reasons, the DEIS has failed to adequately disclose the aquatic impacts of the alternatives.

Existing conditions and activities under the alternatives also affect low flows and surface-subsurface water interactions. These impacts affect aquatic conditions, but are not disclosed in the DEIS.

For these reasons, the DEIS fails to adequately disclose the cumulative impacts on streamflow stability. It also fails to adequately differentiate among the alternatives with respect to impacts on streamflow stability and its aquatic consequences.

Literature cited

Bartholow, J.M., 2000, Estimating cumulative effects of clearcutting on stream temperatures, Rivers, 7: 284-297.

Beschta, R.L., Bilby, R.E., Brown, G.W., Holtby, L.B., and Hofstra, T.D., 1987. Stream temperature and aquatic habitat: Fisheries and forestry interactions. Streamside Management: Forestry and Fishery Interactions, pp. 191-231, Univ. of Wash. Inst. of Forest Resources Contribution No. 57, Seattle, Wash.

Beschta, R.L., Platts, W.S., and Kauffman, B., 1991. Field Review of Fish Habitat Improvement Projects in the Grande Ronde and John Day River Basins of Eastern Oregon. BPA Project No. 91-069, Bonneville Power Admin., Div. of Fish and Wildlife, Portland, Or.

Beschta, R.L., Griffith, J., and Wesche, T.A., 1993 Field Review of Fish Habitat Improvement Projects in Central Idaho. BPA Project No. 84-24; 83-359, Bonneville Power Admin., Div. of Fish and Wildlife, Portland, Or.

Beschta, R.L., Rhodes, J.J., Kauffman, J.B., Gresswell, R.E, Minshall, G.W., Karr, J.R, Perry, D.A., Hauer, F.R., and Frissell, C.A., 2004. Postfire Management on Forested Public Lands of the Western USA. Cons. Bio., 18: 957-967.

Booth, D.B., Hartley, D., and Jackson, R., 2002. Forest cover, impervious-surface area, and mitigation of stormwater impacts. J. Amer. Water Resour. Assoc., 38:835-845.

Booth, D.B., Karr, J.R., Schauman, S., Konrad, K.P., Morley, S.A., Larson, M.G., and Burges, S.J., 2004. Reviving urban streams: Land use, hydrology, biology, and human behavior. J.Amer. Water Resources Assoc. 40:1351-1364.

Bowling, L.C., Storck, P., and Lettenmaier, D.P., 2000. Hydrologic effects of logging in western Washington, United States. Water Resour. Res. 36: 3223-3240.

Clearwater National Forest (CNF), 2003. Roads analysis report. Clearwater National Forest, Orofino, ID.

Cowley, E.R., 2002. Monitoring Current Year Streambank Alteration. US Bureau of Land Management, Idaho State Office, Boise, ID.

(CWWR) Centers for Water and Wildland Resources, 1996. Sierra Nevada Ecosystem Project Report, Summary and Final Report to Congress, Summary and Vol. I-III. Wildland Resources Center Report No. 39, University of California, Davis.

Dose, J.J. and Roper, B.E. 1994. Long-term changes in low-flow channel widths within the South Umpqua watershed, Oregon. Water Resour. Bull., 30: 993-1000.

Dunne, T., Agee, J., Beissinger, S., Dietrich, W., Gray, D., Power, M., Resh, V., Rodrigues, K., 2001. A scientific basis for the prediction of cumulative watershed effects. University of California Wildland Resource Center Report No. 46.

Grant, G.E., Lewis, S., Swanson, F. and McDonnell, J., 2007 (*In Review*). Effects of forest practices on peak flows and consequent channel response in western Oregon: a state-of-science report. Pacific Northwest Research Station, U.S. Department of Agriculture, U.S. Forest Service. Corvallis, OR.

Gucinski, H., Furniss, M.J, Ziemer, R.R and Brookes, M.H., 2001. Forest roads: a synthesis of scientific information. Gen. Tech. Rep. PNW GTR-509. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Hancock, P.J., 2002. Human impacts on the stream-groundwater exchange zone. Env. Mgmt. 29: 763-781.

Hicks, B.J., Beschta, R.L., and Harr, R.D. 1991. Long-term changes in streamflow following logging in western Oregon and associated fisheries implications. Water Resources Bulletin 27: 217-226.

Jones J.A. and Grant. G.E., 1996. Peak flow responses to clear-cutting and roads in small and large basins, western Cascades, Oregon. Water Resour. Res. 32: 959-974.

Karr, J.R., Rhodes, J.J., Minshall, G.W., Hauer, F.R., Beschta, R.L., Frissell,C.A., and Perry, D.A, 2004. Postfire salvage logging's effects on aquatic ecosystems in the American West. BioScience, 54: 1029-1033.

Kauffman, J.B., Thorpe, A.S., Brookshire, J. and Ellingson, L., 2004. Livestock exclusion and belowground ecosystem responses in riparian meadows of eastern Oregon. Ecological Applications: 1671-1679.

Kirkby, M.J. (ed.), 1978. Hillslope Hydrology. John Wiley & Sons, Inc., New York.

La Marche, J.L. and Lettenmaier, D.P., 2001. Effects of forest roads on flood flows in the Deschutes River, Washington. Earth Surf. Process. Landforms, 26: 115-134.

Luce, C.H., 1997. Effectiveness of road ripping in restoring infiltration capacity of forest roads. Restoration Ecology, 5: 265-270.

MacDonald, A. and Ritland, K.W., 1989. Sediment Dynamics in Type 4 and 5 Waters: A Review and Synthesis. TFW-012-89-002, Wash. Dept. of Natural Resour., Olympia, Wash.

MacDonald, L.H. and Coe, D., 2007. The influence of headwater streams on downstream reaches in forested areas. Forest Sci., 53: 148-168.

Maidment, D.R. (ed), 1993. Handbook of Hydrology. McGraw-Hill, New York.

McCullough, D.A., 1999. A review and synthesis of effects of alterations to the water temperature regime on freshwater life stages of salmonids, with special reference to chinook salmon. USEPA Technical Report EPA 910-R-99-010, USEPA, Seattle, WA.

Meehan, W.R. (ed.). 1991. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Am. Fish. Soc. Special Publication 19.

Megahan, W.F., 1972. Subsurface flow interception by a logging road in mountains of central Idaho. Conf. Proceedings: National Symposium on Watersheds in Transition, pp. 321-329. Am. Water Resour. Assoc., Bethesda, MD.

Menning, K., Erman, D.C., Johnson, K.N., Sessions, J. 1996. Modeling aquatic and riparian systems, assessing cumulative watershed effects and limiting watershed disturbance. Sierra Nevada Ecosystem Project Final Report to Congress, Addendum, pp. 33-52. Wildland

Resources Center Report No. 39, Centers for Water and Wildland Resources, University of California, Davis.

Montgomery, D.R. and J.M. Buffington, 1998. Channel processes, classification, and response. *In* R. Naiman and R. Bilby (eds.), River ecology and management: lessons from the Pacific Coastal Region, p. 13-42. Springer-Verlag, New York.

Murphy, M.L. 1995. Forestry impacts on freshwater habitat of anadromous salmonids in the Pacific Northwest and Alaska – requirements for protection and restoration. NOAA Coastal Oceans Program, Decision Analysis Series No. 7. U.S. Dept. Commerce, NOAA, Coastal Ocean Office, Silver Springs, Maryland.

O'Loughlin, E.M., 1986. Prediction of surface saturation zones in natural catchments by topographic analysis. Water Resour. Res., 22: 794-804.

Platts, W.S., 1991. Livestock grazing. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats, pp. 389-424. Am. Fish. Soc. Special Publ. 19.

Reid, Leslie M., 1993. Research and cumulative watershed effects. Gen. Tech. Rep. PSW-GTR-141. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture.

Rhodes, J.J., McCullough, D.A., and Espinosa Jr., F.A., 1994. A Coarse Screening Process for Evaluation of the Effects of Land Management Activities on Salmon Spawning and Rearing Habitat in ESA Consultations. CRITFC Tech. Rept. 94-4, Portland, Or.

Rosgen, D. 1996. Applied river morphology. Wildland Hydrology, Pagosa Springs, CO

Spence, B.C., Lominicky, G.A., Hughes, R.M., and Novitzki, R.P., 1996. An ecosystem approach to salmonid conservation. TR-4501-96-6057. ManTech Environmental Research Services Corporation, Corvallis, Oregon. (available from the National Marine Fisheries Service, Portland, Oregon).

Tague, C. and Band, L, 2001. Simulating the impact road construction and forest harvesting on hydrologic response. Earth Surf. Process Landforms, 26: 135-151.

USFS, 2000. Roadless Area Conservation Final Environmental Impact Statement. USFS, Washington, D.C.

USFS, 2001. Sierra Nevada Forest Plan Amendment FEIS, USFS PSW Region, San Francisco, Ca.

USFS, NMFS, USBLM, USFWS, USNPS, USEPA, 1993. Forest Ecosystem Management: An Ecological, Economic, and Social Assessment. USFS PNW Region, Portland, Or.

Wemple, B.C., Jones, J.A., Grant, G.E., 1996. Channel network extension by logging roads in two basins, Western Cascades, Oregon. Water Resour. Bull., 32: 1195-1679.

Wissmar, R.C., Smith, J.E., McIntosh, B.A., Li, H.W., Reeves, G.H., Sedell, J.R. 1994. Ecological health of river basins in forested regions of eastern Washington and Oregon. Gen. Tech. Rep. PNW-GTR-326. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Curriculum Vitae: Jonathan J. Rhodes

Hydrologist
P.O. Box 15286 •• Portland, OR 97293-5286

EDUCATION

1989: Doctoral candidacy degree in forest hydrology at the Univ. of Wash. Completed all requirements but dissertation.

1985: M.S. in Hydrology and Hydrogeology at the Univ. of Nev.-Reno. Thesis topic: The influence of seasonal stream runoff patterns on water quality.

1981: B.S. in Hydrology and Water Resources at the Univ. of Ariz.

PROFESSIONAL HISTORY

Sept. 2001 – present. Hydrologist. Main duties: Analysis of effects of water and land use on streams and other aquatic resources, including native salmonids and their habitats; diagnosis of watershed and stream conditions; stream monitoring; development of programmatic and site-specific watershed and stream protection measures; project management. Recent projects (and clients): Analysis of potential effects of groundwater pumping on streamflow (Conf. Tribes of the Umatilla Indian Reservation, OR); diagnosis of watershed and stream conditions in an urbanized watershed (West Multnomah Soil and Water Conservation District, OR); diagnosis of effects of grazing on watershed and stream conditions in forested watersheds in N. California (Center for Biological Diversity (CBD)) and Oregon (Or. Natural Desert Assoc.); Coordinator and Aquatic Scientist for the Western Native Trout Campaign (CBD, Pacific Rivers Council (PRC), Biodiversity Conservation Alliance, Trout Unlimited).

Aug. 1990 – Sept. 2001. Consulting hydrologist for non-profit organizations. Past projects (and clients): hydrologic characterization of remnant marsh proposed as urban wildlife refuge/greenspace (Multnomah County Parks Department, OR); review of aquatic effects of: quarry expansion (Friends of Forest Park, OR), urban construction (homeowners consortium, W. Linn, OR); forest manipulations on streamflow (PRC).

Apr. 1989 – Sept. 2001. Senior Scientist-Hydrologist, Columbia River Inter-Tribal Fish Commission. Main duties: Administration and implementation of projects monitoring channel change from land management in Columbia River basin; development of programmatic and site-specific land management plans to ensure protection of watershed integrity, water quality and aquatic resources; development of restoration plans for watersheds degraded by grazing, roads, logging and mining; design of plans for monitoring watershed and stream erosion, sedimentation, water quality, and habitat conditions; review of land management plans for adequacy of protection of aquatic resources; field evaluation of watershed and channel conditions throughout the Columbia Basin; expert witness testimony; development of technical recommendations for policy staff for protection of natal habitat for anadromous fish; review of state and federal aquatic resource monitoring plans; report and proposal writing; and, participation in various state and federal technical work groups.

Aug. '84 -- Apr. '89. Research assistant, College of Forestry, Univ. of Wash. Main duties: analysis and interpretation of water quality-quantity data; technical report writing; design and maintenance of water chemistry and quantity monitoring network in a coastal forested watershed; training in data acquisition techniques; public presentation of findings.

July -- Oct. 1987 and May -- Oct. 1988. Consulting hydrologist, Tahoe Regional Planning Association, CA and NV. Main duties: field delineation and mapping of riparian zones, wetlands, and erosion-prone areas.

June -- Sept. 1985 and July 1986. Research assistant, Dept. of Geophysics, Univ of Wash. Main duties: operation of field station for glacier research on Mt. Olympus, Wash.; measurement of snow and glacier melt rates; mapping of supra- and extra- glacial streams contributing to basal sub-glacial flow rates on surging and non-surging glaciers in the Alaska Range, Alaska.

Jan. 1984. Consultant with C.M. Skau, Reno, NV. Main duties: field evaluation of logging roads for erosion potential and sedimentation risk; recommendations for placement of future roads to minimize erosion and sediment delivery to fish-bearing streams in coastal Northern California; report preparation.

Oct. 1983 -- June 1984. Hydrologic Tech., USGS, Carson City, NV. Main duties: aid in development and calibration of predictive water quality model for the Truckee River; statistical analysis of water quality data; identification and quantification of non-point sources of nutrients to Truckee River, NV.

Aug. 1981 -- Sept. 1983. Research Assistant, Univ. of Nev.-Reno. Main duties: design and installation of instrument network to monitor water chemistry and quantity in a small, forested alpine watershed in the Sierra Nevada; water quality sampling; data interpretation and management; preparation of reports, grant proposals, and publications, computer programming for data reduction and storage; mapping of geology, soils and runoff-producing areas; and, training of field technicians.

Feb. -- May 1981. Water Quality Intern, Pima Assoc. of Gov'ts., Tucson, AZ. Main duties: water quality sampling of agricultural production wells; mapping of groundwater levels; and, coordination of sampling efforts.

PROFESSIONAL SERVICE

Mar. 2007. Invited Panel Speaker, International Environmental Law Conference: "Fuel Treatments & Thinning: Its Impacts and Low Priority Relative to Other Needed Restoration Measures," Univ. of OR, Eugene, OR.

Mar. 2007. Invited Panel Speaker, International Environmental Law Conference: "The Impacts of Livestock Grazing on Water Quality and Trout Habitats," Univ. of OR, Eugene, OR.

Feb. 2005. Invited Guest Lecturer, Lewis and Clark School of Law course on public lands law: "Postfire Watershed Management on Western Public Lands" Portland, OR.

Mar. 2004. Invited Panel Speaker, International Environmental Law Conference: "Postfire Watershed Restoration," Univ. of OR, Eugene, OR.

April 2002. Invited Speaker, Restoring Public Lands: Reclaiming the Concept of Forest Restoration, "Watersheds and Fisheries: Restoration Needs for Trout Habitats," Univ. of CO, Boulder, CO

Mar 2002. Invited Panel Speaker, International Environmental Law Conference: "Soils Impacts and Effects on Trout Habitat," Univ. of OR, Eugene, OR

Mar. 2001. Invited Panel Speaker, International Environmental Law Conference: "NFMA and Salmon Habitat Protection," Univ. of OR, Eugene, OR.

May 2000. Invited speaker, 5th National Tribal Conf. on Environmental Management: "Federal Land Management's Effects on Critical Habitat for Endangered Salmon," Lincoln City, OR

July 1998-2000. Peer Reviewer for N. Amer. J. Fish for papers related to the sedimentation of fish habitat in response to erosion from land uses and fire.

Feb. 1998. Invited Speaker, Oregon AFS Annual meeting: "Adaptive management: Is it really adaptive?" Sunriver, OR

May 1996-2000. Guest lecturer, Oregon State Univ. graduate course on riparian and wetland ecology, Corvallis, OR

Apr.-May 1996. Peer-reviewer for Proceedings of Forest-Fish Conference: Land Management Affecting Aquatic Ecosystems, Proc. Forest-Fish Conf., May 1-4, 1996, Calgary, Alberta, Canada. Nat. Resour. Can., Can. For. Serv. Nort. For. Cent., Edmonton, Alberta. Inf. Rep. NOR-X-356.

Apr. 1995. Invited speaker, Pacific Rivers Council Workshop on Watershed Analysis and Salvage Logging, Wenatchee, Wash.

Apr. 1995. Invited speaker, Oregon State Univ. Dept of Fisheries and Wildlife Seminar, Corvallis, OR

Apr. 1995. Invited speaker, American Fisheries Society North Pacific International Chapter, Annual Meeting, Vancouver B.C., Can.

Mar. 1995. Invited speaker, American Fisheries Society Idaho Chapter Annual Meeting, Boise, ID.

Nov. 1994. Invited speaker, President's Council on Sustainable Development Workshop, Yakima, WA.

Sept. 1994. Invited speaker, Oregon Water Resources Research Institute Streambank Restoration Conference: "Biological Methods to Stabilize Streambanks--From Theory to Practice," Portland, OR.

Mar.-April, 1994. Peer-reviewer for Henjum et al., 1994. Interim Protection for Late Successional Forests, Fisheries, and Watersheds: National Forests East of The Cascade Crest, Oregon and Washington. The Wildlife Soc., Bethesda, MD.

Jan. 1993-Sept. 1995. Member, Oregon Department of Environmental Quality's (ODEQ) Technical Advisory Committee for Triennial Review of the State Water Temperature Standard.

Mar. 1993. Invited speaker, Northwest Scientific Association Symposium: "Cumulative Effects of Land Management Practices on Anadromous Salmonids," La Grande, OR.

Aug. 1992 -- Sept. 1992. Member, Ad Hoc Consultant Selection Committee for Portland Water Bureau Study of Future Water Supply Needs.

May 1992. Invited Speaker, US Forest Service, Pacific Northwest Region, Regional Workshop on Monitoring Soil and Water Resources, Bend, OR.

May 1992. Invited Speaker, Northern Arizona University, School of Forestry, Graduate Seminar Series, Flagstaff, AZ.

Jan. 1991 -- Mar. 1995. Member, Technical Work Group: Upper Grande Ronde River Anadromous Fish Habitat Protection, Restoration and Monitoring Plan.

Aug. 1989 -- Feb. 1990. Member, Technical Advisory Committee to ODEQ for development of definitions for level of beneficial use impairment by nonpoint sources.

May 1989 -- Jan. 1991. Member, Nonpoint Source Technical Advisory Committee to Idaho Department of Environmental Quality: Coordinated Nonpoint Source Monitoring Program For Idaho.

PUBLICATIONS

Peer-Reviewed:

Rhodes, J.J., C.M. Skau, and W.M. Melgin, 1984. Nitrate-nitrogen flux in a forested watershed -- Lake Tahoe, USA. In: Recent Investigations in the Zone of Aeration, Proc. of Inter. Symp., Munich, West Germany, 1984, P. Udluft, B. Merkel, and K. Prosl (Editors), pp. 671-680.

Rhodes, J.J., 1985. A Reconnaissance of Hydrologic Transport of Nitrate in An Undisturbed Forested Watershed Near Lake Tahoe. M.S. thesis, Univ. of Nev. Reno, 254 pp.

Rhodes, J.J., C.M. Skau, and J.C. Brown, 1985. An areally intensive approach to hydrologic nutrient transport in forested watersheds. In: The Forest-Atmosphere Interaction, B.A. Hutchison and B.B. Hicks (Editors), pp. 255-270.

Rhodes, J.J., C.M. Skau, D. Greenlee, and D.L. Brown, 1985. Quantification of nitrate uptake by riparian forests and wetlands in an undisturbed headwaters watershed. US Forest Service Gen. Tech. Rept. RM-120.

Rhodes, J.J., C.M. Skau, and D. Greenlee, 1986. The role of snowcover on diurnal nitrate concentration patterns in streamflow from a forested watershed in the Sierra Nevada, Nevada, USA. In: Proc. of AWRA Symposium: Cold Regions Hydrology, Fairbanks Alaska, 1986, D.L. Kane (Editor), pp. 157-166.

Rhodes, J.J., R.L. Armstrong, and S.G. Warren, 1987. Mode of formation of "ablation hollows" controlled by dirt content of snow, J. Glaciology, **33**: 135-139.

R.L. Edmonds, T.B. Thomas, and J.J. Rhodes, 1991. Canopy and soil modification of precipitation chemistry in a temperate rain forest. Soil Soc. of Amer. J., **55**: 1685-1693.

Rhodes, J.J., McCullough, D.A., and Espinosa Jr., F.A., 1994. A Coarse Screening Process for Evaluation of the Effects of Land Management Activities on Salmon Spawning and Rearing Habitat in ESA Consultations. CRITFC Tech. Rept. 94-4, Portland, OR

Rhodes, J.J. 1995. A Comparison and Evaluation of Existing Land Management Plans Affecting Spawning and Rearing Habitat of Snake River Basin Salmon Species Listed Under the Endangered Species Act. CRITFC Tech. Rept. 95-4, Portland, OR

Rhodes, J.J. 1996. Description and Evaluation of Some Available Models for Estimating the Effects of Land Management Plans on Sediment Delivery, Channel Substrate, and Water Temperature, CRITFC, Portland, OR

Espinosa, F.A., Rhodes, J.J., and McCullough, D. A. 1997. The failure of existing plans to protect salmon habitat on the Clearwater National Forest in Idaho. J. Env. Management **49**: 205-230.

Rhodes, J.J., and Purser, M.D., 1998. Overwinter sedimentation of clean gravels in simulated redds in the upper Grande Ronde River and nearby streams in northeastern Oregon, USA: Implications for the survival of threatened spring chinook salmon, Forest-Fish Conference: Land Management Affecting Aquatic Ecosystems, Proc. Forest-Fish Conf., May 1-4, 1996, Calgary, Alberta, Canada. Nat. Resour. Can., Can. For. Serv. Nort. For. Cent., Edmonton, Alberta. Inf. Rep. NOR-X-356, pp: 403-412.

Beschta, R.L., Rhodes, J.J., Kauffman, J.B., Gresswell, R.E, Minshall, G.W., Karr, J.R, Perry, D.A., Hauer, F.R., and Frissell, C.A., 2004. Postfire Management on Forested Public Lands of the Western USA. Cons. Bio., 18: 957-967.

Karr, J.R., Rhodes, J.J., Minshall, G.W., Hauer, F.R., Beschta, R.L., Frissell,C.A. Perry, D.A, 2004. Postfire Salvage Logging's Effects on Aquatic Ecosystems in the American West. BioScience, 54: 1029-1033.

Rhodes, J.J. and Odion, D.C., 2004. Comment Letter: Evaluation of the Efficacy of Forest Manipulations Still Needed. BioScience, 54: 980.

Rhodes, J.J., 2005. Comment on "Modeling of the interactions between forest vegetation, disturbances, and sediment yields" by Erkan Istanbulluoglu et al. J. Geophys. Res. Earth Surf., Vol. 110, No. F1, F01012 10.1029/2004JF000240

Rhodes, J.J. and Baker, W.L., *In press*. Fire probability, fuel treatment effectiveness and ecological tradeoffs in western U.S. public forests. Open Forest Science Journal.

Technical Reports:

1986. Annual Report on Watershed Studies at Olympic National Park. College of Forestry, Univ. of Wash., Seattle, Wash. (Co-authors: R.L. Edmonds, T.B. Thomas, T.W. Cundy)

1987. Annual Report on Watershed Studies at Olympic National Park. College of Forestry, Univ. of Wash., Seattle, Wash. (Co-authors: R.L. Edmonds, T.B. Thomas, T.W. Cundy)

1988. Annual Report on Watershed Studies at Olympic National Park. College of Forestry, Univ. of Wash., Seattle, Wash. (Co-authors: R.L. Edmonds, T.B. Thomas, T.W. Cundy)

1989. Annual Report on Watershed Studies at Olympic National Park. College of Forestry, Univ. of Wash., Seattle, Wash. (Co-authors: R.L. Edmonds, T.B. Thomas, T.W. Cundy)

1990. Coordinated Nonpoint Source Monitoring Program For Idaho. Idaho Dept. of Environmental Quality, Boise, Idaho. (Co-authors: B. Clark, D. McGreer, W. Reid, T. Burton, W. Low, I. Urnovitz, D. McCullough, T. Litke)

1992. The Upper Grande Ronde River Anadromous Fish Habitat Protection, Restoration and Monitoring Plan. Wallowa-Whitman National Forest, Baker, OR (Co-authors: M. Purser, P. Boehne, R.E. Gill, R.L. Beschta, J.R. Sedell, B. McIntosh, J. Zakel, J.W. Anderson, D. Bryson, S. Howes, R. George).

1992. Salmon Recovery Program for the Columbia River Basin: An Advisory Report for the US Congress, Col. Riv. Inter-Tribal Fish Comm., Portland, OR (Co-authors: P.R. Mundy, D.A. McCullough, M.L. Cuenco, T.W. Backman, D.Dompier, P. O'Toole, S. Whitman, E. Larson, B. Watson, G. James).

1993. A comprehensive approach to restoring habitat conditions needed to protect threatened salmon species in a severely degraded river--The Upper Grande Ronde River Anadromous Fish Habitat Protection, Restoration and Monitoring Plan. USFS Gen. Tech. Rept RM-226, pp. 175-179. (Co-authors: J.W. Anderson, R.L. Beschta, P. Boehne, D. Bryson, R.E. Gill, S. Howes, B. McIntosh, M.D. Purser and J. Zakel).

1993. Dante's Video Guide to Habitat Conditions for Wild Spring Chinook Salmon, Steelhead and Bull Trout in the John Day Basin, Oregon. (Video) Presented at AFS National Meeting, Portland, Or, Aug. 29-31. (Co-authors: R. Taylor and M. Purser).

1995. Wildfire and Salvage Logging: Recommendations for Ecologically Sound Post-Fire Salvage Logging and Other Post-Fire Treatments on Federal Lands in the West. Pacific Rivers Council, Portland, OR (Co-authors: R. Beschta, C. Frissell, R. Gresswell, R. Hauer, J. Karr, G. Minshall, D. Perry).

1998. Adaptive management: Is it really adaptive? Abstracts: Oregon AFS Annual Meeting, Feb. 11-13, 1998, p. 31.

1998. Thinning For Increased Water Yield in the Sierra Nevada: Free Lunch or Pie in the Sky? Pacific Rivers Council, Eugene, OR. (Co-author: M. Purser)

1999. Annual Project Report: Watershed Evaluation and Aquatic Habitat Response to Recent Storms. Bonneville Power Administration (BPA), Portland, OR. (Co-author: C. Huntington)

1999. Annual Project Report: Monitoring Fine Sediment in Salmon Habitat in John Day and Grande Ronde Rivers. BPA, Portland, OR (Co-author: M. Purser)

2000. Annual Project Report: Watershed Evaluation and Aquatic Habitat Response to Recent Storms. BPA,. Portland, OR. (Co-author: C. Huntington)

2000. Annual Project Report: Monitoring Fine Sediment in Salmon Habitat in John Day and Grande Ronde Rivers. (Co-author: M. J. Greene)

2001. Annual Project Report: Monitoring Fine Sediment in Salmon Habitat in John Day and Grande Ronde Rivers. BPA, Portland, OR. (Co-author: M. J. Greene)

2001. Imperiled Western Trout and the Importance of Roadless Areas. Western Native Trout Campaign, Center for Biological Diversity, Tucson, Az. (Co-authors: J. Kessler, C. Bradley, and J. Wood)

2002. Tryon Creek Watershed: Overview of Existing Conditions, Data Gaps, and Recommendations for the Protection and Restoration of Aquatic Resources. West Multnomah Soil and Water Conservation District, Portland, OR

2002. An Analysis of Trout and Salmon Status and Conservation Values of Potential Wilderness Candidates in Idaho and Eastern Washington. Western Native Trout Campaign, Center for Biological Diversity, Tucson, AZ. (Co-authors: C. Bradley, J. Kessler, C. Frissell)

2003. Stream and Fish Habitat Conditions in Tryon Creek: Their Likely Causes and Ramifications for Salmonids. Proceedings of Urban Ecology and Conservation Symposium, January 24, 2003, Portland, OR. Portland State University, Environmental Sciences and Resources, Portland, OR

Semi-Technical Publications:

1993. Dam the analysis--heal streams instead. The Assoc. of Forest Service Employees for Env. Ethics Inner Voice, 5(6): 1, 4-5.

1994. Invited Preface to Northwest Science Special Issue--Environmental History of River Basins in Eastern Oregon and Washington. Northwest Sci., **68**.

PROJECT MANAGEMENT

1993-1996. Technical Assistance Contract with NMFS to produce technical guidance for ESA consultations for effects of land management on critical habitat for listed Columbia basin salmon. Main duties: Co-Primary Investigator; primary author of peer-reviewed reports including proposed ESA consultation guidelines for effects on salmon habitat (Rhodes et al., 1994), evaluation and comparison of compatibility of land management plans with protection of critical salmon habitat (Rhodes, 1995), and evaluation of models for estimating land management effects on salmon habitat (Rhodes, 1996); review and synthesis of available scientific literature; budget preparation and tracking; coordination with subcontractors and grantor representatives. Total budget: $230,000

1998-2000. Watershed Evaluation and Aquatic Habitat Response to Recent Storms. Main duties: Primary Investigator; design and implementation of monitoring methods (erosion, runoff, etc.), coordination with subcontracting fish biologists in 10 watersheds with differing levels of grazing and logging within 3 subbasins in Idaho, Washington, and Oregon; training of field technicians; data analysis and synthesis; subcontract administration; proposal development; technical and progress report preparation; budget development and tracking; coordination with grantor representatives. Total budget: $164,000.

1998-2000. Evaluation of Effects of Grazing on Rate of Salmon Habitat Recovery. Main duties: Primary Investigator; design and implementation of monitoring methods, training of field technician; data analysis and synthesis; proposal development; preparation of progress reports; budget development and tracking; coordination with grantor representatives. Total budget: $73,000

1998-2001. Monitoring Fine Sediment Levels in Salmon Habitat in Grande Ronde and John Day Rivers. Main duties: Primary Investigator; design and implementation of methods for monitoring fine sediment levels in four rivers; field technician training; data analysis and synthesis; subcontract administration; proposal development; progress and technical report preparation; budget development and tracking; coordination with grantor representatives. Total budget: $128,000.

2001-2002. Western Native Trout Campaign, Aquatic Scientist and Coordinator. Main duties: Provide oversight and assure scientific integrity of all reports and work products; coordinate conservation efforts among campaign member organizations; coordinate campaign efforts with other groups working to protect and restore trout habitats and populations; budget tracking; technical and progress report preparation.

HONORS AND AWARDS

1996. Leadership and Excellence. Col. River Inter-Tribal Fish Comm., Portland, OR

1991. Employee of the Year. Col. River Inter-Tribal Fish Comm., Portland, OR

1984. Academic Recruitment Scholarship for Outstanding Graduate Prospect. Univ. of Wash, Seattle, Wash.

1982. Maxey Award for Outstanding Graduate Student Paper in Hydrology. Univ. of Nev.-Reno.

1980. Winslow and Myron Reuben Scholarship for Outstanding Undergraduate in the Earth Sciences. Univ. of Ariz., Tucson, Az.

ADDITIONAL TRAINING

1993. USFWS Water Temperature Modeling via SNTEMP

1991. USFWS Introduction to IFIM Investigations