

## **APPENDIX A**

### **MOJAVE-SOUTHERN GREAT BASIN AREA**

#### **PREAMBLE**

The Standards and Guidelines for grazing administration on BLM lands in southern Nevada apply to livestock grazing. The Mojave-Southern Great Basin Resource Advisory Council (RAC) intends that the Standards and Guidelines will result in a balance of sustainable development and multiple use along with progress, over time, toward attaining desired rangeland conditions. Standards are expressions of physical and biological conditions required for sustaining rangelands for multiple uses. Guidelines point to management actions related to livestock grazing for achieving the Standards. Guidelines are options that move rangeland conditions toward the multiple use Standards. Guidelines are based on science, best rangeland management practices, and public input. Guidelines indicate the types of grazing methods and practices for achieving the Standards for multiple use, are developed for functional watersheds and implemented at the allotment level.

The Mojave-Southern Great Basin Resource Advisory Council recognizes that it will sometimes be a long-term process to restore rangelands to proper functioning condition. In some areas, it may take many years to achieve healthy rangelands.

The Resource Advisory Council may be requested by any party to assist reaching agreement in resolving disputes.

#### **STANDARDS AND GUIDELINES**

##### **STANDARD 1. SOILS:**

Watershed soils and stream banks should have adequate stability to resist accelerated erosion, maintain soil productivity, and sustain the hydrologic cycle.

Soil indicators:

- Ground cover (vegetation, litter, rock, bare ground);
- Surfaces (e.g., biological crusts, pavement); and
- Compaction/infiltration.

Riparian soil indicators:

- Stream bank stability.

All of the above indicators are appropriate to the potential of the ecological site.

#### GUIDELINES:

- 1.1 Upland management practices should maintain or promote adequate vegetative ground cover to achieve the Standards.
- 1.2 Riparian-wetland management practices should maintain or promote sufficient residual vegetation to maintain, improve, or restore functions such as stream flow energy dissipation, sediment capture, groundwater recharge, and streambank stability.
- 1.3 When proper grazing practices alone are not likely to restore areas, land management practices may be designed and implemented where appropriate.
- 1.4 Rangeland management practices should address improvement beyond this Standard, significant progress toward achieving Standards, time necessary for recovery, and time necessary for predicting trends.

#### STANDARD 2. ECOSYSTEM COMPONENTS:

Watersheds should possess the necessary ecological components to achieve State water quality criteria, maintain ecological processes, and sustain appropriate uses.

Riparian and wetlands vegetation should have structural and species diversity characteristic of the stage of stream channel succession in order to provide forage and cover, capture sediment, and capture, retain, and safely release water (watershed function).

##### Upland Indicators:

- Canopy and ground cover, including litter, live vegetation, biological crust, and rock appropriate to the potential of the ecological site.
- Ecological processes are adequate for the vegetative communities.

##### Riparian Indicators:

- Stream side riparian areas are functioning properly when adequate vegetation, large woody debris, or rock is present to dissipate stream energy associated with high water flows.
- Elements indicating proper functioning condition such as avoiding accelerating erosion, capturing sediment, and providing for groundwater recharge and release are determined by the following measurements as appropriate to the site characteristics:
  - Width/Depth ratio;

- Channel roughness;
  - Sinuosity of stream channel;
  - Bank stability;
  - Vegetative cover (amount, spacing, life form); and
  - Other cover (large woody debris, rock).
- Natural springs, seeps, and marsh areas are functioning properly when adequate vegetation is present to facilitate water retention, filtering, and release as indicated by plant species and cover appropriate to the site characteristics.

#### Water Quality Indicators:

- Chemical, physical and biological constituents do not exceed the State water quality Standards.

The above indicators shall be applied to the potential of the ecological site.

#### GUIDELINES:

- 2.1 Management practices should maintain or promote appropriate stream channel morphology and structure consistent with the watershed.
- 2.2 Watershed management practices should maintain, restore or enhance water quality and flow rate to support desired ecological conditions.
- 2.3 Management practices should maintain or promote the physical and biological conditions necessary for achieving surface characteristics and desired natural plant community.
- 2.4 Grazing management practices will consider both the economic and physical environment, and will address all multiple uses including, but not limited to, (i) recreation, (ii) minerals, (iii) cultural resources and values, and (iv) designated wilderness and wilderness study areas.
- 2.5 New livestock facilities will be located away from riparian and wetland areas if they conflict with achieving or maintaining riparian and wetland functions. Existing facilities will be used in a way that does not conflict with achieving or maintaining riparian and wetland functions, or they will be relocated or modified when necessary to mitigate adverse impacts on riparian and wetland functions. The location, relocation, design and use of livestock facilities will consider economic feasibility and benefits to be gained for management of lands outside the riparian area along with the effects on riparian functions.
- 2.6 Subject to all valid existing rights, the design of spring and seep developments shall include provisions to protect ecological functions and processes.
- 2.7 When proper grazing practices alone are not likely to restore areas of low infiltration or

permeability, land management practices may be designed and implemented where appropriate. Grazing on designated ephemeral rangeland watersheds should be allowed only if (i) reliable estimates of production have been made, (ii) an identified level of annual growth or residue to remain on site at the end of the grazing season has been established, and (iii) adverse effects on perennial species and ecosystem processes are avoided.

- 2.8 Rangeland management practices should address improvement beyond these Standards, significant progress toward achieving Standards, time necessary for recovery, and time necessary for predicting trends.

### STANDARD 3. HABITAT AND BIOTA:

Habitats and watersheds should sustain a level of biodiversity appropriate for the area and conducive to appropriate uses. Habitats of special status species should be able to sustain viable populations of those species.

#### Habitat Indicators:

- Vegetation composition (relative abundance of species);
- Vegetation structure (life forms, cover, height, and age classes);
- Vegetation distribution (patchiness, corridors);
- Vegetation productivity; and
- Vegetation nutritional value.

#### Wildlife Indicators:

- Escape terrain;
- Relative abundance;
- Composition;
- Distribution;
- Nutritional value; and
- Edge-patch snags.

The above Indicators shall be applied to the potential of the ecological site.

## GUIDELINES:

- 3.1 Mosaics of plant and animal communities that foster diverse and productive ecosystems should be maintained or achieved.
- 3.2 Management practices should emphasize native species except when others would serve better for attaining desired communities.
- 3.3 Intensity, frequency, season of use and distribution of grazing use should provide for growth, reproduction, and, when environmental conditions permit, seedling establishment of those plant species needed to reach long-term land use plan objectives. Measurements of ecological condition, trend, and utilization will be in accordance with techniques identified in the Nevada Rangeland Handbook.
- 3.4 Grazing management practices should be planned and implemented to provide for integrated use by domestic livestock and wildlife, as well as wild horses and burros inside Herd Management Areas.
- 3.5 Management practices will promote the conservation, restoration and maintenance of habitat for special status species.
- 3.6 Livestock grazing practices will be designed to protect fragile ecosystems of limited distribution and size that support unique sensitive/endemic species or communities. Where these practices are not successful, grazing will be excluded from these areas.
- 3.7 Where grazing practices alone are not likely to achieve habitat objectives, land management practices may be designed and implemented as appropriate.
- 3.8 Vegetation manipulation treatments may be implemented to improve native plant communities, consistent with appropriate land use plans, in areas where identified Standards cannot be achieved through proper grazing management practices alone. Fire is the preferred vegetation manipulation practice on areas historically adapted to fire; treatment of native vegetation with herbicides or through mechanical means will be used only when other management techniques are not effective.
- 3.9 Rangeland management practices should address improvement beyond these Standards, significant progress toward achieving Standards, time necessary for recovery, and time necessary for predicting trends.

## GLOSSARY

Definitions are taken from "A Glossary of Terms Used in Range Management" developed through the Society for Range Management or Bureau of Land Management Technical Reference or from the Dictionary of Ecology, Evolution and Systematics except where noted. Other definitions are from Grazing Administration Regulations Code of Federal Regulations, Chapter 43 Sec. 4100.0-5. Definitions also include meanings that were developed by the Mojave Southern Resource Advisory Council to understand their intent in the Standards and Guidelines.

### -A-

**Annual Growth.** The amount of production of new above ground plant biomass for a given site during a given year.

### -B-

**Biodiversity.** The diversity of organisms in a region; made up of species diversity in individual community-types and the turnover of species across different community-types.

**Biological (Cryptogamic) Crust.** Community of non-vascular primary producers that occur as a "crust" on the surface of soils; made up of a mixture of algae, lichens, mosses, and cyanobacteria (bluegreen algae).

**Biotic.** Refers to living components of an ecosystem, e.g., plants and animals and micro-organisms.

### -C-

**Canopy.** (1) The vertical projection downward of the aerial portion of vegetation, usually expressed as a percent of the ground so occupied; (2) the aerial portion of the overstory vegetation.

**Canopy Cover.** The percentage of ground covered by a vertical projection of the outermost perimeter of the natural spread of foliage of plants. Small openings within the canopy are included. (BLM Technical Reference 4400-7)

**Climate.** The average or prevailing weather conditions of a place over a period of years. (BLM Technical Reference 4400-7)

**Conservation.** The planned management of natural resources; the retention of natural balance, diversity and evolutionary change in the environment.

The use and management of natural resources according to principles that assure their sustained economic and/or social benefits without impairment of environmental quality.

**Cover.** A. (1) The plants or plant parts, living or dead, on the surface of the ground. Vegetative cover or herbage cover is composed of living plants and litter cover of dead parts of plants; (2) The area of ground cover by plants of one or more species.

B. (1) The combined aerial parts of plants and mulch, and (2) shelter and protection for animals and birds. (BLM Manual 4400)

C. (1) Plant material, living (vegetative cover) and dead (litter cover) on the soil surface; (2) the area of ground covered by the canopy projections of a particular plant species, expressed as a scale or as a percentage of total ground surface area.

**Cultural Resources.** A broad, general term meaning any cultural property and any traditional lifeway value. (BLM Manual 8100)

**Cultural property.** A definite location of past human activity, occupation, or use identifiable through field inventory (survey), historical documentation, or oral evidence. (Manual 8100)

-D-

**Desert Pavement.** A cemented, hydrophobic layer of rocks or small pebbles that occurs over time on desert soil surfaces; prevents water infiltration into soils and wind/water erosion of the soil; often covered with a chemical varnish layer.

**Desired Natural Plant Community.** The type of plant community which is desired for a particular ecological site. This could include native and non-native species depending on the desired land use, but as a natural plant community it must have native species adapted to the climate and soil type as dominants or co-dominants in the community.

**Desired Plant Community.** Of the several plant communities that may occupy a site, the one that has been identified through a management plan to best meet the plan's objectives for the site. It must protect the site as a minimum.

**Diversity.** (1) The absolute number of species in a community; species richness; (2) A measure of the number of species and their relative abundance in a community; low diversity refers to few species or unequal abundances, high diversity to many species or equal abundances.

-E-

**Ecological Processes.** Natural functions including the hydrologic cycle, the nutrient cycle, and energy flow. (see also 43 CFR 4180.1(b))

**Ecological Site.** The kind of land with a specific potential natural community and specific physical site characteristics, differing from other kinds of land in its ability to produce vegetation and to respond to management. (BLM Manual 4400)

**Edaphic.** Refers to the soil.

**Endemic Species.** Native to, and restricted to, a particular geographical region, community type, or specific habitat.

**Ephemeral Rangelands.** Rangelands characterized by low, highly seasonal and often episodic

rainfall, resulting in annual plants comprising a significant proportion of annual primary production.

**Erosion.** (v.) Detachment and movement of soil or rock fragments by the action of water, wind, ice or gravity. (n.) The land surface worn away by running water, wind, ice, or other geologic agents, including such processes as gravitational creep.

**Exotic.** An organism or species which is not native to the region in which it is found. Synonym *non-native*: Not native; alien; a species that has been introduced into an area.

#### -F-

**Forage.** The plant material actually consumed by (or available to) grazing animals.

**Fragile Ecosystems.** Uncommon ecosystems of limited distribution and size that support unique sensitive/endemic species or communities; ecosystems that have low resilience to environmental stress or to disturbance.

**Frequency.** The ratio between the number of sample units that contain a species and the total number of sample units.

A quantitative expression of the presence or absence of individuals of a species in a population. It is defined as the percentage of occurrence of a species in a series of samples of uniform size. (BLM Technical Reference 4400-4)

#### -G-

**Grazing Distribution.** Dispersion of livestock grazing within a management unit or area.

**Ground Cover.** The percentage of material, other than bare ground, covering the land surface. It may include live and standing dead vegetation, litter, cobble, gravel, stones and bedrock. Ground cover plus bare ground would total 100 percent. (BLM Technical Reference 4400-4)

**Ground Water.** Subsurface water that is in the zone of saturation. The top surface of the ground water is the "water table." Source of water for wells, seepage, springs.

#### -H-

**Habitat.** The natural abode of a plant or animal, including all biotic, climatic, and edaphic factors affecting life.

**Hydrologic Balance.** The balance between hydrological inputs (infiltration of incident precipitation, run-on) and hydrological outputs (run-off, deep drainage) for an ecological site.

-I-

**Infiltration.** The flow of a fluid into a substance through pores or small openings. It connotes flow into a substance in contradistinction to the word *percolation*. The process by which water seeps into a soil, as influenced by soil texture, aspect and vegetation cover.

**Infiltration Rate.** Maximum rate at which soil under specified conditions can absorb rain or shallow impounded water, expressed in quantity of water absorbed by the soil per unit of time, e.g., inches/hour.

**Integrated Use.** To merge the use of each type of public land use through a series of land management practices.

-L-

**Land Use Plan.** Land use plan means a resource management plan, developed under the provisions of 43 CFR part 1600, or management framework plan. These plans are developed through public participation in accordance with the provisions of the Federal Land Policy and Management Act of 1976 and establish management direction for resource uses of public lands. (43 CFR 4100)

**Litter.** The uppermost layer of organic debris on the soil surface; essentially the freshly fallen or slightly decomposed vegetal material. (BLM Technical Reference 4400-4)

-M-

**Management Objective.** The objectives for which rangeland and rangeland resources are managed which includes specified users accompanied by a description of the desired vegetation and the expected products and/or values.

**Management Plan.** A program of action designed to reach a given set of objectives.

**Marsh.** Flat, wet, treeless areas usually covered by standing water and supporting a native growth of grasses and grasslike plants.

**Monitoring.** The orderly collection, analysis, and interpretation of resource data to evaluate progress toward meeting management objectives. (BLM Technical Reference 4400-7)

**Monitoring.** Monitoring means the periodic observation and orderly collection of data to evaluate: (1) Effects of management actions; and (2) Effectiveness of actions in meeting management objectives. (43 CFR 4100.0.5)

**Morphology.** The form and structure of an organism, with special emphasis on external features.

**Multiple Use.** The management of the public lands and their various resource values so that they are utilized in the combination that will best meet the present and future needs of the American people; making the most judicious use of the land for some or all of these resources

or related services over areas large enough to provide sufficient latitude for periodic adjustments in use to conform to changing needs and conditions; the use of some land for less than uses that takes into account the long-term needs of future generations for renewable and nonrenewable resources, including, but not limited to, recreation, range, timber, minerals watershed, wildlife and fish, natural scenic, scientific and historical values; and harmonious and coordinated management of the various resources without permanent impairment of the productivity of the land and the quality of the environment with consideration being given to the relative values of the resources and not necessarily to the combination of uses that will give the greatest economic return of the greatest unit output. (Federal Land Policy and Management Act)

-N-

**Native Species.** A species which is a part of the original fauna or flora of the area in question. Indigenous; living naturally within a given area and was part of the areas flora or fauna prior to human settlement of the region.

**Naturalized Species.** An exotic or introduced species that has become established and exhibits successful reproduction in an ecosystem.

-P-

**Percolation.** The flow of a liquid through a porous substance.

**Productivity.** The potential rate of incorporation or generation of energy or organic matter (biomass) by an organism, population or trophic unit per unit time per unit area; plant productivity is termed primary production, and animal productivity is termed secondary production.

**Proper Functioning Condition.** Riparian-wetland areas are functioning properly when adequate vegetation, landform, or large woody debris is present to dissipate stream energy associated with high waterflows, thereby reducing erosion and improving water quality; filter sediment, capture bedload, and aid floodplain development; improve flood-water retention and ground-water recharge; develop root masses that stabilized streambank against cutting action; develop diverse ponding and channel characteristics to provide the habitat and the water depth, duration, and temperature necessary for fish production, waterfowl breeding, and other uses; and support greater biodiversity. (BLM Technical Reference 1737-9)

-R-

**Range Improvement.** Range improvement means an authorized physical modification or treatment which is designed to improve production of forage; change vegetation composition; control patterns of use; provide water; stabilize soil and water conditions; restore, protect and improve the condition of rangeland ecosystems to benefit livestock, wild horses and burros, and fish and wildlife. The term includes but is not limited to, structures, treatment projects, and use of mechanical devices or modifications achieved through mechanical means.

**Residual Vegetation.** Amount, cover, and species composition of the vegetation on a site after it has been grazed for a period of time.

**Resource.** Any component of the environment that can be utilized by an organism.

**Riparian.** Pertaining to, living or situated on, the banks of rivers and streams. 'Xeroriparian' refers to being situated on dry washes (ephemeral streams).

#### -S-

**Seep.** Wet areas, normally not flowing, arising from an underground water source.

**Soil.** (1) The unconsolidated mineral and organic material on the immediate surface of the earth that serves as a natural medium for the growth of land plants. (2) The unconsolidated mineral matter on the surface of the earth that has been subjected to and influenced by genetic and environmental factors of parent material, climate (including moisture and temperature effects), macro- and micro-organisms, and topography, all acting over a period of time and producing a product -soil- that differs from the material it was derived in many physical, chemical, biological, and morphological properties and characteristics.

**Soil Productivity.** The organic fertility or capacity of a given area or habitat.

**Species.** A taxon of the rank species; which is the basic unit, and lowest principal category, of biological classification; in the hierarchy of biological classification, the category below genus; a group of organisms formally recognized as distinct from other groups.

**Species Composition.** The proportions of various plant species in relation to the total on a given area. It may be expressed in terms of cover, density, weight, etc. Synonym *Vegetative composition*.

**Surface Characteristics.** The amount of bare ground, litter, rock and basal cover of live vegetation, which may include cryptograms. (Nevada Rangeland Monitoring Handbook)

**Sustained yield.** The achievement and maintenance in perpetuity of a high level annual or regular periodic output of the various renewable resources of the public lands consistent with multiple use. (FLPMA)

#### -T-

**Traditional lifeway values.** The quality of being useful in or important to the maintenance of a specified social and/or cultural group's traditional systems of (a) religious belief, (b) cultural practice, or (c) social interaction, not closely identified with definite locations. Another group's shared values are abstract, nonmaterial, ascribed ideas that one cannot know about without being told. (BLM Manual 8100)

**Trend.** The direction of change in ecological status or resource value rating observed over time. Trend in ecological status should be described as *toward*, or *away from* the potential natural community, or as not apparent. (BLM Technical Reference 4400-4)

-U-

**Upland.** Terrestrial ecosystems located away from riparian zones, wetlands, springs, seeps and dry washes; ecosystems made up of vegetation not in contact with groundwater or other permanent water sources.

-V-

**Vegetative Life Form.** The characteristic structural features and method of perennation of a plant species, e.g., annuals, perennial forbs, shrubs, trees and succulents.

-W-

**Watershed.** (1) A total area of land above a given point on a waterway that contributes runoff water to the flow at that point. (2) A major subdivision of a drainage basin.

**Wetlands.** Areas characterized by soils that are usually saturated or ponded, i.e., hydric soils, that support mostly water-loving plants (hydrophytic plants).

In areas of arid low lying land that is submerged or inundated periodically by water, and is characterized by hydric soils that support mostly water-loving (hydrophytic) plants.