



---

**CENTRAL BASIN AND RANGE  
RAPID ECOREGIONAL ASSESSMENT  
FINAL MEMORANDUM I-2-C**

---

**Prepared for:**

Department of the Interior  
Bureau of Land Management  
Rapid Ecoregional Assessments

**Submission Date: November 19, 2010**

**Submitted to:**

Department of the Interior  
Bureau of Land Management, BC-662  
Building 50, Denver Federal Center  
P.O. Box 25047  
Denver, Colorado 80225-0047  
Attn: Craig Goodwin, Ecoregional Assessment Project Manager

**Submitted by:**

NatureServe  
1101 Wilson Boulevard, 15<sup>th</sup> Floor  
Arlington, Virginia 22209

---

Patrick Crist, Principal Investigator



# Contents

---

Executive Summary	4
Task 2 Objectives .....	4
Data Identification, Management and Evaluation .....	4
Data Evaluation Results for CEs .....	5
CE Class I: Terrestrial Coarse Filter .....	5
CE Class II: Terrestrial Fine Filter.....	6
CE Class III: Physical Feature – Sensitive Soils.....	7
CE Class IV: Aquatic Coarse Filter .....	7
CE Class V: Aquatic Fine Filter .....	7
Data Evaluation Results for CAs.....	8
Data Evaluation Results for Managed Lands .....	9
Data Evaluation Results for Management Questions .....	9
Introduction	10
Task 2 Objectives .....	10
Memorandum I-2-b Organization .....	10
Data Identification, Management and Evaluation	10
Secure File Transfer .....	10
SharePoint Site – Data Management .....	11
Data Management and Tracking.....	11
Data Evaluation .....	12
Ongoing Use of Master Data List.....	12
Identified Data Sources and Data Sets	12
Data Sources .....	12
Data Evaluation Results for Conservation Elements (CEs)	13
CE Class I: Terrestrial Coarse Filter .....	13
CE Class II: Terrestrial Fine Filter .....	15
CE Class III: Physical Feature - Sensitive Soils.....	17
CE Class IV: Aquatic Coarse Filter.....	17
CE Class V: Aquatic Fine Filter .....	19
Data Evaluation Results for CAs	19
Class I: Wildfire.....	19
Class II: Development .....	21
Urbanization .....	22
Infrastructure .....	22
Energy development.....	23

Hydrologic Change Agents .....	24
Mining .....	24
Military use/expansion areas .....	24
Air quality impacts (non attainment areas and dust).....	25
Recreation (OHV use, other intensive recreation, land sales, etc.).....	25
Refuse Management (landfills, sewage sludge disposal, nuclear disposal, etc.) .....	25
Agriculture .....	25
Exotic ungulate grazing.....	26
Class III: Invasives.....	26
Terrestrial Invasive Species .....	26
Aquatic Invasive Species.....	27
Class IV: Climate change .....	29
Data Evaluation Results for Managed Lands and Sites .....	32
PL Class I: Sites of High Biodiversity.....	32
PL Class II: Specially Designated Areas of Ecological or Cultural Value .....	32
PL Class III: Other Managed Lands .....	33
Summary Data Gaps and Recommendations for CEs, CAs, PLs, and MQs .....	33
CE Data Gaps and Recommendations .....	33
CA Data Gaps and Recommendations .....	33
Recommendations for Management Question Revisions.....	34
References .....	36
Appendix I: Master Data Table for the Central Basin and Range REA .....	40
Appendix II. Coarse-filter Conservation Elements for the Central Basins and Ranges Ecoregion.....	78
Appendix III: Current Draft of Fine-Filter Conservation Elements for the Central Basin and Range REA .....	79
Appendix IV. Management Questions: Implications from Data Evaluation.....	104

## Tables

---

Table 1. Data Suitability for CEs.....	33
Table 2. Summary of Data Suitability for CAs.....	34

## **Executive Summary**

Rapid Ecoregional Assessments (REAs) are the first step in the Bureau's Landscape Approach. REAs are intended to synthesize existing knowledge and information applicable to all lands and waters within the ecoregion. This synthesis aims to inform subsequent decision making, implementation, and monitoring by BLM and partners within the ecoregion, and should interact with ongoing scientific research as a foundation for science-based land management. REAs are organized into a series of phases and component tasks. Phase 1 includes tasks that clarify the scope, expected data and analysis approaches to be used, and culminating in a detailed workplan for the assessment. Phase 2 completes the preparation of data, conducts agreed-upon analyses, and documents assessment results. This memorandum summarizes the work, decisions, and remaining issues to be resolved for Task 2, Phase 1 for the Central Basin and Range Ecoregion. Here we conduct the assessment of data availability and quality representing the candidate conservation elements and change agents needed to answer the management questions. This memorandum is the final version (I-2-C) which has been revised and finalized by incorporating comments provided at AMT Workshop 2 or submitted separately to BLM.

### **Task 2 Objectives**

The objectives of Task 2 were:

1. Identify available data for the REA and obtain samples or metadata
2. Evaluate the data for utility (content, scale, completeness)
3. Evaluate the data quality (precision, consistency, documentation)
4. Make recommendations about data to be applied
5. Identify data gaps and proposed revisions to management questions, conservation elements, and change agents

### **Data Identification, Management and Evaluation**

NatureServe established a secure file transfer site for the BLM REA work that is being used for transferring data between NatureServe, NatureServe sub-contractors, and data sources. NatureServe has also created a secure collaborative workspace for the REA project team. The Data Management component of this SharePoint site includes resources such as technical instructions and documentation, and a "Master Data List" that NatureServe is using to track work status, conduct data evaluations, and prepare materials for reporting and creating tables. To create the Master Data List, NatureServe initially imported to our SharePoint site the spreadsheet provided by BLM "Att6.2-DMP-DataLayers.xlsx". NatureServe has added a number of attributes to track BLM requirements, as well as for internal data management and tracking purposes.

To ensure standardization and high quality products for BLM, many attributes in the Master Data List were configured as 'controlled value lists' with a menu of values to choose from or "Yes/No" check boxes. Full documentation for the Master Data List was created with definitions for all attributes, information about which are required, and when appropriate examples for the data entry.

The Master Data List is NatureServe's primary tool for managing information about the individual data sets as well as tracking status of the work being conducted. These include:

- information about the data set (filename, data source, citation, description, data type, scale, ISO category, currentness, data agreements, data restrictions / sensitivity, metadata )
- information about data management (filename and location where data resides on NatureServe's servers)
- work status (person requesting the data; data acquisition status and date; who needs to assess the data set; and review status)
- how data will be used in the REA analyses (type of CE, CA, or place; applicable REA(s))

The Master Data List is also NatureServe's primary tool for conducting the Phase I, Task 2 Data Quality Evaluation. To conduct this data evaluation, NatureServe started with the materials in "Appendix 7: Data Quality Evaluation Worksheet" and enhanced these by including a *Comments* field for each of the eleven Data Quality Evaluation criteria. This *Comments* field allows the expert conducting the data review to explain the assignment of one of the following confidence ratings: Very High, High, Moderate, Low, and Unknown. NatureServe's evaluation also includes information on the intended use of the data, and the suitability for these uses. Based on the information in the data evaluation attributes, NatureServe then assigns an Overall Data Confidence Rating score, again accompanied with comments where relevant.

The data evaluation process employed by NatureServe also encompasses metadata. The Metadata review includes an evaluation of whether the metadata are incomplete (missing key information), minimally complete (has abstract, purpose, currentness, scale, projection, attribute definitions, and contacts), or accepted. The metadata are reviewed to ensure that the projection / coordinates and datum (as appropriate) are provided.

The SharePoint system that NatureServe has developed for data management, tracking, and evaluation is both powerful and very flexible. NatureServe plans to build upon the existing structure to conduct subsequent evaluations for the REA, including the Phase I Task 3 identification, evaluation and recommendation of Models, Methods, and Tools to conduct the assessment.

### **Data Evaluation Results for CEs**

As established in memorandum I-1-C, a "coarse filter/fine filter approach" characterizes our method for CE identification; intending to provide an effective focus for the assessment. This applies both to criteria for selection of component elements, and to the various means of their treatment for analysis. Representative ecological types form our initial focus of assessment, and will be treated through mapping, modeling, and various assessment methods. Here these are described under CE Class I – Terrestrial Coarse Filter and CE Class IV Aquatic/Wetland Coarse Filter. Additionally, the desired CE of "sensitive soils" is addressed under CE Class III – Physical Features. Species data sets are summarized below within CE Class II – Terrestrial Fine Filter and CE Class V Aquatic/Wetland Fine Filter.

#### ***CE Class I: Terrestrial Coarse Filter***

The terrestrial "coarse filter" includes 17 terrestrial ecological system types and communities that express the predominant ecological pattern and dynamics of uplands across the ecoregion. NatureServe ecological classifications provided the basis for several current national or regional map products. These include ReGAP efforts from the southwest and California. Similarly, the national inter-agency LANDFIRE effort uses the same classification as the basis for their conceptual state-and-transition, vegetation dynamics models and spatial models aimed at characterizing fire regimes. LANDFIRE Existing Vegetation Type (EVT) classifies and maps types closely aligned with the ReGAP efforts. In 2009, NatureServe compiled ReGAP and LANDFIRE EVT to produce a composite national map of the current land cover and terrestrial ecological systems. In that effort (NatureServe 2009), numerous edits were completed and documented to reconcile the various map inputs into an integrated whole. **While this NatureServe (2009) map retains some error, as identified in this project review, we recommend that this map be used for this REA.** We propose to complete additional review and refinement of this map using other ancillary map layers to produce a best-available current distribution for terrestrial coarse-filter elements.

We intend to use several thousand georeferenced samples for spatial modeling of the predominant terrestrial coarse filter units under past, current, and future climate regimes. The LANDFIRE Reference Database (LFRDB) will be augmented with sample data consolidated and labeled for the SW ReGAP and CA ReGAP efforts. **We recommend use of these reference samples, totaling approximately 14,000 samples, for the REA study area.**

Part of our assessment of terrestrial coarse-filter CEs includes assessment of long-term trends in extent for each type; where we desire a mapped representation of each unit as it might occur today had no major land conversions occurred. The LANDFIRE Biophysical Settings (BpS) layer depicts, through inductive modeling, ‘potential’ or ‘historical’ distributions of terrestrial ecological system types assuming natural fire regimes have been unaltered. **The LANDFIRE BpS layer – with additional review and refinement – is what we recommend for use in this REA. Additionally, we will investigate linkages between existing NRCS Ecological Site Descriptions (and any mapped versions) for integration with these BpS maps.**

Ecological integrity is measured through a variety of means. One approach uses mapped ecological classification concepts as a focus. Criteria to evaluate a given example of a coarse-filter CE are documented through conceptual ‘state-and-transition’ vegetation dynamics models that reflect assumptions about succession and disturbance for a given type. Complementary to these ‘state-and-transition’ models, NatureServe has established and implemented methods for gauging the quality of ‘occurrences’ of each CE. Known as “element occurrence ranking criteria” measures of location size, condition, and landscape context are integrated to describe relative quality or condition against an assumed unaltered reference condition. These criteria are available for selected shrubland and riparian types (Appendix II). **We recommend use of these available ecological integrity criteria as inputs to our effort in this REA.**

Approaches to evaluating ecological integrity can also include development of spatial models to reflect patterns of land conversion that directly affect habitats and species. Three existing spatial models exist to gauge landscape conditions relevant to this REA: the Human Footprint model (2008), the NatureServe Landscape Condition model (2009), and the Theobald Natural Landscapes layer (2010). Each of these layers would be adequate for use in the REA though we acknowledge and can clarify concerns about the latter expressed by USGS reviewers. During Task 3, we will finalize the uses of these layers, and consider options for developing new iterations of these models.

### ***CE Class II: Terrestrial Fine Filter***

The “fine-filter” includes species that, due to their conservation status and/or specificity in their habitat requirements, are likely vulnerable to being impacted or lost from the ecoregion unless resource management is directed towards their particular needs. For species to be addressed in this assessment, we proposed, and the AMT accepted, several selection criteria for inclusion and treatment in the assessment. We continue to apply these criteria in an ongoing effort to finalize our list and approaches that will be used to handle all species meeting our criteria for inclusion, and that effort will be concluded during Phase I of this REA. Appendix III provides a summary of data for representing currently known locations for individual candidate species. These locational data fall into several categories. **Natural Heritage Programs** from this ecoregion maintain several thousand location records derived from field surveys over recent decades. A total of 15,309 records currently exist for our draft list of species CEs within this ecoregion (Appendix III).

A second major source of locational data for species CEs are habitat maps for all terrestrial vertebrates developed through **Gap Analysis projects** during the CA GAP project of the 1990s and the SW ReGAP completed in 2005. Species such as greater sage-grouse (*Centrocercus urophasianus*) will be addressed through the best-available locational data for various habitat components and subpopulation locations. We are still pursuing all best available data for sage grouse. **Critical habitat designations from the Fish and Wildlife Service** include several species from the CBR, including Mexican spotted owl (*Strix occidentalis lucida*), desert tortoise (*Gopherus agassizii*), and the Fish Slough milk-vetch (*Astragalus lentiginosus* var. *piscinensis*). Seasonal habitat data for **mule deer** exist as polygonal data sets at approximately 1:250,000 scale. These data should be adequate for purposes of the REA.

One additional category of habitat information for species CEs includes **identified corridors and crucial habitats as designated through state efforts coordinated by the WGA Western States**

**Decision Support System (DSS) initiative.** We anticipate gaining access to these data in collaboration with the WGA-sponsored Southwest States DSS project.

***CE Class III: Physical Feature – Sensitive Soils***

For this REA, sensitive soils can be depicted across the ecoregion by combining several spatial data sets. First SSURGO data are available for significant portions of the ecoregion. Most variables identified by BLM can be extracted from SSURGO data for a meaningful representation of this CE which. However, given incompleteness of SSURGO in this area, we will utilize draft soil map information as it becomes available from UT, NV, and CA state offices of NRCS. For UT, these include compilations of NPS and FS sources. We will coordinate with BLM and NRCS scientists to resolve availability issues. During Task 3 we will explore limitations of current data and explore additional modeling needs for these features using 10m<sup>2</sup> digital elevation for landform models and the spatially coarser STATSGO soils data (possibly further augmented by surficial material lithology data see e.g., Sayre et al. 2009).

***CE Class IV: Aquatic Coarse Filter***

Aquatic CEs combine what are commonly referred to as ‘aquatic’ habitats (streams, rivers, lakes, etc.) with ‘wetland’ communities (marsh, swamp, floodplain bottomlands) and ‘riparian’ communities (mosaic of wetland and intermittently flooded habitats). The **NatureServe composite ecological systems map** (NatureServe 2009) depicts current distributions of the primary wetland and riparian components of aquatic CEs. We propose to complete additional review and refinement of this map using several primary data sources. These include **SSURGO**, where available, for depicting hydric soils with natural land cover; National Wetland Inventory (**NWI**) for wetlands locations; and **NHD Plus** (1:24K scale data) for streams, lakes, intermittent washes, and playas. **Desert spring and seep** locations exist primarily for Nevada, but we continue to identify data from surrounding states.

As with terrestrial CEs, ecological integrity for aquatic CEs is measured through a variety of means. NatureServe has established and implemented methods for gauging the quality of individual occurrences of each CE, as described above for terrestrial CEs. Available criteria for aquatic CEs pertain to wetland and riparian ecological system types from the CBR and adjacent ecoregions. **We recommend use of these available ecological integrity criteria as inputs to our effort in this REA.**

***CE Class V: Aquatic Fine Filter***

Similarly referenced above under the terrestrial fine-filter, **Natural Heritage Programs** from this ecoregion maintain several thousand location records derived from field surveys over recent decades. A total of 2,192 records currently exist for our draft list of aquatic species CEs within this ecoregion (Appendix III). Critical habitat designations from the Fish and Wildlife Service include 10 fish species from the CBR. EcoAnalysts Inc., has conducted taxonomic identification of aquatic macro invertebrates, including natives and invasives, for hundreds of projects and hundreds of clients in the Western USA. State Game and Fish agencies also should have additional location and habitat data for aquatic species of concern to the REA. We will explore their availability within the context of discussions with the WGA-sponsored Southwest DSS effort during Phase I.

**Summary of data suitability for CEs.**

<b>Conservation Element Category</b>	<b>Number of Elements</b>	<b>Data Suitable?</b>
Basin Dryland Ecosystems	10	Yes
Montane Dryland Ecosystems	7	Yes
Basin Wet Ecosystems	6	Yes

Conservation Element Category	Number of Elements	Data Suitable?
Montane Wet Ecosystems	3	Yes
Nested Terrestrial Habitat-Based Species Assemblages	TBD	high probability
Nested Aquatic Habitat-Based Species Assemblages	TBD	Yes
Individual Species	TBD	high probability
<b>Desired Conservation Elements</b>		
Mule Deer		Yes
Sensitive Soils		high probability

### **Data Evaluation Results for CAs**

We evaluated data to represent the four CA classes: I – Wildfire, II – Development, III – Invasives, and IV—Climate Change. Sufficient comprehensive data sets exist to model the Wildfire and Climate Change classes. For other CAs, there are critical data gaps for which we are still pursuing data sources and will also investigate modeling of these in Task 3. The data availability for the Development and Invasives classes is more limited, however, some Development subclasses are well represented in the extant data. In the Development class, sufficient data exists to adequately depict or readily model urbanization (current and for 2025), infrastructure, energy development (current and potential), air quality impacts, and hydrology. Centroid locations of mining and refuse management are available and the spatial footprint of these features may be approximated with supporting land use/land cover (LU/LC) data (to be explored in Task 3). While we currently have limited data for recreation, more detailed information is forthcoming from the NOC and will be evaluated along with potential to model OHV distribution in particular will be investigated in Task 3.

Surface disturbances within military use areas (large extents of bare ground, urbanized areas) can be detected using satellite-derived LU/LC classifications. Noise impacts from low-flying and super-sonic aircraft may be approximated with flight path and low-flying areas however there are not definite links between these areas and noise impacts to wildlife. We have therefore recommended removing aircraft noise consideration from our assessment.

Given the complex nature and potential effects of exotic ungulate grazing, we have very limited data on this CA and have proposed simplifying the treatment of exotic ungulate grazing from our original proposal during Task 1.

While aquatic invasives are adequately represented, data for their terrestrial counterparts was not as forthcoming or comprehensive. The NOC is planning to provide additional data. Regardless, modeling these species or building upon existing models produced by the Nevada Natural Heritage program will be investigated in Task 3. This effort may incorporate disparate sources of location data from counties, the BLM and the LANDFIRE vegetation reference plots. We will also conduct further investigation of the invasive vulnerability component of The Human Footprint map.

### **Summary of data suitability for CAs.**

Change Agent Class	Number of Subclasses	Data Suitable?
Wildfire	2	Yes
Development	10	Yes for most subclasses
Invasives	2	Limited but suitable for several

<b>Change Agent Class</b>	<b>Number of Subclasses</b>	<b>Data Suitable?</b>
Climate	TBD	Yes/resolution issues

**Data Evaluation Results for Managed Lands**

We found adequate data to represent managed lands, which we categorize as: I—Sites of High Biodiversity, II—Specially Designated Areas of Ecological or Cultural Value, and III—Other Managed Lands. We will evaluate crucial habitats and any other similar information created by the Southwest States DSS project as it comes available and seek input from the AMT on utility of those areas as assessment units.

**Data Evaluation Results for Management Questions**

Treatment of individual management questions (MQs) is described in Appendix IV. Generally, data appears available and suitable to answer most of the MQs though several data sets are yet to be acquired and evaluated. MQs related to exotic ungulate grazing are most tenuous from our data evaluation to date.

# Task 2 Identify, Evaluate, and Recommend Potential Data

---

## **Introduction**

Rapid Ecoregional Assessments (REAs) are the first step in the Bureau's Landscape Approach. REAs are intended to synthesize existing knowledge and information applicable to all lands and waters within the ecoregion. This synthesis aims to inform subsequent decision making, implementation, and monitoring by BLM and partners within the ecoregion, and should interact with ongoing scientific research as a foundation for science-based land management. REAs are organized into a series of phases and component tasks. Phase 1 includes tasks that clarify the scope, expected data and analysis approaches to be used, and culminating in a detailed workplan for the assessment. Phase 2 completes the preparation of data, conducts agreed-upon analyses, and documents assessment results. This memorandum summarizes the work and decisions for Task 2, Phase 1 for the Central Basin and Range Ecoregion. Here we conduct the evaluation of data availability and quality representing the candidate conservation elements and change agents needed to answer the management questions. This memorandum is the final version (I-2-b) which has been revised and finalized by incorporating comments provided at AMT Workshop 2 or submitted separately to BLM.

## **Task 2 Objectives**

The objectives of Task 1 were:

1. Identify available data for the REA and obtain samples or metadata
2. Evaluate the data for utility (content, scale, completeness)
3. Evaluate the data quality (precision, consistency, documentation)
4. Make recommendations about data to be applied
5. Identify data gaps and proposed revisions to management questions, conservation elements, and change agents

## **Memorandum I-2-b Organization**

This memorandum summarizes our evaluation of data availability and quality to represent the conservation elements and change agents needed to answer the management questions. Additionally, data that reflect locations of managed lands, specially designated lands, and area of high significance from existing natural resource prioritization efforts (e.g., SWAPs) are also addressed. The memorandum is organized according to the objectives. Details are provided in tables in the appendices.

## **Data Identification, Management and Evaluation**

### **Secure File Transfer**

NatureServe established a secure file transfer site for the BLM REA work that is being used for transferring data between NatureServe, NatureServe sub-contractors, and data sources. The secure file upload requires a username and password, and files placed in this repository can only be retrieved by NatureServe data management staff. This upload resource is being used to allow people to contribute data in a secure manner. For datasets that NatureServe needs to share with REA subcontractors, NatureServe has established a secure file download site that requires a different username and password.

All usernames and passwords are tightly controlled and only distributed to the relevant project team members.

### **SharePoint Site – Data Management**

Based on the materials developed for Phase I Task 1, NatureServe identified the Conservation Elements (CEs), Change Agents (CAs), Places (PLs), and other data desired to evaluate for possible inclusion in the assessment. Working closely with BLM to minimize redundancy in data requests, the responsibility for identifying data sets was assigned to various team members based on areas of expertise. When possible, we obtained the full data set plus all supporting metadata and reports. When the data were not available, we requested and obtained at a minimum metadata and supporting materials, with sample data as available. As each member of the team worked through their list of data sets, the information was entered in the Master Data List (described below) and the appropriate team experts notified so they could begin the data quality evaluation process.

Using Microsoft SharePoint software, NatureServe has created a secure collaborative workspace for the REA project team. The Data Management component of this SharePoint site includes resources such as technical instructions and documentation, including data management guideline materials provided by BLM, and a “Master Data List” that NatureServe is using to track work status, conduct data evaluations, and prepare materials for reporting and creating tables.

All members of the NatureServe REA team received training via Web-Ex in the proper use of the BLM REA project SharePoint site, and additional support is available as needed by the project Data Management lead and NatureServe IT staff.

To create the Master Data List, NatureServe initially imported to SharePoint the spreadsheet provided by BLM “Att6.2-DMP-DataLayers.xlsx”. After reviewing the materials in the document “Rapid Ecoregional Assessment (REA) Data Management Plan: Contractor Guidance”, NatureServe added attributes from the following appendices (from BLM’s data management guidelines) critical for achieving compliance with those guidelines:

- Appendix 7: Data Quality Evaluation Worksheet
- Appendix 8: QA/QC Checklist
- Appendix 9: Pre-Acquisition Data Assessment Worksheet

In addition, the NatureServe project team added attributes to the Master Data List for internal data management and tracking purposes.

To ensure standardization and high quality products for BLM, many attributes in the Master Data List were configured as controlled value lists with a menu of values to choose from or “Yes/No” check boxes. Full documentation for the Master Data List was created with definitions for all attributes, information about which are required, and when appropriate examples for the data entry.

The SharePoint site allows the NatureServe team the flexibility to have multiple people working collaboratively on the Master Data List and allows customization through filters and creating “views” so that individual users can focus on any subset of attributes and/or data records of interest. Because SharePoint is fully integrated with other Microsoft software, NatureServe can export from the Master Data List to Excel and create tables for reports.

### **Data Management and Tracking**

The Master Data List is NatureServe’s primary tool for managing information about the individual data sets as well as tracking status of the work being conducted. These include:

- information about the data set (filename, data source, citation, description, data type, scale, ISO category, currentness, data agreements, data restrictions / sensitivity, metadata )

- information about data management (filename and location where data resides on NatureServe’s servers)
- work status (person requesting the data; data acquisition status and date; who needs to assess the data set; and review status)
- how data will be used in the REA analyses (type of CE, CA, or place; applicable REA(s))

### **Data Evaluation**

The Master Data List is also NatureServe’s primary tool for conducting the Phase I, Task 2 Data Quality Evaluation. To conduct this data evaluation, NatureServe started with the materials in “Appendix 7: Data Quality Evaluation Worksheet” and enhanced these by including a *Comments* field for each of the eleven Data Quality Evaluation criteria. This *Comments* field allows the expert conducting the data review to explain the assignment of one of the following confidence ratings: Very High, High, Moderate, Low, and Unknown. NatureServe’s evaluation also includes information on the intended use of the data, and the suitability for these uses. Based on the information in the data evaluation attributes, NatureServe then assigns an Overall Data Confidence Rating score, again accompanied with comments where relevant.

The data evaluation process employed by NatureServe also encompasses metadata. The Metadata review includes an evaluation of whether the metadata are incomplete (missing key information), minimally complete (has abstract, purpose, currentness, scale, projection, attribute definitions, and contacts), or accepted. The metadata are reviewed to ensure that the projection / coordinates and datum (as appropriate) are provided. And the reviewer can enter comments about the metadata, particularly if there are areas that are incomplete or questions that need to be resolved.

### **Ongoing Use of Master Data List**

The SharePoint system that NatureServe has developed for data management, tracking, and evaluation is both powerful and very flexible. NatureServe plans to build upon the existing structure to conduct subsequent evaluations for the REA, including the Phase I Task 3 identification, evaluation and recommendation of Models, Methods, and Tools to conduct the assessment.

In addition, the information already captured in the Master Data List provides the foundation for the Phase II Task 1 compilation and generation of source data sets. We have already begun tracking which data sets have been requested, acquired, and their physical management. This will be expanded to include generated data sets, as well as the scripts and modeling processes used. We will build on the existing “metadata” attributes to track the creation and review of metadata for generated data sets, and will apply the existing Data Quality Evaluation to these generated data sets.

## **Identified Data Sources and Data Sets**

Appendix I identifies and characterizes all data sets evaluated in this Task. Details on the evaluation are described under the CE and CA sections below and their respective appendices as well as data evaluation forms delivered separately to BLM. To date, we have evaluated over close to 200 data sets and recommended many dozens as suitable for the REA.

### **Data Sources**

We identified the following primary data sources and obtained sample data and or metadata:

- BLM
- USGS
- EPA

- LANDFIRE
- Natural Heritage Programs
- NatureServe
- The Nature Conservancy
- NRCS
- State Wildlife Agencies
- State Water Quality agencies
- NREL
- SAGEMAP

## **Data Evaluation Results for Conservation Elements (CEs)**

All of the described data sets below are proposed for use in the REA following our evaluation unless otherwise described. Conservation Element (CE) data sets were identified and evaluated; with results detailed in Appendices II and III. Here we summarize our evaluation and results by CE Class; with categories reflecting major CE types, their distribution, and ecological integrity. Base biophysical data most strongly tied to CE distributions and are listed within CE Classes I-V. For this report we have combined “core” and “desired” CEs within each of these categories.

As established in memorandum I-1-C, a “coarse filter/fine filter approach” characterizes our method for CE identification; intending to provide an effective focus for the assessment. This applies both to criteria for selection of component elements, and to the various means of their treatment for analysis. Representative ecological types form our initial focus of assessment and will be treated through mapping, modeling, and various assessment methods. These are described under CE Class I – Terrestrial Coarse Filter and CE Class IV Aquatic/Wetland Coarse Filter. Additionally, the desired CE of “sensitive soils” is addressed under CE Class III – Physical Features. Species data sets are summarized below within CE Class II – Terrestrial Fine Filter and CE Class V Aquatic/Wetland Fine Filter.

### **CE Class I: Terrestrial Coarse Filter**

The terrestrial “coarse filter” includes 17 terrestrial ecological system types and communities that express the predominant ecological pattern and dynamics of uplands across the ecoregion. These classified units: a) characterize each component of the ecoregion conceptual model, b) define the vast majority of this ecoregion’s lands and waters, and c) reflect described ecological types with distributions concentrated within this ecoregion. By treating these in our assessment we aim to adequately treat the habitat requirements of most characteristic native species, ecological functions, and ecosystem services. Ecological models (both conceptual and spatial) for these coarse filter elements will form a major focus for this ecoregional assessment. Here we briefly summarize data sets applicable to mapping the location and extent (current and probable/historical) of terrestrial coarse filter units. Additionally, we summarize data sets for use in documenting their natural ecological dynamics and integrity.

Among numerous older regional vegetation maps, the “Sagestitch” effort under SAGEMAP reconciled existing vegetation maps from the 1990s aiming to depict rangewide distributions of sagebrush and related vegetation (Comer et al. 2002). These maps may be useful for review and refinement of more current map products. NatureServe ecological classifications provided the basis for several current national or regional map products (see <http://www.natureserve.org/explorer/> for more detailed descriptions of ecosystem types listed for this REA). These include ReGAP efforts from the southwest (Lowry et al. 2007) (including NV, AZ, and UT) and CA ReGAP (in progress) and the Northwest ReGAP effort (also in progress). The Northwest ReGAP land cover products include only a small portion of the CBR REA area in southern Idaho. It incorporated the Shrubmap effort for map

zones overlapping with SW ReGAP in NV. Similarly, the national inter-agency LANDFIRE effort uses the same classification as the basis for their conceptual state-and-transition, vegetation dynamics models and spatial models aimed at characterizing fire regimes (<http://www.landfire.gov/>). LANDFIRE Existing Vegetation Type (EVT) classifies and maps types closely aligned with the ReGAP efforts. In these cases, they also used common input data sets with the ReGAP efforts, including field reference samples and imagery. However, within this project area, there are considerable discrepancies between LANDFIRE EVT and SW ReGAP. We trace many of these to sample plot labeling error since there are distinct differences between expert-labeled ReGAP samples, and subsequent auto-key labels applied by LAMNFIRE (to the sample plot).

In 2009, NatureServe compiled ReGAP and LANDFIRE EVT (for California in this project area) to produce a composite national map of the current land cover and terrestrial ecological systems. In that effort (NatureServe 2009), numerous edits were completed and documented to reconcile the various map inputs into an integrated whole. **While this NatureServe (2009) map retains some error, as identified in this project review, we recommend that this map be used for this REA.** We propose to complete additional review and refinement of this map using other ancillary map layers to produce a best-available current distribution for terrestrial coarse-filter elements. Additional local data sets, such as existing vegetation maps from districts within the Humboldt-Toiyabe National Forest, will be accessed to assist with this review and refinement of the ecoregional coverage.

Reference sample data from field surveys identify the vegetation type, physiognomy, and plant species composition. We intend to use several thousand georeferenced samples for spatial modeling of the predominant terrestrial coarse filter units under past, current, and future climate regimes. The LANDFIRE reference database (LFRDB) was developed between 2004 and 2009, consolidating field samples from across federal and non-federal sources for use in spatial modeling. The LFRDB will be reviewed, updated, and augmented (for certain sparsely vegetated and wetland/riparian types) with sample data consolidated and labeled for the SW ReGAP and CA ReGAP efforts. **We recommend use of these reference samples, totaling approximately 14,000 samples, for the REA study area.** The LFRDB and ReGAP data will also provide reference samples for invasive plant species assessment detailed below. See Appendix II for summary statistics on reference samples available for each coarse-filter CE.

Part of our assessment of terrestrial coarse-filter CEs includes assessment of long-term trends in extent for each type; where we desire a mapped representation of each unit as it might occur today had no major land conversions occurred. Three primary data sets exist for this purpose. The LANDFIRE Biophysical Settings (BpS) layer depicts, through inductive modeling, ‘potential’ or ‘historical’ distributions of terrestrial ecological system types assuming natural fire regimes have been unaltered. A second national “footprint” map from USGS (Sayre et al. 2009) aims at the same goal, but through deductive modeling with a more limited set of national spatial data inputs. NatureServe subsequently completed a third model for the Great Basin Integrated Landscape Monitoring effort (Comer et al. 2009) which experimentally combined inductive modeling approach but using the national spatial data inputs from the USGS “footprint” map. While the latter data sets are suitable national-scaled analysis, **the LANDFIRE BpS layer – with additional review and refinement – is what we recommend for use in this REA.** During Task 3, we will investigate the utility of incorporating available data sets now provided through the NASA TOPS effort (<http://ecocast.arc.nasa.gov/>), such as ASTER-derived land surface temperature (25m), SRTM-derived topography, or SCAN-derived soil moisture observations, into BpS map refinements. Additionally, NRCS Ecological Site Descriptions, where developed and mapped using SSURGO data, may provide additional useful information for both conceptual models and map refinements to this BpS layer. **We will investigate linkages between existing NRCS Ecological Site Descriptions (and any mapped versions) for integration with these BpS maps.**

Ecological integrity is measured through a variety of means. One approach uses mapped ecological classification concepts as a focus. Criteria to evaluate a given example of a coarse-filter CE are documented through conceptual ‘state-and-transition’ vegetation dynamics models that reflect assumptions about succession and disturbance for a given type. These are available in several forms, and will be referenced more fully below under CA Class I – Wildfire. Complementary to these ‘state-and-transition’ models are criteria to integrate assumptions about ecological condition for each type. NatureServe has established and implemented methods for gauging the quality of ‘occurrences’ of each CE. Known as “element occurrence ranking criteria” measures of location size, condition, and landscape context are integrated to describe relative quality or condition against an assumed unaltered reference condition. NatureServe methods have evolved over the past decade, and for this REA, some criteria are available from the adjacent Utah High Plateaus ecoregion that were developed using NatureServe standards *circa* 2000. More recent work from the CO and WA Natural Heritage programs include criteria under more recent 2008 NatureServe standards. These criteria are available for selected sagebrush shrubland and montane riparian types that may be readily adapted for use in the CBR (Appendix II). **We recommend use of these available ecological integrity criteria as inputs to our effort in this REA.** This existing information provides input primarily to conceptual modeling, where we state our current assumptions about key ecological attributes that drive ecological processes and support a given recognizable biotic assemblage. For example, these conceptual models make statements about expected natural fire frequency, intensity, and spatial character. They may document current knowledge of hydrologic flow patterns that produce recognizable patterns in riparian vegetation. They may state assumptions about the expected diversity of native plant species one would tend to encounter, and observations on the effects of certain invasive species introductions into the system type. Given these assumptions, measurable criteria and indicators are established for evaluation of the ecological system, either as individual patches, or across a regional distribution. For purposes of this REA, we aim to evaluate established criteria that may be readily applied with available data. In most instance, we will be limited to applying indicators of ecological integrity that can be measured through remote sensing and spatial modeling.

Spatial models intending to integrate human alterations and ecological effects within this ecoregion have been developed. The Human Footprint in the West map depicts an ‘ecological footprint’ using 14 land cover variables, including land cover classes and transportation corridors at a base resolution of 180m<sup>2</sup> (Leu et al. 2008). Following an identical logic, NatureServe completed a similar national model of Landscape Condition using 17 variables and a base map resolution of 90m<sup>2</sup> (Comer and Hak 2009) including both ‘direct impact’ measures and a ‘distance decay’ function for each input layer. Theobald’s Natural Landscapes layer (2010) detailed Class II: Development section below, provides a third option for consideration. Each of these layers would be adequate for use in the REA though we acknowledge and will address concerns about the latter voiced by USGS AMT reviewers. During Task 3, we will finalize our selection and proposed use of these layers, and propose modified forms of applying these types of models. In most instances, we anticipate being able to create spatial models that depict a) the current location of a given CE, b) a spatial model of apparent landscape conditions that tend to effect the ecological integrity of the CE at any given location, and c) summary information organized into watershed units, regular spatial grids, or other spatial reporting unit, to indicate the relative condition of the CE. To the degree that these same inputs can be developed for each time series of the REA (current, mid-century, and perhaps one date in between), reporting on ecological integrity of a similar nature will be feasible.

### **CE Class II: Terrestrial Fine Filter**

The “fine-filter” includes species that, due to their conservation status and/or specificity in their habitat requirements, are likely vulnerable to being impacted or lost from the ecoregion unless resource

management is directed towards their particular needs. For species to be addressed in this assessment, we proposed, and the AMT accepted, several selection criteria for inclusion and treatment in the assessment. These criteria include:

- a. All taxa listed under Federal or State protective legislation (including species, subspecies, or designated subpopulations)
- b. Full species with NatureServe Global Conservation Status rank of G1-G3<sup>1</sup>
- c. Full species or subspecies listed as BLM Special Status and those listed by applicable SWAPs with habitat included within the ecoregion
- d. Full species and subspecies scored as *Vulnerable* within the ecoregion according to the NatureServe Climate Change Vulnerability Index (CCVI)<sup>2</sup>.

One additional species, mule deer (*Odocoileus hemionus*), was included as a desired conservation element. Appendix III includes a current draft list for the ecoregion for species under criteria a-d above, and has had approximately 110 taxa added since Memo I-2-a was issued. The additional taxa are those we have determined to probably occur in the CBR and are listed by BLM as “sensitive” or “special status” from AZ, CA, NV and UT, or are animals listed in the relevant SWAPs that were not previously on our list. During Task 3, this list will be reviewed by local experts for their inclusion within the ecoregion. We anticipate a number of taxa now on this list will be removed after we determine the details of their distribution. Finalizing the list of species meeting these criteria is an ongoing effort to be concluded during Phase I of this REA. We have established several distinct approaches to treating species that meet established criteria for inclusion in the REA. These include:

- Species assumed to be adequately **represented indirectly through the assessment of major “coarse-filter”** ecological systems of the ecoregion. For example, species strongly affiliated with desert springs may be adequately treated in the REA through assessment of desert springs themselves.
- Species assumed to be adequately **represented indirectly as ecologically-based assemblages**. That is, due to similar group behavior and habitat requirement, a recognizable species assemblage is defined and treated as the unit of analysis. Examples could include bat hibernacula, treating multiple species of bats; all or some of whom are of conservation concern. Similarly, migratory bird stopover sites or raptor nesting/foraging zones could also be treated as multi-species assemblages.
- Species which should be **best addressed as individuals** in the assessment. These include those species meeting our criteria for assessment that cannot be presumed to be included in the previous two categories. This will tend to include many major ‘landscape’ species that range over wide areas within the ecoregion and with clearly distinct habitat requirements from all other taxa of concern.
- Species of concern from the latter category that have **very narrow distributions; limited to one BLM management jurisdiction**, we are gathering current locational information, but will not aim to develop conceptual models for these elements. We will continue to work with the AMT to determine appropriate means to spatially represent these elements within this REA.

It also remains an ongoing effort to finalize which approach will be used to handle all species meeting our criteria for inclusion, and that effort will be concluded during Phase I of this REA. Our team will further consult previous relevant work (e.g., Wisdom et al. 2004) and local expertise. Appendix III provides a summary of data for representing currently known locations for individual species. These locational data fall into several categories. Natural Heritage Programs from this ecoregion maintain several thousand location records derived from field surveys over recent decades. These data include

---

<sup>1</sup> See <http://www.natureserve.org/explorer/ranking.htm> for NatureServe Conservation Status Rank definitions

<sup>2</sup> See <http://www.natureserve.org/prodServices/climatechange/ccvi.jsp> for more on the NatureServe CCVI

field ‘observations’ and ‘element occurrences’ of species populations; the latter resulting from a systematic processing of ‘observations’ into standardized representations that consider distances separating each observation. A total of 15,309 records currently exist for our draft list of species CEs within this ecoregion (Appendix III).

A second major source of locational data for species CEs are habitat maps for all terrestrial vertebrates developed through Gap Analysis projects during the CA GAP project of the 1990s and the SW ReGAP completed in 2005. Appendix III references CEs for which we have data from these efforts. Some species have had much greater attention and data developed for their conservation. Species such as greater sage-grouse (*Centrocercus urophasianus*) will be addressed through the best-available locational data for various habitat components and subpopulation locations. We are still pursuing all best available data for sage grouse. Critical habitat designations from the Fish and Wildlife Service include several species from the CBR, including Mexican spotted owl (*Strix occidentalis lucida*), desert tortoise (*Gopherus agassizii*), and the Fish Slough milk-vetch (*Astragalus lentiginosus* var. *piscinensis*). Seasonal habitat data for mule deer exist as polygonal data sets at approximately 1:250,000 scale developed by Utah State University Extension & RS/GIS Laboratory & National Fish and Wildlife Foundation. These data should be adequate for purposes of the REA.

One additional category of habitat information for species CEs includes identified corridors and crucial habitats as designated through state efforts coordinated by the Western Governor’s Association Western States Decision Support System (DSS) initiative. We anticipate gaining access to these data in collaboration with the WGA-sponsored Southwest States DSS project.

### **CE Class III: Physical Feature - Sensitive Soils**

From current BLM definition: “Sensitive soils” are those identified as having characteristics that make them extremely susceptible to impacts or they may be more difficult to restore or reclaim after disturbance -- characteristics such as high wind or water erosion hazard (steep slopes), moderate to high salinity, low nutrient levels, low water holding capacity (droughty), or high water table (wetland/riparian soils). Information used to identify sensitive soils includes NRCS published soil surveys, ecological site descriptions, local monitoring records and research studies.”

For this REA, sensitive soils can be depicted across the ecoregion by combining several spatial data sets. First SSURGO data are available for significant portions of the ecoregion. Most variables listed above are tracked in some form by polygon and can be extracted from SSURGO data for a meaningful representation of this CE. However, given incompleteness of SSURGO in this area, we will utilize draft soil map information as it becomes available from UT, NV, and CA state offices of NRCS. For UT, these include compilations of NPS and FS sources. We will coordinate with BLM and NRCS scientists to resolve availability issues. During Task 3 we will explore limitations of current data and explore additional modeling needs for these features using 10m<sup>2</sup> digital elevation for landform models and the spatially coarser STATSGO soils data (possibly further augmented by surficial material lithology data see e.g., Sayre et al. 2009).

Additional discussion has centered on the potential treatment of biotic soil crusts. We agreed that treatment of soil crusts is best included within the assessment of ecological integrity for coarse filter CEs where these crusts play a significant role. During task 3 we will review current material (e.g., Rosentrater and Pellant *in prep.*) and document feasible methods for treatment of this issue.

### **CE Class IV: Aquatic Coarse Filter**

As established in memorandum I-1-c, Aquatic CEs nest from the ecoregion-wide conceptual model that defines all “wet” ecosystem types. These combine what are commonly referred to as ‘aquatic’

habitats (streams, rivers, lakes, etc.) with ‘wetland’ communities (marsh, swamp, floodplain bottomlands) with ‘riparian’ communities (mosaic of wetland and intermittently flooded habitats). Our aim is to provide a map depicting historical and current distributions for each of the nine coarse-filter CEs. The NatureServe composite ecological systems map (NatureServe 2009) depicts current distributions of the primary wetland and riparian components of aquatic CEs. Again, this coverage was derived largely from the SW ReGAP and LANDFIRE EVT maps. The LANDFIRE Biophysical Settings (BpS) map depicts in a generalize fashion, the ‘potential’ or ‘historical’ distribution of the CEs. We propose to complete additional review and refinement of these two maps using several primary data sources. These include SSURGO, where available, for depicting hydric soils with natural land cover; National Wetland Inventory (NWI) for wetlands locations; and NHD Plus (1:100K and 1:24K scale data) for streams, lakes, intermittent washes, and playas. Desert spring and seep locations exist primarily for Nevada, but we continue to identify data from surrounding states.

As with terrestrial CEs, ecological integrity for aquatic CEs is measured through a variety of means. NatureServe has established and implemented methods for gauging the quality of individual occurrences of each CE, as described above for terrestrial CEs. These “element occurrence ranking criteria” specify measures of the size, condition, and landscape context with which to describe the relative quality or condition of any occurrence of a CE in comparison to an assumed unaltered reference condition. Available criteria for aquatic CEs pertain to wetland and riparian ecological system types from the CBR and adjacent ecoregions. NatureServe methods have evolved over the past decade; for this REA, some criteria are available from the adjacent Utah High Plateaus ecoregion that were developed using NatureServe standards *circa* 2000. More recent work from the CO and WA Natural Heritage programs include criteria under more recent 2008 NatureServe standards. These criteria are available for selected riparian and other wetland types (Appendix II). **We recommend use of these available ecological integrity criteria as inputs to our effort in this REA.**

The element occurrence ranking criteria include information on both the biotic and abiotic (physical habitat) condition of a CE occurrence and information on its landscape context. The identification of these criteria rests on a conceptual ecological model for each CE. For terrestrial and wetland (including riparian) CEs, these models are often state-transition models, as noted above. For aquatic CEs or the strictly aquatic components of combined aquatic-riparian-wetland CEs, these models more often are causal diagrams such as those pioneered by Karr et al. (1986). These “ecological integrity diagrams” identify the key biotic attributes of a CE: a) key abiotic attributes of the CE affecting its biotic attributes; b) key external drivers – aspects of the “landscape context” –affecting the biotic and abiotic attributes of the CE; and c) the causal linkages among them. The key aquatic attributes and drivers identified through these models will be combined with the element occurrence ranking criteria for riparian and wetland CEs to produce integrated lists of criteria for combined aquatic-riparian-wetland CEs. Although development of such ecological integrity models for aquatic-riparian-wetland CEs will take place during Phase I, Task 3, we have framed informal, preliminary versions to guide identification of data with which to assess the biotic condition, abiotic condition, and the status of critical aspects of landscape context for the strictly aquatic components of combined aquatic-riparian-wetland CEs.

Specifically, we have identified sources for data on:

- Biotic condition: aquatic bioassessment data from federal and state monitoring programs; and data on native aquatic species distributions and aquatic non-native (nuisance) species distributions (see Invasives CA discussion below).
- Abiotic condition: data on the proportion of annual stream flow resulting from groundwater discharge (baseflow); the spatial extent of perennial versus intermittent flow; water quality; the distribution of dams; and habitat quality.
- Landscape context: data on near-stream and watershed land use (see discussion of Landscape Condition for terrestrial CEs, above), water use in the surrounding surface watershed and

contributing groundwater zone, atmospheric deposition of N (nitrogen), a representative potential acidification agent as well as a nutrient) and Hg (mercury), a representative potential bioaccumulative pollutant). To support the analysis of landscape context, we have also identified sources of data with which to identify the basin fill aquifers potentially responsible for sustaining base flow or base water elevations in aquatic CEs, and the watershed zones within each HUC potentially most responsible for generating surface runoff to streams and recharge to basin fill aquifers.

#### **Additional dataset for assessing aquatic coarse-filter**

Desert Research Institute Springs Ecosystems database: Dr. Don Sada of DRI has collected data from more than 2000 springs in the desert southwest including BLM's Mojave and Central Basin and Range ecoregions. This database includes endemic and invasive macroinvertebrate and fish locations and environmental variables associated with these taxa. Many of the springs have never been sampled or the historic data are outdated. Dr. Sada is willing to compile most of the data into a useable format for BLM and NatureServe pending funding. NatureServe has contacted Dr. Sada and asked him for a one to two page summary of his database, the amount of funding he is requesting and an estimated delivery date. We will provide this information to the NOC when we receive it from Dr. Sada.

#### **CE Class V: Aquatic Fine Filter**

Similarly referenced above under the terrestrial fine-filter, Natural Heritage Programs from this ecoregion maintain several thousand location records derived from field surveys over recent decades. These data include field 'observations' and 'element occurrences' of aquatic species (fish and aquatic invertebrate) populations; the latter resulting from a systematic processing of 'observations' into standardized representations that consider distances separating each observation. A total of 2,192 records currently exist for our draft list of aquatic species CEs within this ecoregion (Appendix III). Critical habitat designations from the Fish and Wildlife Service include 10 fish species from the CBR. EcoAnalysts Inc., (included on our consultant team) has conducted taxonomic identification of aquatic macro invertebrates, including natives and invasives, for hundreds of projects and hundreds of clients in the Western USA. State Game and Fish agencies also should have additional location and habitat data for aquatic species of concern to the REA. We will explore their availability within the context of discussions with the WGA-sponsored Southwest DSS effort. Additionally, EcoAnalysts Inc. has conducted taxonomic identification of aquatic macro invertebrates, including natives and invasives, for hundreds of projects and hundreds of clients in the Western USA. With additional refinement of our assessment approach, we may pursue additional data acquisition from this source.

Ecological integrity assessment for the aquatic fine-filter will be subsumed within the analysis of the aquatic coarse filter. Those data sets were reviewed in the previous section.

### **Data Evaluation Results for CAs**

Data sets evaluated and results for CAs are detailed in Appendix IV. Here we summarize our evaluation and results by CA Class. All of the described data sets below are proposed for use in the REA following our evaluation unless otherwise described.

#### **Class I: Wildfire**

We identified and evaluated LANDFIRE's ([www.LANDFIRE.gov](http://www.LANDFIRE.gov)) geospatial layers and data products to represent this CA class. We conclude that LANDFIRE is suitable for the REA purposes. LANDFIRE products describe existing vegetation composition and structure, potential vegetation,

surface and canopy fuel characteristics, simulated historical fire regimes, and current departure from simulated historical vegetation conditions. LANDFIRE data sets and models are based on peer-reviewed science and create consistent and comprehensive fire-ecology products that are standardized across the entire United States. LANDFIRE data products consist of over 50 spatial data layers in the form of maps and other data that support a range of land management analysis and modeling.

Specific data layer products within the database include:

***Fire Regime Condition Class***

Fire regime condition class (FRCC) is a discrete metric that quantifies the amount that current vegetation has departed from the simulated historical vegetation reference conditions. We have noted discrepancies in FRCC map products along map zone boundaries. These result from application of models with conditions within map zones (i.e., the land area across the boundary is ‘unknown’ to the model). During Task 3 we will investigate options to address this issue.

***Fire Regime Condition Class departure***

The (FRCC) Departure Index data product uses a range from 0 to 100 to depict the degree to which current vegetation has departed from simulated historical vegetation reference conditions. FRCC departure reflects changes in community structure and fire frequency and severity.

***Mean Fire Return Interval***

Mean Fire Return Interval layer quantifies the average period between fires under the presumed historical fire regime. This frequency is derived from vegetation and disturbance dynamics simulations using LANDSUM.

***Percent of all fires that are low severity***

The Percent of Low-severity Fire layer quantifies the amount of low-severity fires relative to mixed- and replacement-severity fires under the presumed historical fire regime. These data are critical for parameterizing VDDT state-and-transition models. We have noted concern over burn severity map outputs and will review each layer in detail during Task 3 methods testing.

***Percent of all fires are stand replacement severity***

The Percent of Replacement-severity Fire layer quantifies the amount of replacement-severity fires relative to low- and mixed-severity fires under the presumed historical fire regime. These data are critical for parameterizing VDDT state-and-transition models.

***Percent of all fires that are mixed severity***

The Percent of Mixed-severity Fire layer quantifies the amount of mixed-severity fires relative to low- and replacement-severity fires under the presumed historical fire regime. These data are critical for parameterizing VDDT state-and-transition models.

***Environmental Site Potential***

The LANDFIRE Environmental Site Potential (ESP) layer represents the vegetation that could be supported at a given site based on the biophysical environment.

***Biophysical Settings***

The Biophysical Settings (BpS) layer represents the vegetation that may have been dominant on the landscape prior to Euro-American settlement and is based on both the current biophysical environment and an approximation of the historical disturbance regime. Some have noted apparent inconsistencies within this layer. During Task 3 we will a) integrate available Ecological Site Descriptions with coarse filter CE conceptual models, and b) investigate options for improvement of BpS map layers where apparent error is identified.

***Existing Vegetation***

The Existing Vegetation Type (EVT) layer represents the vegetation currently present at a given site. These data are classified using the NatureServe terrestrial ecological system classification taxonomy. This layer map only be used for portions CA, in combination with other data layers and with additional edits applied to error-prone areas.

***LANDFIRE National Vegetation Dynamics Development Tool (VDDT) models***

This data library provides access to quantitative state-and-transition models for each mapped BPS. Outputs from these models were used to produce the BPS, FRCC departure, and other modeled data layers distributed by LANDFIRE.

#### ***LANDFIRE Rapid Assessment VDDT models***

These models were created to support the LANDFIRE rapid Assessment. This rapid assessment was superseded by the National LANDFIRE Assessment. However, these models are useful for understanding the dynamics of larger areas, and the common dynamics of similar community types.

#### ***The Nature Conservancy's VDDT models***

The Nature Conservancy (TNC) offices in Nevada and Idaho have created a suite of VDDT models that reflect current vegetation. When appropriate, these models are built upon the foundation of the LANDFIRE models with the addition of current (typically anthropogenic) vegetative states and changes in disturbance regimes. When available, these will form a foundation for the VDDT modeling in this effort.

#### ***Monitoring Trends in Burn Severity ([www.mtbs.gov](http://www.mtbs.gov))***

The monitoring trends in burn severity database provides maps of the burn severity and perimeters of all wildfires across all lands in the US for the period spanning 1984-2010. The MTBS is a multi-agency project to track trends in wildfire frequency, size, and severity. We also noted during AMT discussion that Lanfire EVT and EV Height and EV Cover reflect early 2000s, time periods. More recent wildfires can be depicted from burn perimeter data to update these layers. This is an ongoing effort of Landfire “refresh” but we will investigate status of these updates during Task 3.

#### ***The Fire Effects Information System ([www.feic.gov](http://www.feic.gov))***

The FEIC is a compendium of research reports and other publications relating the effects of fire on native plant and animal species, invasive species, ecological communities, and soils. The FEIC is a useful source for understanding fire effects on biodiversity, and for identifying parameter values for VDDT models.

### **Class II: Development**

At the AMT meeting it was decided to incorporate development in scenarios representing current (including development applications submitted to BLM as of May 1, 2011), intermediate (2025 including planned and high potential energy development and infrastructure), and 2060 (climate changes). The 2025 scenario will include all current development and the 2060 scenario will include development in the previous two scenarios.

Several data sets have been identified and evaluated to represent the development CA class. This class contains a large number of subclasses which are further detailed below. Two data sets were evaluated for summarizing overall human modification of the landscape. While we intend to spatially represent each CA class and subclass individually, we will explore in Task 3 the utility of using a summary index of human modification for a more generalized assessment or component of ecological integrity modeling. Both model the influence of anthropogenic disturbance in the in the CBR but were developed at broader scales.

The first data set the Human Footprint (Leu et al. 2008) was developed by the USGS Snake River Field Station. The map focuses on shrubland ecosystems and combines models of habitat use by predators (ravens, crows) closely associated with human presence and the risk of invasive plant infestation (also closely associated with human presence) to estimate the total influence of human activities.

The second data set is the Natural Landscapes (NL) (Theobald 2010). NL is a multi-scale, integrated metric that incorporates national data sets on land cover, housing density, road existence, and highway traffic volume to measure the dynamics of natural landscapes in the conterminous U.S. The advantage of this is metric is that it provides a simple, robust measure of landscape dynamics that has a direct physical

interpretation related to the proportion of natural habitat affected at a location. In addition it represents landscapes as a gradient of conditions rather than a predicated patch/matrix definition. Furthermore it measures the spatial context of natural areas, incorporates land conversion, residential use, transportation infrastructure (including traffic or use), and resource extraction activities.

The NL metric is similar to other approaches that evaluate the effect of humans on natural landscapes such as the Human Footprint (Leu et al. 2008) in that it uses surrogate spatial data on land cover, population, and roads, as well as relying on heuristically derived estimates of human-dominated cover types. NL differs in that it is a simpler metric that has a direct physical interpretation related to proportion of natural cover at a location, examines the broader, landscape-scale pattern to differentiate the spatial context, and assumes that impacts decline continuously as a function of distance, rather than using abrupt “distance bands” or “buffers.” NL also does not rely on pre-established critical scales and avoids the persistent problem of the arbitrariness of defining a patch. As such, this latter database is recommended as a reference for human disturbance caused by development. We acknowledge concerns expressed by USGS reviewers about the NL dataset and will further address those in Task 3.

## **Urbanization**

The Integrated Climate Land Use System (ICLUS) project has developed national scenarios of housing density that are logically consistent with IPCC emissions storylines. It uses a cohort-component methodology to represent population growth in the U.S. Spatial allocation is accomplished using SERGoM (Theobald 2005), a hierarchical (national to state to county), deterministic model that calculates the number of additional housing units needed in each county to meet the demand specified by population projections from the demographic model, based on the ratio of housing units to population (downscaled from census tract to block).

Housing units are spatially allocated within a county in response to the spatial pattern of land ownership, previous growth patterns, and travel time accessibility. The model is dynamic in that as new urban core areas emerge, the model re-calculates travel time from these areas. SERGoM was created using refined land ownership, transportation, and groundwater well density using 2009 data, and by weighting housing units by NLCD 2001 cover types (US EPA 2009; Bierwagen et al. in press).

Other data sets that are suggested for urban development include SILVIS housing density and LANDSCAN, but these are not based on open source demographic/population projections and do not include the detailed spatial data on land ownership, accessibility, and groundwater density to allocate housing units. For these reasons we only evaluated the ICLUS/SERGoM layer which we determined is adequate for use in the REA.

## **Infrastructure**

### ***Roads***

The NOC is preparing a new product, the “linear disturbance” map that was developed at the BLM field office level. Data managers at the NOC have indicated that this will be the most detailed data set of roads and will be ready early in 2011. We recommend this data set in place of the 2009 Tiger/Line shapefiles and National Transportation Atlas Database (NTAB). ***Pipelines***

The NOC has indicated that the BLM Linear Disturbance maps may contain pipelines at a fine scale. However if there are discrepancies or gaps in the data set, NatureServe recommends an augmented National Pipeline Mapping System (NPMS) data set which is available on the SAGEMAP web site. This data set includes all major gas and hazardous liquid transmission for the CBR. We have reviewed the FGDC compliant metadata for this data set and recommend it for use. Geospatial data regarding future pipelines in the CBR such as the Ruby Pipeline Project have been requested but not yet obtained.

### ***Transmission lines***

Transmission lines are another component of the BLM Linear Disturbance Maps. This data set will be fully evaluated for completeness and accuracy upon receipt. Other transmission lines data sets consist

of market significant transmission lines. However useful, this layer unfortunately lacks smaller branch transmission lines that represent the bulk of transmission lines on the landscape. If the BLM Linear Disturbance maps lack this component, we will obtain more data sets from USGS SAGEMAP for review. Point locations of communications towers have been obtained and will be considered as a part of the transmission infrastructure. Extensive improvements to the electrical grid are planned for the CBR to accommodate new renewable energy projects. To represent these changes we propose using the Section 368 corridor maps provided by West-wide Energy Corridor Programmatic EIS (DOE & BLM 2008). We will continue to examine other energy corridors relating to specific renewable energy sectors as we identify and evaluate them. The project PI, Crist, is a member of the SPSG Environmental Data Task Force (EDTF) and will use this connection to obtain data if possible on planned transmission corridors should they come available during the course of the project.

#### ***Water transmission***

The USGS NHDplus layer has specific categories identifying canals, ditches and other artificial paths used for water transmission. Querying this data set will create an adequate water transmission layer.

#### ***Railroads***

Railroad networks are less spatially and thematically complex than roads. We recommend using the railroads layer from the NTAB for the ecoregion if this information is not included in the BLM Linear Disturbance Maps.

## **Energy development**

### ***Renewable Energy Development***

#### **Wind**

The BLM provided maps of pending, authorized and closed wind leases for the CBR. Also provided were the annual average wind resource potential maps at 50m height for the states of the CBR (NREL 1986). Produced by NREL, this data set from 1986 is being replaced by a high resolution wind resource map showing the predicted mean annual wind speeds at 80m height (AWS Truewind 2010). This new data set presents the most accurate picture of wind resource potential for the region. We have recently requested this data but it has not been received in time for this evaluation.

#### **Solar**

Pending and closed solar energy leases for the CBR were provided by the BLM. We recommend using the Solar Energy Study Areas that identify areas currently being evaluated in the Solar Energy PEIS (ANL 2009). Also available are solar energy resource maps which show direct normal solar radiation for areas of 1% and 3% slope (SUNY & NREL 2007). These will provide some indication of the areas most likely to be developed for solar energy, especially concentrated solar power facilities. The Solar PEIS shows the areas most likely to be developed in the short term.

#### **Geothermal**

We obtained from the BLM maps of producing and non-producing geothermal leases as well as a potential geothermal energy layer. From the Great Basin Center for Geothermal Energy we obtained operating geothermal plants and the map of Geothermal Favorability and Exploration Activity (Zehner et al 2009). These data sets adequately show current and future siting of geothermal generators.

#### **Biomass**

We reviewed the national biomass potential maps and found them to be too coarse for use in the REA. The AMT indicated there is significant interest in CBR to harvest Pinyon-Juniper (PJ) areas for biomass and as a restoration tool. The AMT also manifested a concern that biomass harvests may target old-growth PJ forests. Readily available data are inadequate to represent the potential biomass locations (current and future expansion beyond old growth) however we will continue to evaluate this as a potential change agent through proposed modeling to be assessed in Task 3.

#### ***Extractive energy development (oil, gas)***

The BLM provided maps of oil and gas leases and agreements for the CBR. Communication with the NOC has indicated that detailed oil and gas maps detailing well locations are pending. The EPCA Phase III Inventory GIS data files (DOI et al 2008) are recommended for evaluating areas likely to be impacted in the future by further extractive energy development. The detailed oil and gas maps and the EPCA will sufficiently depict the extent of this activity.

## **Hydrologic Change Agents**

### ***Groundwater withdrawals***

Data on current intensities of groundwater withdrawal within the CBR will be assembled from data developed by the USGS for its Southwest Principal Aquifers (SWPA) study (McKinney & Anning 2009), specifically data on municipal and agricultural withdrawals. Projections of future intensities will build on the results of the assessment of future development, incorporating present estimates of the rates of municipal per-capita and agricultural per-acre consumptive use.

### ***Altered Surface Flow Connectivity***

Data on present surface flow connectivity within the CBR will be assembled from the U.S. Army Corps of Engineers National Inventory of Dams (NID), the download for which needs to be carried out by a governmental agency. At present we have no strong basis for projecting future dam distributions, but in general the construction of dams is strongly disfavored at both the state and federal levels; if any changes take place in dam distribution they will likely involve the removal of dams, particularly ones with high hazard ratings. We will assess the changes to flow connectivity by examining the consequences of removing high-hazard dams, as identified in the NID.

### ***Altered Surface Flow***

Surface flow change can result either from human withdrawals and return flows, or from the results of climate change. Since all surface water rights are fully appropriated in the CBR, as they are throughout the arid west, we do not forecast changes in surface water withdrawals or return flows. As noted above, we will use the projections of future development as the basis for projecting future water demand for the CBR, and estimate the extent to which any increases in demand could be met through either surface or groundwater resources. We will carry out a separate assessment of the likely changes in surface hydrology (and groundwater recharge) resulting from climate change, as discussed elsewhere in this memo.

## **Mining**

The BLM provided maps of solid mineral leases for the CBR. We also acquired a data set from the USGS Mineral Resource Data System (MRDS) of all mine sites and mine processing facilities for the ecoregion. These same state entities can also inform us of which mine and quarries are currently active in the state, an important factor that the MRDS does not report. The MRDS is largely derived from 7.5 minute USGS quadrangles, however, it is comprised of point data which does not reflect the surface disturbance spatial extent. We will need to identify another data set or model surface disturbance if we intend to identify the total surface footprint of mines and their supporting infrastructure. In Task 3 we will explore modeling the footprint by associating the point locations to “barren” land cover classes from 30 m land cover data or using them to derive a relative mining impact layer. Large active mines (e.g. open pit) mines may be detected with existing satellite derived LU/LC maps.

## **Military use/expansion areas**

Geospatial data pertaining to impacts or management of natural resources on military reservations was not readily available for the CBR. Heavily disturbed areas on military reservations will likely need to be extracted from general land use/land cover maps such as the National Landcover Data set (NLCD)

or NatureServe ecological systems map. Currently there are no publicly announced plans to expand any military bases in the CBR so expansion boundaries are unavailable. The NOC has indicated that they have three military flight data sets (from the FAA): no-fly zones, low flying areas and flight paths. These three layers may approximate areas of elevated noise from aircraft and serve to identify incompatible use areas, specifically areas where the DOD may object to the development of wind turbines. However the correlation between these designated flight zones and disruptive elevated noise levels is somewhat tenuous. The AMT indicated that their primary concern are the flight and training areas incompatibility with renewable energy development. In task three we will continue to look for military data by reaching out to a contact at Fallon AFB and the DoD-led Western Regional Partnership. .

### **Air quality impacts (non attainment areas and dust)**

We will use National Atmospheric Deposition Program (NADP) data on Nitrogen as a stand-in for all air pollutants that involve acid deposition and result in nutrient enrichment once buffered. We will use NHDPlus and USGS-Nitrogen Groundwater Risk data sets as cross-checks on the NADP regional estimates. We will use NADP data on Mercury as a stand-in for all air pollutants that can bio-accumulate and cause physiological or developmental harm.

### **Recreation (OHV use, other intensive recreation, land sales, etc.)**

We recommend using modeled estimates of dispersed recreational use via a method documented in *Network and Accessibility Methods to Estimate the Human Use of Ecosystems* (Theobald 2008). This approach will be thoroughly reviewed and evaluated in Task 3 as well as data from BLM on recreation sites and managed areas. Pending review of these data sets we will provide a recommendation of the extent to which we can incorporate effects of site-based recreation. We evaluated the US Forest Service National Visitor Use Monitoring data set and determined that because there is no comparable data set on BLM, NPS, USFWS and other public lands -- these data are not suitable to be used in the REA.

### **Refuse Management (landfills, sewage sludge disposal, nuclear disposal, etc.)**

From the USGS SAGEMAP site, we obtained the locations of landfills and waste transfer stations in 11 western states. Data was obtained from state and federal agencies in GIS, tabular, and map format. The data is in point format which leaves us without the spatial extent of landfills. This has created a similar situation identified with the mine resource data- a lack of a total footprint area for each feature and likewise we will investigate modeling potential to represent this CA. Data for mining slurry lagoons has also been obtained from the NV Dept of Environmental Protection. Similar data has not been obtained yet from Utah or California. Data regarding sewage sludge disposal, nuclear disposal, etc. have not been identified.

### **Agriculture**

#### ***Crops, orchards, irrigated pasture***

A useful resource for evaluating agriculture at a fine scale is the USDA Common Land Unit, the smallest unit of land that has a permanent, contiguous boundary, a common land cover and land management, a common owner and a common producer in agricultural land associated with USDA farm program. However the Food, Conservation, and Energy Act of 2008 restricts access to this information to certain departments of the USDA. The alternative is the 2007 Agricultural Census of the United States (USDA 2007) which is only spatially explicit down to the county level or 1:21,000,000 which is too coarse for the REA. We recommend that agricultural areas be identified through an existing raster data set such as NatureServe's ecological systems map which identifies these areas with a sufficiently high level of accuracy and precision.

## **Exotic ungulate grazing**

### ***Wild Horses and Burros***

Data exist to answer most of the Management Questions posed at a relatively coarse scale. Although spatial data for Herd Areas (HA) and Herd Management Areas (HMA) boundaries are believed to be accurate, tabular data on wild horse and burro numbers are presented on an HMA or HA basis. In Nevada these areas range from 4,000 to more than 1,000,000 acres. Tabular data on wild horses and burros include numbers of each by HMAs and HAs for each FY from 2005-2009. The AMT recommended against using the tabular data due to concerns about the accuracy of this information. After discussing this CA with the AMT, we clarified that few assumptions can be made using the data as-is to answer questions about whether units are exceeding AUM. Instead we will answer questions about the location and the likely integrity changes to HAs and HMAs using them as assessment and reporting units.

### ***Livestock***

Spatial data provided by BLM for allotment and pasture (pastures are areas within allotments) boundaries are believed to be the most accurate available. Tabular data on livestock Animal Unit Months (AUMs) and season of use are being assembled from the Rangeland Administration System (RAS) by the NOC. NOC indicated that it is revising and quality-testing this data and that only some livestock data will become available in an appropriate time frame (species of grazer (e.g. sheep or cattle) and permitted AUMs at each allotment).

Drs. David Pyke and Cam Aldridge of the USGS are currently leading an effort to improve accuracy of the BLM allotment data in the Western US. This data may be available to incorporate early in 2011 and we will conduct a rapid evaluation of them if received in time.

Authorized use data adjusted for actual use, spatial and temporal variation, and monitoring data would be at a coarse scale. The effects of livestock would need to be analyzed over the extent of the allotments boundaries, which range up to thousands of acres in size. Authorized use data could be adjusted for 1) actual use based on billing records for each allotment, 2) spatial and temporal distribution within allotments based on textual information contained in ten-year and annual grazing permits and permit decisions, and 3) actual use based on monitoring data. This information would need to be assembled from BLM field offices and is beyond the REA scope.

As with wild horses and burros, the current data is insufficient to draw conclusions about appropriate AUM so we will likewise treat grazing allotments as assessment and reporting units only.

The AMT expressed interest in artificial water source locations for stock and wildlife. The field offices are currently gathering this data in the field but it will not be available during the REA timeframe. Ecoregion-wide data on illegal grazing on allotments or feral cattle grazing are unavailable.

## **Class III: Invasives**

### **Terrestrial Invasive Species**

Comprehensive mapped data on terrestrial invasive species are non-existent for the ecoregion. Given the diversity and abundance of weeds in this ecoregion, this is no great surprise. We do anticipate organizing weed species as assemblages, i.e., annual grasses, perennial grasses and forbs, etc.; in order to amass sufficient sample data for modeling of units that are meaningful for addressing management questions,

We have located a few data sets that cover a small area (e.g., single county) for many species, and a few data sets that cover a larger area for single species (namely cheatgrass and tamarisk). For covering cheatgrass (*Bromus tectorum*) we have three sources. The Annual Grass Index (Peterson 2006) used Landsat data from 2004 for Northern Nevada (which was based on training data and is predominately *Bromus tectorum* but also included *Bromus arvensis*, *Poa bulbosa*, *Taeniatherum caput-medusae*, *Vulpia*

*microstachys*, and *Vulpia octoflora*). The *Bromus tectorum* Estimated Percent Cover Model (Peterson 2003) estimated cover from satellite imagery in April and June 2001. In addition we have the 2,325 survey points of *Bromus tectorum* presence/absence from 2004 & 2006 (Bradley and Mustard 2006). We will seek out the weed data set being developed by Doug Ramsey at UT State University identified by the AMT.

The Southwest Exotic Plant Mapping Program (SWEMP) has >11,000 records that coincide with the Terrestrial Invasive Species change agent list for the CBR. These are point location data that need further data quality evaluation in order to evaluate fully.

SWEMP	
Count	Scientific_Name
9092	<i>Bromus tectorum</i>
731	<i>Elaeagnus angustifolia</i>
997	<i>Halogeton glomeratus</i>
347	<i>Salsola kali</i>
260	<i>Sisymbrium altissimum</i>

We still lack ecoregion wide data for *Taeniantherum caput-medusae*, *Acroptilon* spp., *Centaurea* spp., *Nasturtium officinale*, *Cirsium* spp and *Caduus* spp. The NOC recently provided a weed infestation map with 30,952 polygon locations in the CBR. This layer is currently undergoing evaluation but certainly provides a valuable resource. The Nevada and Arizona natural heritage programs have also provided exotic species data.

For *Tamarix* we have the Colorado River Basin Tamarisk and Russian Olive Assessment data (Tamarisk Coalition 2009). This database is a compilation of many sources. However very little, if any of this data set coincides with the Central Basin Ecoregional boundaries. We don't have a source of mapped data of *Tamarix* for the Central Basin.

Additional sources of data on invasive species locations occur in the SWReGAP, CA ReGAP and LANDFIRE reference sample databases. These sources have geo-referenced points that include exotic species such as tamarisk (*Tamarix*), Russian olive (*Elaeagnus angustifolia*), cheatgrass (*Bromus tectorum*), peppergrass (*Lepidium*) and others. In addition, these data contain geo-referenced points representing invasive species assemblages, for example "Introduced Riparian Vegetation" and "Exotic Annual Grassland." The geographic extent and abundance of exotic species point locations in these databases needs to be evaluated. These data sources are also listed as data sources in the CE section (above), and can be located in the list of CE data sources, Appendix I.

We searched for mapped invasive species data from the Extension Service with University of Nevada, University of Las Vegas, Utah State University, and the University of Arizona to no avail. State weed councils (CA, AZ, NV, and UT) have abundant information defining noxious or invasive plants and status ranks (how aggressively "invasive" a species may be). But specific location and mapped data was not available through these sources.

A component of the USGS Human Footprint map (Lue et al 2008) includes an exotic plant invasion risk model that predicted the potential spread of exotic plants according to anthropogenic features. This will be evaluated as a potential resource for modeling potential spread in Task 3.

### Aquatic Invasive Species

Unlike many ecological and environmental databases (e.g. real-time weather data); current, complete, and verifiable site location databases of aquatic invasive species are dependent on timely observation and reporting by qualified biologists or taxonomists. There are often large lag times between when a private citizen, researcher, or manager observes an aquatic invasive species, when it is reported to

the appropriate agency, and when it is verified and entered into a useable database. There are also large differences in observational and survey effort between water- body types. Invasive species are more likely to be reported and monitored in easily accessible or popular fisheries than in other locations. Isolated remote small springs/seeps are seldom visited; unless they provide known habitat for a listed native species. In such remote springs/seeps, an invasive species could go unreported for many years. Detectability and recognition of invasive species is also problematic. Most private citizens and many biologists are unfamiliar with invasive species identification or may assume that an invasive species is native. Invasive species may also be cryptic, highly evasive, or occur at low or undetectable densities, further reducing timely verification and reporting. Finally, monetary funding for surveys and compiling databases of invasive species is lacking.

Given these restraints, the most comprehensive available databases on aquatic invasive species are maintained by the USGS Nonindigenous Aquatic Species Program. These databases contain high quality point or HUC locations and brief descriptions for the majority of invasive species that were selected as representative change agents (CA) in the ecoregion. However, the USGS NAS databases are not exhaustive and additional databases have been selected to complement or supplement these databases. These include the Montana State University New Zealand Mudsail in the Western USA database (which includes reported locations in our ecoregion) and a USGS Fort Collins, CO database containing locations of known *Didymosphenia geminata* (didymo) locations in our ecoregion. More recent up- to- date didymo location data have been requested from USGS at Fort Collins and is pending. In addition, we are in the process of obtaining an extremely useful database from the Desert Research Institute, Reno, NV. This database contains high quality information on invasive and native species and environmental data from well over 2000 springs in the southwestern USA, including springs from our ecoregion. Historic ecological data from many of these springs that are now in the Desert Research Institute database are were highly outdated or have never been reported. Also, most of the state fish and game departments and state and federal water quality monitoring program databases that we are using in the aquatic Conservation Elements analysis also contain data on aquatic invasive species. For example, US EPA National Lake Assessment, Nevada Division of Environmental Protection Bureau of Water Quality Planning, and Utah State University-Western Center for Monitoring & Assessment of Freshwater Ecosystems databases are being utilized in the aquatic Conservation Elements analysis and will also be used in our aquatic invasive species Change Agent analysis. However, state and federal water quality data sampling and collection methods are not specifically focused on invasive species and may overlook or under represent invasive species locations.

#### **Additional datasets for assessing aquatic invasives**

Desert Research Institute Springs Ecosystems database: Dr. Don Sada of DRI has collected data from more than 2000 springs in the desert southwest including BLM's Mojave and Central Basin and Range ecoregions. This database includes endemic and invasive macroinvertebrate and fish locations and environmental variables associated with these taxa. Many of the springs have never been sampled or the historic data are outdated. Dr. Sada is willing to compile most of the data into a useable format for BLM and NatureServe pending funding. NatureServe has contacted Dr. Sada and asked him for a one to two page summary of his database, the amount of funding he is requesting and an estimated delivery date. We will provide this information to the NOC when we receive it from Dr. Sada.

USGS *Didymosphenia geminata* (didymo) database: Dr. Sarah Spaulding, USGS, Fort Collins, CO, has provided NatureServe and BLM with known locations of didymo in the western USA, as of 2007. She also has additional didymo locations that have been reported since 2007 but the data is not in a useable format. Since didymo is rapidly spreading throughout the western USA, including BLM's Mojave and Central Basin and Range ecoregions, the acquisition of the most recent data is critical in order to evaluate its spread. Dr. Spaulding is willing to compile the most recent data into a useable format for BLM and NatureServe pending funding. NatureServe has contacted Dr. Spaulding and asked her for a one to two page summary of her database, the amount of funding she is requesting, and an

estimated delivery date. We will provide this information to the NOC when we receive it from Dr. Spaulding.

#### **Class IV: Climate change**

Datasets available for climate change effects modeling are divided into two categories: current and future time periods. The BLM recommended dataset for analyzing current climatic patterns is the PRISM dataset (Daly et al 2002), which is widely recognized as the most accurate spatial climate dataset available within the domain of the conterminous U.S. PRISM is currently the official climatology of the U.S. Dept. of Agriculture. Future climate change effects will be modeled using dynamically downscaled model outputs generated by the USGS (which we were not required to evaluate for Task 2). Factors in these current and future spatial climate datasets relevant to the objectives of the CBR REA include the spatial resolution of the available data, the temporal extent of available records to analyze recent historical climatic variability, the climate parameters available in the current as compared to the future, the temporal resolution with which these current and future climate variables have been measured (i.e., daily, monthly, etc.), and the degree of uncertainty that remains based on the limited number of future climate datasets available for climate change effects modeling.

PRISM is available at several spatial resolutions. The finest resolution for the freely available PRISM dataset is a 2.5 min grid (4km). While higher resolution spatial climate information is always desirable given that plants and animals interact with climate at relatively fine spatial scales, the 4km spatial scale of the PRISM data is an appropriate resolution for the suite of climate change effects analyses that will be conducted for the CBR REA (see addendum below). The future climate models to be obtained from the USGS are produced at 15km grid resolution – significantly coarser than the 4km PRISM data for current climate. It must be recognized that any current climate analyses or current species distribution models will be produced at a finer resolution than equivalent future analyses, as an unavoidable limitation of the coarser resolution climate model outputs. This tension between the finer resolution of current climate datasets and the coarser resolution of global climate models is longstanding, and it will likely be several decades before global or even regional climate models can produce native outputs at sufficiently fine spatial resolution for detailed ecological impacts analyses. The availability of multiple dynamically downscaled climate model outputs for impacts analyses is unprecedented, and it represents a huge advance over the far more prevalent and simplistic statistically down-scaled climate model outputs currently in use for most all ecological forecasting efforts.

The first task in the CBR REA climate change effects modeling is to quantify observed historical climatic variability for the distribution of major CEs across the ecoregion. The results of this analysis will be highly dependent on the availability of long term reliable climatic records. In this regard, the PRISM dataset is highly appropriate and will perform well (see addendum below). PRISM currently offers monthly climatic variables dating back to 1895 that have been vetted and published (Daly et al 2002; Gibson et al 2002). This represent a time series sufficiently long to capture climatic variability caused by decadal-range oscillation patterns such as ENSO (El Nino – La Nina cycles).

There is a significant difference in the climate parameters available in the PRISM dataset for the present, and the very large number of climate and biophysical variables archived from the dynamically downscaled climate model outputs. PRISM offers only monthly measures of: maximum temperature, minimum temperature, total precipitation, and dew point temperatures. Over 70 climatic and biophysical variables recorded at 3 hourly and 6 hourly intervals are produced by the dynamical models generated by the USGS. As of this writing, we have received significant new information about the climate model dataset that will help determine the specific variables we will request for climate change effects modeling. At a minimum, the same four parameters available in PRISM averaged into monthly time steps will be requested. In addition, the 3-hourly and 6-hourly model outputs will allow an analysis of extreme temperature and precipitation events as produced by the regional climate model outputs. Extreme events

are very important drivers of climate impacts on plants and animals. We are currently contributing to a multi-ERA working group to establish a clear path forward on this front.

Finally, the evaluation of the degree of uncertainty in climate-driven future ecological impacts is limited by the number of available climate model outputs. At this stage, we know we will have 3 independently generated climate model runs. Each run represents 3 global circulation models that are used as boundary conditions to feed a single regional climate model, all of which are run under a single scenario of future greenhouse gas emissions. While, as stated earlier, it is unprecedented to be conducting ecological impacts analyses with multiple dynamically downscaled climate model outputs, it must be recognized that the data still represent a relatively small sampling of future climate space. The degree of uncertainty can be evaluated by the degree of model agreement across these 3 independent model outputs, but in cases where there may be relatively little agreement, climate driven ecological impacts may remain somewhat uncertain.

### **Alternative Climate Data for Climate Change Effects Assessment**

Measuring impacts of climatic change requires an understanding of the current climates to which target conservation elements are adapted. Weather station data providing specific measurements of localized climates has only been available for about the last century, and only in the last 50 years or so has the density and quality of weather station data been sufficient to produce region-specific 'normal' climatologies. The PRISM group at Oregon State University has generated decadal averages monthly temperature and precipitation for the conterminous U.S. from the 1890's to the present day at a resolution of 4km. These are the BLM-recommended climate data for characterizing current climates in the ecoregions under assessment.

However, the influence of climate change on species distributions and interactions is likely to be mediated by microclimatic patterns, just as the fine scale patterns of current species distributions are also strongly influenced by microclimate. Climate data that more accurately reflect microclimatic features such as cool air drainage down valleys, or temperature and precipitation differences on north vs. south facing slopes, will offer a better understanding of how future climatic changes might influence conservation elements.

The PRISM group has generated a time-series climatology for the conterminous U.S. at 800m resolution. Acquisition of this dataset, which would cost roughly \$5000-7000, would allow 1) a characterization of historical climate normals at a spatial scale that more closely approximates how plant and animal species interact with local climates, and 2) the analysis of climate anomalies (also called "departure analysis"), that is, a measure of the magnitude and directionality of climatic changes already observed, at fine spatial scale. In addition, the 800m PRISM dataset is considered a more accurate product, even though it requires additional interpolation. In tests comparing the two climate datasets for the 10 counties around the San Francisco Bay Area, CA, significant errors were encountered in the 4km as compared to the 800m in several mountainous areas (D. Ackerly, personal communication, Nov 2010).

The 800m PRISM data will be particularly useful in conjunction with the future climate model outputs of the Flint and Flint (2007) Basin Characterization Model (see below) that has been driven with the GFDL and PCM climate models under the A2 and B1 emissions scenarios. The Flint and Flint future climate datasets are produced at a spatial resolution of 270m. Together, a characterization of climate norms at 800m and climate futures at 270m would result in climate impacts analyses using the very best possible spatial climate datasets available.

The PRISM 800m spatial climate data is available now for purchase from the PRISM group at Oregon State University. Because the dataset covers the entire lower 48 states, a single purchase will serve climate analysis for all ecoregions outside Alaska and Pacific Islands.

### **Climate-Hydrologic Effects Assessment**

We will also assess the impacts of climate change on aquatic coarse-filter CEs. The USGS has developed data to assess the likely impacts of climate change on the watershed hydrology of large areas

of the southwestern US. This work has been conducted by Flint and Flint, and incorporates the Basin Characterization Model (BCM) methodology that they developed to assess the impacts of historic-to-current climate variation on watershed hydrology (Flint and Flint 2007). They used the 4-km PRISM dataset as their historical reference for precipitation and max and min air temperature. Using these 4-km data, they ran their BCM historically for the interior desert southwest (1940-2007) and for California (1895-2009). The model produces output at a 270-m grid resolution for monthly precipitation, max and min air temperature, potential evapotranspiration, actual evapotranspiration, excess water, snow accumulation and melt, sublimation, soil storage, recharge, runoff, climatic water deficit. This study (Flint and Flint 2007), with its 270-m grid resolution, provides crucial information for the assessment of current condition and ecological integrity for aquatic coarse-filter CEs. Importantly, its fine spatial resolution makes it possible to aggregate the BCM output data effectively by 5- or 6-field HUC and link the BCM output to NHD data layers.

The data from the Flint and Flint (2007) study have already been made publicly available. However, the spatial boundaries for the earlier study do not cover 100% of the CBR and MBR domains. Specifically, its interior southwest (“Great Basin”) spatial domain covers most of CBR except along one portion of the northern periphery; and their “Great Basin” and “California” domains together cover most of the MBR except for the NW corner of Arizona. As a result, one area for potential investment of resources would be to support incorporation of these orphan areas into the master domains. Additionally, we would like the BLM to be aware that the Flint and Flint team could run the same analyses on other ecoregions.

The Flint and Flint team further, as noted above, has applied their BCM methodology to climate data downscaled to the same 270-m grid resolution. Lorraine Flint (personal communication 10/15/10) has stated that their team has downscaled and bias-corrected four climate future scenarios, using the PCM and GFDL climate models coupled to the A2 and B1 greenhouse gas emissions scenarios for the continental US, resulting in precipitation and maximum and minimum air temperatures again at a 4-km grid resolution. The Flints have used their BCM model to calculate local climate and watershed condition variables at the 270-m scale for the same Great Basin and California domains, again resulting in monthly estimates for each parameter. Unfortunately the USGS presently does not have funding to complete all four climate futures analyses for their Great Basin; they have so far completed only the GFDL-A2 analysis for this domain, although they have completed all four analyses for California. They presently anticipate having the resulting five output datasets available for public use “by this winter.” They would need funding to run PCM-A2, PCM-B1, and GFDL-B1 for the interior southwest on our REA timeframe; and would also need funding to run both their historic and GFDL-A2 models for the portions of CBR and MBR that their domains do not presently cover.

There are four other factors to consider, with respect to the potential costs versus benefits of having the USGS complete its planned climate futures analyses, and filling in the orphaned areas of CBR and MBR for historic conditions:

1. The only climate future scenario available from the Flint and Flint team for the majority of CBR and similarly for MBR is the GFDL-A2 model. This is only one of many climate future models, and may not produce results consistent with other down-scaled models. The down-scaling methodology used by the Flint and Flint team also is not a dynamic methodology, and so their results may not be fully comparable to those produced by the Hostetler team. On the other hand, the 15-km scale of the Hostetler team output lends itself poorly to aggregation at the HUC scale or linking to NHD data layers, for analysis of aquatic coarse-filter CEs.
2. Any new runs by the Flint and Flint team will result in what the USGS officially designates as “preliminary” data. Such preliminary data cannot be released to the public until fully reviewed by the USGS, but can be released to any client that has actually contracted for the study. So, if

the BLM were the client, the Flints could provide them with the preliminary data in advance of public release, and the BLM would then be able to share the data with us and other REA contractors.

3. There is a fair amount of serious data management involved in working with the USGS BCM output, because the datasets are so huge. The Flint and Flint team is willing to help with that, but it would be better if a client could help with funding.
4. Finally, the Flints have expressed interest in completing their BCM modeling for other ecoregions. Lorraine Flint (personal communication, November 2010) has stated that it would take them “about a month” to do each additional ecoregion that the BLM might request, both the historic and the future scenario runs. This clearly does not affect CBR and MBR, but could be valuable for future REAs. Running the BCM for additional ecoregions under a “client” relationship with the BLM would, as noted above, make the results of these additional runs available more rapidly for REA purposes.

## **Data Evaluation Results for Managed Lands and Sites**

We refer to these classes as *Places*, being neither a CE nor CA; thus the PL abbreviation below.

### **PL Class I: Sites of High Biodiversity**

Areas of High Biodiversity are represented in the data by previous analyses characterizing locations with concentrated at-risk biodiversity or locations where a prioritization exercise has identified areas of high conservation significance. Criteria for previous prioritization exercises vary, and those variations can reflect on their suitable usage for the REA. This class may overlap spatially with the subsequent two PL classes (II and II) but they differ in that the latter categories include established legal boundaries for land and water units (e.g., ACECs). Areas of high biodiversity significance most frequently imply more flexible boundary definition and suggest the need for future field verification prior to settling upon new legal or management designations.

Crucial habitats, as defined through the Western Governor’s Association (WGA) Western States Decision Support System (DSS) efforts, often fall into this category. We have yet to evaluate these data; as they will become available through the Southwest DSS effort. Ecoregional assessments (ERAs) conducted by The Nature Conservancy (TNC) include the identification of priority conservation areas. These “portfolio sites” equate with area of high biodiversity sites. The primary TNC effort for the CBR includes their Great Basin assessment (Nachlinger et al 2001), but adjacent assessments include sites that overlap the CBR boundaries. By compiling information on “coarse-filter” and “fine-filter” CEs, evaluating their condition, establishing representation goals, and factoring in existing protected areas ERAs identified an efficient land allocation to achieve their stated representation goals. NatureServe has acquired the entire U.S. data set from TNC to represent these sites in the REA. **We recommend using these site boundaries as a potential spatial reporting unit for this REA.**

Two additional data sources in this category include Important Bird Areas, identified by Audubon and by the American Bird Conservancy. In many instances, the IBAs were already factored into previous TNC assessments. However, as we acquire these data, we will determine their relative applicability to this REA.

### **PL Class II: Specially Designated Areas of Ecological or Cultural Value**

Many of these areas are special classifications of BLM and US Forest Service lands: wilderness areas, wilderness study areas, and the regions only national conservation area, the Black Rock Desert-High Rock Canyon Emigrant Trails. We will also take into account unique BLM lands distinctions such

as Areas of Critical Environmental Concern (ACEC). By their special nature, USFWS National Wildlife Refuges and National Parks are also included in this category. All of this data is best represented in the Protected Area Database of the U.S. (PADUS) version 1.1 which has been obtained and evaluated. This data set will be verified against the BLM Surface Management Agency (SMA) maps.

**PL Class III: Other Managed Lands**

Other managed lands include the majority of the area of federal or state managed lands in the CBR characterized by management for multiple uses. The AMT has requested that we use BLM’s SMA maps provided by the NOC to identify all managed lands..

**Summary Data Gaps and Recommendations for CEs, CAs, PLs, and MQs**

We summarize the key data gaps and revisions by REA component:

**CE Data Gaps and Recommendations**

It is unlikely that there will be substantial data gaps for Conservation Elements in the REA. As noted throughout the sections above, considerable effort is needed to combine and rectify existing data sets to meet the needs for the REA. At this time we do not recommend any changes to the proposed conservation elements.

**Table 1. Summary of data suitability for CEs.**

<b>Conservation Element Category</b>	<b>Number of Elements</b>	<b>Data Suitable?</b>
Basin Dryland Ecosystems	10	yes
Montane Dryland Ecosystems	7	yes
Basin Wet Ecosystems	6	yes
Montane Wet Ecosystems	3	yes
Nested Terrestrial Habitat-Based Species Assemblages	TBD	high probability
Nested Aquatic Habitat-Based Species Assemblages	TBD	yes
Individual Species	TBD	high probability
<b>Desired Conservation Elements</b>		
Mule Deer		yes
Soils of Conservation Concern (high readability)		high probability

**CA Data Gaps and Recommendations**

Sufficient comprehensive data sets exist to model the Wildfire and Climate Change classes. For other CAs, there are critical data gaps that will require additional research and AMT guidance.

**Table 2. Summary of data suitability for CAs.**

<b>Change Agent Class</b>	<b>Number of Subclasses</b>	<b>Data Suitable?</b>
Wildfire	2	Yes
Development	10	Yes for most subclasses
Invasives	2	Limited but suitable for several
Climate	TBD	Yes/resolution issues

**Managed Lands Data Gaps and Recommendations**

We did not identify any data gaps at this stage but we will need to evaluate the Crucial Habitats data from the Southwest States DSS project when it is available.

**Recommendations for Management Question Revisions**

Treatment of individual management questions (MQs) is described in Appendix IV. Generally, data appears available and suitable to answer most of the MQs. Further acquisition and assessment of data is needed and ability to assess the MQs through models will be treated in Task 3. The table below summarizes MQs or categories of MQs that are or may be impacted by data availability.

**Table 3. Summary of Management Question Implications**

<b>Management Question</b>	<b>Issue/Recommendation</b>
MQs involving climate change	Large differences in climate change data resolution may impact ability to deliver products at desired resolutions
MQs involving exotic grazers	Data on actual distribution of grazers non-existent or inadequate to answer questions requiring such data. Drop MQs related to these parameters.
Of these water resources, what is their surface water/groundwater connectivity?	Not directly measurable at regional scale; we will use surrogates (see memo Appendix IV
What is the natural range of variation in high and low water levels or flows (e.g., frequency, timing, duration of high and low water levels or flows)?	Not directly measurable at regional scale; we will use surrogates (see memo Appendix IV
What is the current distribution of invasive species included as CAs?	Most data are highly localized or state-level and will likely require modeling for many species
Where are areas of planned or potential development (outside of current urban areas)(e.g., under lease, plans of operation, governmental planning), including transmission corridors?	Development plans of private industry not readily available unless already in NEPA process

Management Question	Issue/Recommendation
Where are areas with groundwater resources available to sustain renewable energy projects that would not degrade aquatic ecosystems that also depend on these groundwater resources.	This may be too fine-detailed a question to be answered with a REA. See Mem 2 Appendix IV for details and suggested approach
Where are areas under leases of water rights?	We have not identified a consistent set of data with which to assess the spatial distribution of either surface or ground-water use rights, and will need to clarify with the BLM what is needed here.
Where are artificial water bodies including evaporation ponds, etc.?	Not sure how we would distinguish "artificial" except as impoundments behind dams (US Army Corps NID)
Where are the areas showing ecological effects from existing surface water exploitation?	We have to rely on comparisons of historic <u>published</u> records (rather than GIS data) on the distribution of perennial flows and perennial water levels in springs, to records of their distribution today; we have not identified GIS data layers for this purpose.
MQs dealing with military use and constraints	We will address how available military restricted areas may affect energy development but will not address questions related to military area expansion unless provided.

## References

- AWS Truewind & NREL. 2009. Predicted mean annual wind speeds at 80-m height. AWS Truewind & National Renewable Energy Laboratory.<http://www.windpoweringamerica.gov/index.asp>
- Belnap, J., J.H. Kaltenecker, R. Rosentreter, J. Williams, S. Leonard and D. Eldridge. 2001. Biological Soil Crusts: Ecology and Management. USDI Bureau of Land Management National Science and Technology Center, Tech. Ref. 1730-2.
- Bierwagen, B., D.M. Theobald, C.R. Pyke, A. Choate, P. Groth, J.V. Thomas, and P. Morefield. (In press, accepted 12 October 2010). Land-Use Scenarios: National-Scale Housing-Density Scenarios Consistent with Climate Change Storylines. Proceedings of the National Academy of Sciences.
- Comer, P., J. Kagan, M. Heiner, & C. Tobalske. 2002. Sagebrush Vegetation in the Western United States. Map 1: 2,000,000 scale. Compact Disc with metadata for the USGS Forest and Rangeland Ecosystems Science Center, Boise, Idaho. The Nature Conservancy. Boulder CO, USA.
- Comer, P.J., J. Hak, and G. Mendiguran. 2009. Alternative Methods for Mapping Terrestrial Ecosystems in the Great Basin of the Western United States. Report to the U.S. Geological Survey, Biological Resources Division. 37 p.
- Comer, P.J. & J Hak. 2009. NatureServe Landscape Condition Model. Technical documentation for NatureServe Vista decision support software engineering. NatureServe, Boulder CO.
- Daly, C., W. P. Gibson, G.H. Taylor, G. L. Johnson, P. Pasteris. 2002. A knowledge-based approach to the statistical mapping of climate. *Climate Research*, 22: 99-113.
- DOI. 2008. Inventory of Onshore Federal Oil and Natural Gas Resources and Restrictions to Their Development. Prepared by the U.S. Departments of the Interior, Agriculture and Energy. [http://www.blm.gov/wo/st/en/prog/energy/oil\\_and\\_gas/EPCA\\_III.html](http://www.blm.gov/wo/st/en/prog/energy/oil_and_gas/EPCA_III.html)
- DOE & BLM. 2008. Programmatic Environmental Impact Statement, Designation of Energy Corridors on Federal Land in the 11 Western States (DOE/EIS-0386).<http://corridoreis.anl.gov/documents/fpeis/index.cfm>
- Enserink, M. 1999. Biological invaders sweep in. *Science*. 285(5435): 1834-1836.
- Erman, N.A. 2002. Lessons from a long-term study of springs and spring invertebrates (Sierra Nevada, California, USA) and implications for conservation and management. In: Sada D.W., Sharpe, S.E., editors; 2002; Las Vegas, NV.
- Gibson, W.P., C. Daly, T. Kittel, D. Nychka, C. Johns, N. Rosenbloom, A. McNab, and G. Taylor. 2002. Development of a 103-year high-resolution climate data set for the conterminous United States. In: Proc., 13th AMS Conf. on Applied Climatology, Amer. Meteorological Soc., Portland, OR, May 13-16, 181-183.
- Hall, R. O., M. F. Dybdahl, and M. C. VanderLoop. 2006. Extremely high secondary production of introduced snails in rivers. *Ecological Applications*. 16 (3): 1121-1131.

- Hershler, R. and D. W. Sada. 2002. Biogeography of Great Basin aquatic snails of the genus *Pyrgulopsis*. *Smithsonian Contributions to the Earth Sciences* 33:255-276.
- Leu, M., Hanser, S.E., Knick, S.T. 2008. The Human Footprint in the West: A Large-Scale Analysis of Anthropogenic Impacts. *Ecological Applications* 18: 1119-1139.
- Lowry, J. R.D. Ramsey, K. Thomas, D. Schrupp, T. Sajwaj, J. Kirby, E. Waller, S. Schrader, S. Falzarano, L. Langs, G. Manis, C. Wallace, K. Schulz, P. Comer, K. Pohs, W. Rieth, C. Velasquez, B. Wolk, W. Kepner, K. Boykin, L. O'Brian, D. Bradford, B. Thompson, and J. Prior-Magee. 2007. Mapping moderate-scale land-cover over very large geographic areas within a collaborative framework: a case study of the Southwest Regional Gap Analysis Project (SWReGAP). *Remote Sensing and Environment* 108: 59-73.
- NatureServe. 2009. Terrestrial Ecological Systems of the Conterminous United States. Version 2.7. Completed in cooperation with USGS Gap Analysis Program and inter-agency LANDFIRE. MMU approx. 2 hectares. NatureServe, Arlington, VA, USA. Digital map.
- NREL. 2010. Concentrating Solar Power Resource Maps. <http://www.nrel.gov/csp/maps.html>. National Renewable Energy Laboratory.
- NREL. 2008a. Photovoltaic Solar Resource Map of the United States. National Renewable Energy Laboratory. [http://www.nrel.gov/gis/data\\_analysis.html](http://www.nrel.gov/gis/data_analysis.html)
- NREL. 2008b. Biomass Resources in the United States. National Renewable Energy Laboratory. <http://www.nrel.gov/docs/fy06osti/39181.pdf>.
- NREL. 2005. A Geographic Perspective on the Current Biomass Resource Availability in the United States. National Renewable Energy Laboratory. <http://www.nrel.gov/docs/fy06osti/39181.pdf>
- NREL. 1986. Wind Energy Resource Atlas of the United States. National Renewable Energy Laboratory. <http://rredc.nrel.gov/wind/pubs/atlas/>
- Peterson, E. B. 2006. A map of invasive annual grasses in Nevada derived from multitemporal Landsat 5 TM imagery. Report for the U.S.D.I. Bureau of Land Management, Nevada State Office, Reno, by the Nevada Natural Heritage Program, Carson City, Nevada.
- Richards, D. C. personal communication. EcoAnalysts. Inc., Bozeman, MT
- Rosentrater, R. & M. Pellant. *In prep.*. Site potential for biological soil crusts development based on biological and physical factors. Internal draft BLM document.
- Sada, D.W, Williams J.E., Silvey J.C., Halford A., Ramakka J., Summers P., and L. Lewis. 2001. A guide to managing, restoring, and conserving springs in the Western United States. Denver: Bureau of Land Management. Report nr 1737-17. 70 p. \
- Sayre, R., P. Comer, H. Warner, and J. Cress. 2009. A new map of standardized terrestrial ecosystems of the conterminous United States: U.S. Geological Survey Professional Paper 1768, 17 p.
- Shepard, W.D. 1993. Desert springs-both rare and endangered. *Aquatic Conservation: Marine and Freshwater Ecosystems* 3(4):351-359.

- Spaulding, S.A., and L. Elwell. 2007. Increase in nuisance blooms and geographic expansion of the freshwater diatom *Didymosphenia geminata*: U.S. Geological Survey Open-File Report 2007-1425, 38 p.
- Theobald, D.M. 2005. Landscape patterns of exurban growth in the USA from 1980 to 2020. *Ecology and Society* 10(1): 32. [online] URL: <http://www.ecologyandsociety.org/vol10/iss1/art32/>.
- Theobald, D.M. 2008. *Network and accessibility methods to estimate the human use of ecosystems*. Proceedings of the International Conference on Geographic Information Science AGILE (Association of Geographic Information Labs Europe), Girona, Spain. 7 pgs.
- Thompson, B.C., Matusik-Rowan P.L., and K. G. Boykin. 2002. Prioritizing conservation potential of arid-land montane natural springs and associated riparian areas. *Journal of Arid Environments* 50(4):527-547.
- U. S. Geological Survey. NonIndigenous Aquatic Species website: <http://nas.er.usgs.gov/default.aspx>
- U.S. Environmental Protection Agency (EPA); Bierwagen, B., D.M. Theobald, C.R. Pyke, A. Choate, P. Groth, J.V. Thomas, and P. Morefield). 2009 Land-Use Scenarios: National-Scale Housing-Density Scenarios Consistent with Climate Change Storylines. Global Change Research Program, National Center for Environmental Assessment, Washington, DC; EPA/600/R-08/076F.
- USDA. 2007. Agriculture Census of the United States. US Department of Agriculture, National Agricultural Statistics Service. <http://www.agcensus.usda.gov/>
- Wisdom, M. J., G. Hayward, S. Shelly, C.D. Hargis, R.S. Holthausen, J. Epifiano, L. Parker, and J. Kershner. 2004. Using species groups and focal species for assessment of species at risk in forest planning. La Grande OR: U.S. Forest Service, Pacific Northwest Research Station.

# Appendices

---

**Appendix I: Master Data Table for the Central Basin and Range REA**

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class I Terrestrial Coarse Filter	California ReGAP Land Cover	in review	CA ReGAP Land Cover of terrestrial ecological systems and land cover, circa 2003, released in 2009	Lennartz, S., et al., 2008. Final Report on Land Cover Mapping Methods for California Map Zones 3, 4, 5, 6, 12, and 13.	accepted	USGS	Yes	CE models of current distribution within CA	Good for use in combination with other sources for CE distribution models
CE Class I Terrestrial Coarse Filter	NWReGAP Land Cover	review not needed	Existing land cover that would cover extreme southern NV portion of Central Basin and Range ecoregion		rejected	USGS	Yes	CE current distribution	has already been incorporated into composite map coverage to be used
CE Class I Terrestrial Coarse Filter	SWReGAP Land Cover	review finished	Land Cover map of NV, UT, AZ, UT, CO, and NM, based on NatureServe ecological systems classification; circa 2001.	Lowry, J. R.D. Ramsey, K. Thomas, D. Schrupp, T. Sajwaj, J. Kirby, E. Waller, S. Schrader, S. Falzarano, L. Langs, G. Manis, C. Wallace, K. Schulz, P. Comer, K. Pohs, W. Rieth, C. Velasquez, B. Wolk, W. Kepner, K. Boykin, L. O'Brian, D. Bradford, B. Thompson, and J. Prior-Magee. 2007. Mapping moderate-scale land-cover over very large geographic areas within a collaborative framework: a case study of the Southwest Regional Gap Analysis Project (SWReGAP). Remote Sensing and Environment 108: 59-73.	accepted	USGS	Yes	Current terrestrial coarse filter CE distribution.	already included as part of NatureServe 2009 map
CE Class I Terrestrial Coarse Filter	Terrestrial Ecosystems of the Conterminous United States	review finished	The U.S. Geological Survey (USGS) modeled the distribution of terrestrial ecosystems for the contiguous United States using a standardized, deductive approach to associate unique physical environments with ecological systems characterized in NatureServe's terrestrial ecological systems classification. this map depicts predicted biophysical settings that might support each ecological system type; regardless of current land use/land cover.	Sayre, R., P. Comer, H. Warner, and J. Cress. 2009. A new map of standardized terrestrial ecosystems of the conterminous United States: U.S. Geological Survey Professional Paper 1768, 17 p.	accepted	USGS	Yes	For use as biophysical setting representing terrestrial coarse filter CEs	This data layer best suited to applications at multi-ecoregion-national scaled analysis. More precise and accurate data sets exist, and/or may be readily combined to serve intended purposes for these REAs.
CE Class I Terrestrial Coarse Filter	The Human Footprint in the West	in review	Map of the human footprint for the western United States from an analysis of 14 landscape structure and anthropogenic features.	Leu, M., Hanser, S.E., Knick, S.T. 2008. The Human Footprint in the West: A Large-Scale Analysis of Anthropogenic Impacts. Ecological Applications 18: 1119-1139.	(empty)	USGS	Yes	Characterizing current condition of terrestrial CEs	Likely suitable. Will be investigated along other modeling options in Task 3.
CE Class I Terrestrial Coarse Filter	LANDFIRE Biophysical Settings	review finished	The Biophysical Settings (BpS) layer represents the vegetation that may have been dominant on the landscape prior to Euro-American settlement and is based on both the current biophysical environment and an approximation of the historical disturbance regime. <a href="http://www.landfire.gov/version_comparison.php">http://www.landfire.gov/version_comparison.php</a>	<a href="http://www.landfire.gov/NationalProductDescriptions20.php">http://www.landfire.gov/NationalProductDescriptions20.php</a>	accepted	USFS LANDFIRE	Yes	Assessment of long-term trends in extent for Coarse-filter CEs; assessment of fire regime departure	Moderate to high, with additional review and potential refinement. Will be brought together with NRCS Ecological Site Descriptions/soil-based maps.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class I Terrestrial Coarse Filter	LANDFIRE Existing Vegetation Type	review finished	The Existing Vegetation Type (EVT) layer represents the vegetation currently present; as defined by NatureServe terrestrial ecological systems classification (with some modifications).	<a href="http://www.landfire.gov/NationalProductDescriptions21.php">http://www.landfire.gov/NationalProductDescriptions21.php</a>	accepted	inter-agency LANDFIRE	Yes	use in combination with ReGAP-based NatureServe land cover sources to refine ecoregion map of CE distribution and condition	Suitable for this purpose; error in sample plot attribution introduced error into these EVT maps. These were very early LANDFIRE map zones.
CE Class I Terrestrial Coarse Filter	LANDFIRE Reference Vegetation Data	review finished	Georeferenced & labeled samples of vegetation gathered by Landfire to use as training data for their mapping & modeling efforts. Each sample is labeled with an ecological system. Includes species composition & cover, structural variables, some disturbance information, and calculated fuels data. Environmental data (elev, aspect, slope, soils, etc) are not included.	LANDFIRE Reference Database - Overview <a href="http://www.landfire.gov/NationalProductDescriptions27.php">http://www.landfire.gov/NationalProductDescriptions27.php</a>  General Technical Report RMRS-GTR-92: Integrating Ecosystem Sampling, Gradient Modeling, Remote Sensing, and Ecosystem Simulation to Create Spatially Explicit Landscape Inventories	accepted	LANDFIRE	Yes	Input for spatial models of current distributions scenarios	High; but label errors between ReGAP and LF labeling detected, and will be reassessed for project uses.
CE Class I Terrestrial Coarse Filter	NatureServe Terrestrial Ecosystems and Land Cover	review finished	Composite national map combining and reconciling ReGAP map products in the SE, SW, and NW with LANDFIRE EVT nationally. Review, editing, and documentation completed by NatureServe. Includes imbedded thematic links to US-NVC, NWI, NLCD, and other land cover classifications.	NatureServe. 2009. Terrestrial Ecological Systems of the Conterminous United States. Version 2.7. Completed in cooperation with USGS Gap Analysis Program and inter-agency LANDFIRE. MMU approx. 2 hectares. NatureServe, Arlington, VA, USA. Digital map.	accepted	USGS GAP, LANDFIRE, NatureServe	Yes	Current distribution of terrestrial CEs	Suitable for this use, with additional review and refinement
CE Class I Terrestrial Coarse Filter	NatureServe Landscape Condition	in review	NatureServe Level I (remotely sensed/modeled) measure of current condition/integrity for terrestrial CEs	Comer, P.J. & J Hak. 2009. NatureServe Landscape Condition Model. Technical documentation for NatureServe Vista decision support software engineering. NatureServe, Boulder CO.	accepted		Yes	For overlay with current CE distributions to gauge current ecological integrity, and as a 'resistance surface' for modeling landscape connectivity for CE distributions.	Highly suitable for certain CEs; can be updated and refined easily with local data.
CE Class I Terrestrial Coarse Filter	SageMap	in review	2002 integration of classification and available map data to depict sagebrush and related vegetation across the inter-mountain West.		rejected	USGS		For evaluation and refinement of composite map of current CE distributions.	suitable for use
CE Class I Terrestrial Coarse Filter	Ecological Integrity Criteria	in review	Descriptive text, metrics, and thresholds for gauging ecological integrity of examples for upland and wetland ecological systems. These reflect 2008 standards established by NatureServe		accepted	NatureServe		For use in conceptual and spatial modeling of integrity for coarse filter CEs	suitable for selected types
CE Class I Terrestrial Coarse Filter	NatureServe Element Occurrence Ranking Criteria for Ecological Systems	in review	Criteria to rank occurrences for ecological integrity, based on NatureServe 2000 data standards.		accepted			Input to conceptual models of ecological integrity for coarse-filter CEs	suitable for use

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class I Terrestrial Coarse Filter	Landfire Rapid Assessment models	in review	These are models created during the LANDFIRE rapid assessment stage. These models, by and large, have been superceded by the LANDFIRE national models. However, they are valuable for reference			Interagency: the USFS is the lead Agency		For input to refinement of existing conceptual models	suitable for this use
CE Class I Terrestrial Coarse Filter	Humboldt-Toyabe NF Existing Vegetation	need to review	existing vegetation maps by FS district (15 maps)	various		Forest Service		review and refinement of ecoregion-wide EVT map	will review during Task 3
CE Class I Terrestrial Coarse Filter	Great Basin Integrated Landscape Monitoring - NatureServe BpS model	review not needed	Using USGS national 'footprint' inputs (bioclimate, landform, surficial lithology) we completed a series of new maps through inductive modeling, using subsets of available sample data to simulate alternative mapping approaches given varying quantities of availability for georeferenced samples.	Comer, P.J., J. Hak, and G. Mendiguran. 2009. Alternative Methods for Mapping Terrestrial Ecosystems in the Great Basin of the Western United States. Report to the U.S. Geological Survey. 37 p.	accepted	NatureServe	Yes	for review and refinement of Landfire BpS maps	suitable for use
CE Class I Terrestrial Coarse Filter	SW-ReGap - Land Cover Field-Based Map Training Points	in review	This database represents the training point and quality control check for training site data collected in the landcover mapping effort for the Southwest Regional GAP Analysis Project. Field surveys were conducted between 2002 and 2004 throughout the region.	Lowry, J. H., Jr., R. D. Ramsey, K. Boykin, D. Bradford, P. Comer, S. Falzarano, W. Kepner, J. Kirby, L. Langs, J. Prior-Magee, G. Manis, L. O'Brien, T. Sajwaj, K. A. Thomas, W. Rieth, S. Schrader, D. Schrupp, K. Schulz, B. Thompson, C. Velasquez, C. Wallace, E. Waller and B. Wolk. 2005. Southwest Regional Gap Analysis Project: Final Report on Land Cover Mapping Methods, RS/GIS Laboratory, Utah State University, Logan, Utah.	accepted	USGS National Gap Analysis Program	Yes	Interpretation, training and validation of ecological systems mapping.	Sites offer the best source for precision identification of ecological system types and locations. Additional use in long-term monitoring efforts and community changes due to development and/or climate change.
CE Class I Terrestrial Coarse Filter	California ReGap - Training Points	in review	Fine scale on the ground documentation of ecological systems. Used to develop land cover maps for CA ReGap.	Lennartz, S., et al., 2008. Final Report on Land Cover Mapping Methods for California Map Zones 3, 4, 5, 6, 12, and 13.		USGS National Gap Analysis Program	No	Interpretation, training and validation of ecological systems mapping.	Sites offer the best source for precision identification of ecological system types and locations. Additional use in long-term monitoring efforts and community changes due to development and/or climate change.
CE Class II Terrestrial Fine Filter	Black-tailed Prairie dog Colonies, 1970 - 2002	need to review	This data represents a merging of all historic and current occupied and unoccupied Black-tailed Prairie Dog colony polygons acquired through March of 2003. Data quality ranges from hand drawn digitized maps to meter accurate GPS surveyed polygons. Data		(empty)	BLM	Yes		
CE Class II Terrestrial Fine Filter	Breeding Bird Survey (BBS)	review finished	This data consist of a series of data files that summarize population change and relative abundance for North American Birds from North American Breeding Bird Survey (BBS) data.		accepted	USGS	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class II Terrestrial Fine Filter	Sage Grouse Habitat of the West	need to review	Sage Grouse habitat. Was delivered as "Bruces National Project", but only a partial delivery was obtained from BLM. The primary raster file was not delivered, only the pyramid layer.		(empty)	BLM	No		
CE Class II Terrestrial Fine Filter	Christmas Bird Count	need to review	An annual hemispheric early-winter bird census.			USGS	Yes		
CE Class II Terrestrial Fine Filter	Core Sage Grouse	need to review			(empty)	BLM	Yes		
PL Class II Specially Designated Areas of Ecological Value	Desert Tortoise critical habitat	need to review	Designation of critical habitat was based on those areas recommended for recovery of the desert tortoise in the Draft Recovery Plan for the Desert Tortoise (Mojave Population) (U.S. Fish and Wildlife Service, 1993).		accepted	USFWS	Yes		
CE Class II Terrestrial Fine Filter	Desert Tortoise predicted habitat	need to review	Predicted habitat potential index values for desert tortoise ( <i>Gopherus agassizii</i> ) in the Mojave and parts of the Sonoran Deserts of Arizona, Nevada, Utah, and Arizona.		accepted	USGS	Yes		
CE Class II Terrestrial Fine Filter	Desert Tortoise suitable habitat	in review	Suitable habitat of Desert Tortoise ( <i>Gopherus agassizii</i> ) in Arizona. Digitized from 1:100,000 scale manuscripts prepared by Field Office Wildlife Specialists or digitized on-screen and edited at 1:100,000 scale or larger by GIS specialists. The criteria		accepted	BLM	Yes		
CE Class II Terrestrial Fine Filter	Mule Deer Locations	in review	Delphi (expert opinion) approach to map all mule and black-tailed deer habitat in North America and Mexico. Six categories of mule deer habitat were delineated, with 18 factors limiting or otherwise affecting the habitat. Classes include Year-around Population, Winter concentration, Winter range, Summer range, Limited range, and Other important habitat.		accepted	Utah State University Extension	No	representation of current seasonal habitats for Mule deer CE	Relatively coarse spatial resolution, but adequate.
CE Class II Terrestrial Fine Filter	Transport Atlas Bird and Mammal distributions	need to review	This data set contains distribution information for all birds and mammals occurring in the Western Hemisphere, as well as Native US fish by watershed.			NatureServe	Yes		
CE Class II Terrestrial Fine Filter	Sage Grouse lek locations	need to review	A westwide compilation of state sage grouse lek point datasets for year 2006		(empty)	BLM	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class II Terrestrial Fine Filter	SWReGAP Vertebrate Habitat Models	in review	This dataset contains ratings of the suitability of habitat for the predicted distributions of 817 native terrestrial vertebrate species in the 5-State SWReGAP project area. Of these models, 234 are for BLM species in Mojave and/or Central Basin ecoregions.	Boykin, K. G., B. C. Thompson, R. A. Deitner, D. Schrupp, D. Bradford, L. O'Brien, C. Drost, S. Propeck-Gray, W. Rieth, K. Thomas, W. Kepner, J. Lowry, C. Cross, B. Jones, T. Hamer, C. Mettenbrink, K.J. Oakes, J. Prior-Magee, K. Schulz, J. J. Wynne, C. King, J. Puttere, S. Schrader, and Z. Schwenke. 2007. Predicted animal habitat distributions and species richness. Chapter 3 in J.S. Prior-Magee, et al., eds. outhwest Regional Gap Analysis Final Report. U.S. Geological Survey, Gap Analysis Program, Moscow, ID.		USGS GAP	Yes	Representing distributions of known and potential habitat for CE terrestrial species	High to Medium
CE Class II Terrestrial Fine Filter	Threatened and Endangered Species	in review	See "critical habitat". Need to evaluate what NS already has. Have requested additional endangered species data for Mojave.		(empty)	FWS	Yes		
CE Class II Terrestrial Fine Filter	CAGAP Vertebrate Habitat Models	need to review	This dataset contains ratings of the suitability of habitat for the predicted distributions of 455 native terrestrial vertebrate species in California. Of these models, 159 are for BLM species in the Mojave and/or Central Basin ecoregions.	Davis, F. W., D. M. Stoms, A. D. Hollander, K. A. Thomas, P. A. Stine, D. Odion, M. I. Borchert, J. H. Thorne, M. V. Gray, R. E. Walker, K. Warner, and J. Graae. 1998. The California Gap Analysis Project--Final Report. University of California, Santa Barbara, CA. [ <a href="http://www.biogeog.ucsb.edu/projects/gap/gap_rep.html">http://www.biogeog.ucsb.edu/projects/gap/gap_rep.html</a> ]  Chapter 3. PREDICTED ANIMAL DISTRIBUTIONS AND SPECIES RICHNESS <a href="http://www.biogeog.ucsb.edu/projects/gap/report/gap_rep_ch3.html">http://www.biogeog.ucsb.edu/projects/gap/report/gap_rep_ch3.html</a>		USGS			
CE Class II Terrestrial Fine Filter	Desert Tortoise Habitat Model	in review	A quantitative habitat model for the desert tortoise using an extensive set of field-collected presence data. Sixteen environmental data layers were converted into a grid covering the study area and merged with the desert tortoise presence data that we gathered for input into the Maxent habitat-modeling algorithm. This model provides output of the statistical probability of habitat potential that can be used to map potential areas of desert tortoise habitat. This type of analysis, while robust in its predictions of habitat, does not account for anthropogenic changes that may have altered habitat with relatively high potential into areas with lower	Nussear, K.E., Esque, T.C., Inman, R.D., Gass, Leila, Thomas, K.A., Wallace, C.S.A., Blainey, J.B., Miller, D.M., and Webb, R.H., 2009, Modeling habitat of the desert tortoise ( <i>Gopherus agassizii</i> ) in the Mojave and parts of the Sonoran Deserts of California, Nevada, Utah, and Arizona: U.S. Geological Survey Open-File Report 2009-1102, 18 p.		USGS	Yes	Tortoise habitat distribution	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
			potential.						
CE Class II Terrestrial Fine Filter	NatureServe Terrestrial Element Occurrence Data for CA, NV & UT		NatureServe, in collaboration with its member Natural Heritage Programs and Conservation Data Centres, maintains a database of rare and imperiled species and plant communities across the United States and Canada. The Element Occurrence (EO) records that form the core of the NatureServe database include information on the location, status, characteristics, numbers, condition, and distribution of elements of biological diversity using established Natural Heritage Methodology developed by NatureServe and The Nature Conservancy (TNC). An Element Occurrence (EO) is an area of land and/or water in which a species or natural community is, or was, present. An EO should have practical conservation value for the Element as evidenced by potential continued (or historical) presence and/or regular recurrence at a given location.			NatureServe	Yes	Location of CEs, input to distribution models	suitable for use
CE Class III Physical Feature	Elevation Derivatives for National Applications (EDNA)	need to review	Email contact at this website with polygon file of extract area, and description of project. Derivatives: Filled DEM, Sinks, Shaded Relief, Slope, Flow direction, Flow Accumulation, Aspect, Contours, Compound Topo Index, Reach Catchment Seedpoints, Reach			USGS			
CE Class III Physical Feature	Geology	in review	Geologic map of the United States (exclusive of Alaska and Hawaii)			USGS			
CE Class III Physical Feature	Gravity anomaly data (Bouguer anomaly)	in review	The grid of gravity anomaly data for the conterminous United States and adjacent marine areas was constructed from National Information Mapping Agency (NIMA) gravity data files.			USGS	Yes		
CE Class III Physical Feature	Land Surface Forms	in review	This dataset was derived from the NED based on various neighborhood analysis using a 1-km2 analysis window.	Sayre, R., P. Comer, H. Warner, and J. Cress. 2009. A new map of standardized terrestrial ecosystems of the conterminous United States: U.S. Geological Survey Professional Paper 1768, 17 p.		USGS	Yes		
CE Class III Physical Feature	Magnetic anomaly maps and data for North America	review not needed	Digital data grids for the magnetic anomaly map of North America.			USGS	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class III Physical Feature	800 m PRISM Monthly Temperature	need to review				Oregon State	Yes		
CE Class III Physical Feature	National Elevation Dataset - 30 m	review not needed	The National Elevation Dataset (NED) is the primary elevation data product produced and distributed by the USGS.	Citation_Information: Originator: U.S. Geological Survey (USGS) Publication_Date: 2009 Title: National Elevation Dataset (NED) Edition: 2 Geospatial_Data_Presentation_Form: raster digital data Publication_Information: Publication_Place: Sioux Falls, SD Publisher: U.S. Geological Survey Online_Linkage: <a href="http://nationalmap.gov">http://nationalmap.gov</a> Online_Linkage: <a href="http://seamless.usgs.gov">http://seamless.usgs.gov</a>	accepted	USGS	Yes	Spatially adequate for most modeling purposes and represents the best complete data set for the region.	The intended uses of the data are utilized by the scientific and resource management communities for global change research, hydrologic modeling, resource monitoring, mapping and visualization applications.
CE Class III Physical Feature	National Elevation Dataset - 10 m	review not needed	The National Elevation Dataset (NED) is the primary elevation data product produced and distributed by the USGS.	Citation_Information: Originator: U.S. Geological Survey (USGS) Publication_Date: 2009 Title: National Elevation Dataset (NED) Edition: 2 Geospatial_Data_Presentation_Form: raster digital data Publication_Information: Publication_Place: Sioux Falls, SD Publisher: U.S. Geological Survey Online_Linkage: <a href="http://nationalmap.gov">http://nationalmap.gov</a> Online_Linkage: <a href="http://seamless.usgs.gov">http://seamless.usgs.gov</a>	accepted	USGS	Yes	Spatially adequate for the majority of modeling purposes. Limited, but useful, potential for systems/species with specialized gradient relationships.	The intended uses of the data are utilized by the scientific and resource management communities for global change research, hydrologic modeling, resource monitoring, mapping and visualization applications.
CE Class III Physical Feature	NWS CPC Datasets	review not needed	Soil moisture, evaporation, precipitation, runoff, temperature			NWS, CPC	Yes		
CE Class III Physical Feature	SSURGO	need to review	we're waiting to hear back from our USGS partners who did the Eastern US and last I talked to them were filling in the big holes in the SSURGO data in the west.- JH			NRCS	Yes		
CE Class III Physical Feature	STATSGO2: US General Soil Map					NRCS	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class III Physical Feature	Surficial Materials Lithology	need to review	This dataset was derived from the 28 lithology classes identified in the USGS map "Surficial Materials in the conterminous United States". These were generalized and reclassified into a set of 18 lithologies that typically control or influence the distrib	Sayre, R., P. Comer, H. Warner, and J. Cress. 2009. A new map of standardized terrestrial ecosystems of the conterminous United States: U.S. Geological Survey Professional Paper 1768, 17 p.		USGS	Yes		
CE Class III Physical Feature	Base Lithology	need to review				USGS	No		
CE Class III Physical Feature	Landform	in review	Topographic position of the landscape derived from the 30m NED.			USGS	No		
CE Class III Physical Feature	Ombrotypes	need to review				USGS			
CE Class III Physical Feature	thermotypes	need to review				USGS			
CE Class IV Aquatic/Wetland Coarse Filter	Aquifers	review not needed	This map layer contains the shallowest principal aquifers of the conterminous United States, Hawaii, Puerto Rico, and the U.S. Virgin Islands, portrayed as polygons.			USGS	Yes	Not intended for use.	
CE Class IV Aquatic/Wetland Coarse Filter	Computed Topographic Index	in review	Combo of flow accumulation and slope for defining wetness zones			EPA			
CE Class IV Aquatic/Wetland Coarse Filter	USGS drought-detection wells gwwst0x020	in review	This map layer shows the locations of wells maintained by the U.S. Geological Survey (USGS) that are used to monitor the effects of droughts and other climate variability on ground-water levels.			USGS	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	Hydrodrologic Units	in review				USGS	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class IV Aquatic/Wetland Coarse Filter	Irrigation canals	need to review	Can pull from the NHD when we get the complete data.		(empty)			Will represent canals and significant ditches. Will be derived from NHD.	NHD will provide the most suitable map available.
CE Class IV Aquatic/Wetland Coarse Filter	Nation Hydrography Dataset - 1:100,000 -- aka NHDPlus	need to review		See the NHDPlus User Guide, USEPA, USGS and Horizon Systems Corporation, January 26, 2010. User Guide, description of NHDPlus, and metadata are all available at <a href="http://www.horizon-systems.com/nhdplus/">http://www.horizon-systems.com/nhdplus/</a> .	accepted	USGS	Yes	The NHDPlus data system provides the foundation for several assessments of aquatic CE occurrence condition, based on additional attributes for the system generated by the USGS (listed separately). NHDPlus makes it possible to use these additional data layers to assess aspects of catchment hydrology, climate and deposition of air pollutants.	This is a crucial platform for several assessments.
CE Class IV Aquatic/Wetland Coarse Filter	Nation Hydrography Dataset - 1:24,000	review not needed				USGS	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	National Wetlands Inventory (NWI)	need to review	Hydrologic Units down to the 6th order		accepted	USFWS	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	NWIS	review not needed	NWIS supports the acquisition, processing, storage and dissemination of information about water quantity and quality collected at over 1.5 million sites around the U.S. As a long-term database and information delivery system, NWIS provides continual access to data collected over the last 100+ years, as well as real-time data on streamflow, etc.			USGS	No		
CE Class IV Aquatic/Wetland Coarse Filter	Topographic Moisture Potential	need to review	This dataset was derived to help contribute substrate moisture regimes and was based on the derivation of ground moisture potential using a combination of computed topographic characteristics and mapped National Wetland Inventory boundaries.	Sayre, R., P. Comer, H. Warner, and J. Cress. 2009. A new map of standardized terrestrial ecosystems of the conterminous United States: U.S. Geological Survey Professional Paper 1768, 17 p.		USGS	Yes		
CE Class IV Aquatic/Wetland	U.S. Army Corps of Engineers Navigable Waterway Network (Line)	review not needed	The National Waterway Network is a comprehensive network database of the nation's navigable waterways.			Bureau of Transportation	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
Coarse Filter						Statistics			
CE Class IV Aquatic/Wetland Coarse Filter	U.S. Army Corps of Engineers Navigable Waterway Network (Node)	review not needed	The National Waterway Network is a comprehensive network database of the nation's navigable waterways.			Bureau of Transportation Statistics	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	Watershed Boundary Database	need to review			(empty)	NRCS			
CE Class IV Aquatic/Wetland Coarse Filter	>2000 springs Biological/Environmental database: Desert Research Institute	in review	Dr. Don Sada, Desert Research Institute, NV is compiling a database on biotic and environmental conditions for almost 5000 springs in our ecoregion. He is willing to work with us pending future discussions.		(empty)	Desert Research Institute University Nevada Reno	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	Nevada DEP Stream Bioassessment Data	need to review	Nevada began its Bioassessment Program in the year 2000 and has continued to collect biological, chemical and physical habitat information on an annual basis throughout Nevada.			Nevada Division Environmental Protection Bureau of Water Quality Planning, Bioassessment Program	Unkn own	These data will meet two needs: (1) The assessment of current biotic condition in stream/river ecosystem CEs; and (2) the assessment of aquatic nuisance species distributions among CEs and their associated HUCs.	If the data meet standards set by EPA Western Streams Assessment for sampling design, field methods, and analysis, they can be included in the baseline assessment.
CE Class IV Aquatic/Wetland Coarse Filter	USEPA National Lakes Assessment	need to review	EPA and its state and tribal partners have conducted a survey of the nation's lakes, ponds and reservoirs. This National Lakes Assessment is designed to provide statistically valid regional and national estimates of the condition of lakes. It uses a probability-based sampling design to represent the condition of all lakes in similar regions sharing similar ecological characteristics. Consistent sampling and analytical procedures ensure			USEPA	Unkn own		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
			that the results can be compared across the country.						
CE Class IV Aquatic/Wetland Coarse Filter	Lakes, Playas, and Other Water Bodies of Nevada	need to review	NV Heritage ecologist attributed layer of lakes, playas, rivers Categories include mud playa, salt playa,. Also designated Major (large) and Minor (small)			Nevada Natural Heritage Program			
CE Class IV Aquatic/Wetland Coarse Filter	NV Spring Terrestrial Vegetation Dataset	need to review	171 vegetation plot taken at Springs throughout NV by the heritage program. Included Plant Association Name and EO Rank information.						
CE Class IV Aquatic/Wetland Coarse Filter	National Atmospheric Deposition Program (NADP) Atmospheric Deposition	in review	The National Atmospheric Deposition Program (NADP) monitors precipitation chemistry. The program is a cooperative effort between many different group, including federal, state, tribal and local governmental agencies, educational institutions, private companies, and non-governmental agencies. See URL <a href="http://nadp.sws.uiuc.edu/Default.aspx">http://nadp.sws.uiuc.edu/Default.aspx</a> for details			USGS	Yes	These data, along with regional estimation model output from the NADP website, support the assessment of the threat(s) posed by atmospheric deposition as a CA.	These are the best data to use for the assessment of atmospheric deposition rates and their spatial variation.
CE Class IV Aquatic/Wetland Coarse Filter	California Groundwater Basins from Calif. DWR	review finished	Map of California Groundwater Basins and identification numbers linked to CDWR Bulletin 118 (2003) for technical info on each basin. See <a href="http://www.water.ca.gov/groundwater/bulletin118/gwbasin_maps_descriptions.cfm">http://www.water.ca.gov/groundwater/bulletin118/gwbasin_maps_descriptions.cfm</a>			California Department of Water Resources	Yes	These data will be used to delineate groundwater basins, in conjunction with the extraction of information from CDWR Bulletin 118 (2003) to identify aquifers that significantly affect the hydrology of spring/seep and stream/river CEs. The CDWR data provide a state-specific backup to the data in the USGS Southwest Principal Aquifers study.	These data are a highly suitable backup to using data from the USGS Southwest Principal Aquifers study. We'll use whichever is more precise.
CE Class IV Aquatic/Wetland Coarse Filter	Arizona Groundwater Basins	review finished	Arizona groundwater basins, <a href="http://www.azwater.gov/azdwr/GIS/">http://www.azwater.gov/azdwr/GIS/</a> : "The data provide base information for use in GIS systems to aid in assessment for a variety of planning and analysis purposes and to provide a geographic view with corresponding data. 'Groundwater basin' means an area which, as nearly as known facts permit as determined by the director pursuant to this chapter, may be designated so as to enclose a relatively hydrologically distinct body or related bodies of groundwater, which shall be described horizontally by surface description."		rejected	Arizona Department of Water Resources	Yes	These data are to be used as a backup to data from the USGS Southwest Principal Aquifers study, to delineate groundwater basins that have significant bearing on aquatic CEs, specifically, springs and seeps, and streams and rivers.	These data are highly suitable at the state level, as a substitute for aquifers delineated by the USGS Southwest Principal Aquifers study. We'll use whichever is more precise.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class IV Aquatic/Wetland Coarse Filter	Arizona Groundwater Sub-Basins	review finished	Arizona groundwater subbasins, <a href="http://www.azwater.gov/azdwr/GIS/">http://www.azwater.gov/azdwr/GIS/</a> : "The data provide base information for use in GIS systems to aid in assessment for a variety of planning and analysis purposes and to provide a geographic view with corresponding data. 'Subbasin' means an area which, as nearly as known facts permit as determined by the director pursuant to this chapter, may be designated so as to enclose a relatively hydrologically distinct body of groundwater within a groundwater basin, which shall be described horizontally by surface description."			Arizona Department of Water Resources	Yes	This provides a more detailed view of groundwater distribution, by sub-basin, in Arizona. It is to be used as an Arizona-specific "backup" to the data from the USGS Southwest Principal Aquifers study. The sub-basin polygons are nested within the groundwater Basin polygons, represented in a separate dataset.	This is a state-specific dataset, highly suitable for use as a backup to using the USGS Southwest Principal Aquifers data, whichever is the more precise.
CE Class IV Aquatic/Wetland Coarse Filter	Arizona Groundwater Site Inventory	review finished	Arizona Groundwater Site Inventory (GWSI) database, <a href="http://www.azwater.gov/azdwr/GIS/">http://www.azwater.gov/azdwr/GIS/</a> is ADWR's main repository for state-wide groundwater data. The GWSI consists of field-verified data regarding wells and springs collected by personnel from Hydrology Division's Basic Data Section, the U.S. Geological Survey, and other co-operating agencies.			Arizona Department of Water Resources	Yes	Unless we have comparable data from the other states in either CBR or MBR, this becomes a localized dataset. It's purpose is to help assess the intensity of groundwater use, as a backup to using the data in the Southwest Principal Aquifers study.	This provides backup data for purposes of assessing the threats posed by groundwater extraction to stream and spring CEs.
CE Class IV Aquatic/Wetland Coarse Filter	Stream baseflow index grid for the conterminous US-USGS	review finished	This 1-kilometer raster (grid) dataset for the conterminous United States was created by interpolating base-flow index (BFI) values estimated at U.S. Geological Survey (USGS) streamgages. Base flow is the component of streamflow that can be attributed to ground-water discharge into streams. For all documentation and citations, see <a href="http://water.usgs.gov/GIS/metadata/usgswrd/XML/bfi48grd.xml">http://water.usgs.gov/GIS/metadata/usgswrd/XML/bfi48grd.xml</a> and <a href="http://ks.water.usgs.gov/pubs/abstracts/of.03-263.htm">http://ks.water.usgs.gov/pubs/abstracts/of.03-263.htm</a> (the latter site also provides contact information for the dataset author).	Wolock, D.M., 2003b, Estimated mean annual natural ground-water recharge estimates in the conterminous United States: U.S. Geological Survey Open-File Report 03-311, digital dataset, available on the World Wide Web, accessed August 20, 2003, at URL <a href="http://water.usgs.gov/lookup/getspatial?rech48grd">http://water.usgs.gov/lookup/getspatial?rech48grd</a>		USGS	Yes	Allows assessment of stream baseflow by aggregating the gridded data by HUC. This in turn is a crucial component of stream hydrology for arid lands streams, which we can therefore assess for current conditions	Very high; the USGS "BFI" method is well established and well documented.
CE Class IV Aquatic/Wetland Coarse Filter	Nitrate contamination, probability for recently recharged ground waters in the Conterminous US	in review	This data set is a national map of predicted probability of nitrate contamination of shallow ground waters based on a logistic regression (LR) model. The LR model was used to predict the probability of nitrate contamination exceeding 4 mg/L in predominantly shallow, recently recharged ground waters of the US. For all documentation and citations, see <a href="http://water.usgs.gov/GIS/metadata/usgswrd/XML/gwrisk.xml">http://water.usgs.gov/GIS/metadata/usgswrd/XML/gwrisk.xml</a>	Nolan, B.T., Hitt, K.J., and Ruddy, B.C., 2002, Probability of nitrate contamination of recently recharged ground waters in the conterminous United States. Environmental Science and Technology Volume 36, Number 10, Pages 2138-2145.		USGS		Provides a means to assess potential for altered nutrient regime in streams, springs and wetlands, in absence of field data on nutrient levels.	Strongly suitable, but may not be very informative for many areas of CBR and MBR where nearby sources of nitrate are sparse.
CE Class IV Aquatic/Wetland Coarse Filter	Hydrographic data for Great Basin groundwater systems, 1:1,000,000	review not needed	This three-part data set consists of 1:1,000,000-scale (a) areas where shallow ground water is consumed by evapotranspiration (ET); (b) hydrographic area and major flow system boundaries and polygons; and (c) large springs for the Great Basin. The source is Harrill, J.R., Gates, J.S., and Thomas, J.M., 1988, Major ground-water flow systems in the Great Basin region of	Harrill, J.R., Gates, J.S., and Thomas, J.M., 1988, Major ground-water flow systems in the Great Basin region of Nevada, Utah, and adjacent states: U.S. Geological Survey Hydrologic Investigations Atlas HA-694-C, scale 1:1,000,000		USGS	Yes	To identify which aquifers/watersheds contribute the water that supports the crucial baseflow in streams and water levels/discharge rates in springs.	This is an older dataset that will be compared to the newer SWPA dataset from USGS, to identify which is best for providing basic information on surface-groundwater interactions in the CBR.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
			Nevada, Utah, and adjacent states: U.S. Geological Survey Hydrologic Investigations Atlas HA-694-C, scale 1:1,000,000						
CE Class IV Aquatic/Wetland Coarse Filter	Hydrologic Attributes for NHDPlus Catchments (Version 1.1) for the Conterminous United States	review finished	These are datasets developed by the USGS as attributes for NHDPlus Catchments. They provide data on catchment atmospheric deposition (2 datasets) and catchment hydrology (6 datasets).	BFI: Wolock, D.M., 2003, Base-flow index grid for the conterminous United States: U.S. Geological Survey Open-File Report 03-263, digital data set, available at <a href="http://water.usgs.gov/lookup/getspatial?bfi48grd">http://water.usgs.gov/lookup/getspatial?bfi48grd</a> .		USGS	Yes	The two atmospheric deposition datasets provide a single-year snapshot of nitrate deposition, as a backup to using estimates directly from the NADP database, to assess this CA. The data will be aggregated by HUC for CA analysis. The six hydrologic datasets provide information on related to runoff and recharge behavior by catchment. They also will be aggregated by HUC, but for CE condition analysis. Additionally, they provide a "Plan B" for assessing the impacts of climate change on runoff and recharge, as a backup to our using the Flint et al. USGS modeled (forecast) data.	The two atmospheric deposition datasets provide a highly suitable backup to working directly with NADP data and regionalized deposition estimates for this CA analysis. The baseflow, runoff and recharge datasets provide a key means for characterizing the hydrology of stream ecosystem CEs. And the six hydrologic datasets together provide a highly suitable backup to working directly with the Flint et al. USGS forecast data on the impacts of climate change, especially if the GCM, emissions scenario, or timestep choices built into the Flint et al. data are not compatible with those used in the rest of the CBR and MBR REAs.
CE Class IV Aquatic/Wetland Coarse Filter	USGS Southwest Principal Aquifers (SWPA) study data	in review	These are five datasets from the USGS Southwest Principal Aquifers (SWPA) study, published in 2008 as Geospatial Data to Support Analysis of Water-Quality Conditions in Basin-Fill Aquifers in the Southwestern United States, U.S. Geological Survey Scientific Investigations Report (SIR) 2008-5239.	McKinney, T.S., and Anning, D.W., 2009, Geospatial data to support analysis of water-quality conditions in basin-fill aquifers in the southwestern United States: U.S. Geological Survey Scientific Investigations Report 2008-5239, 16 p. Available at <a href="http://pubs.er.usgs.gov/sir/2008/5239">http://pubs.er.usgs.gov/sir/2008/5239</a> .		USGS	Yes	These data identify and delineate the aquifers on which spring/seep and stream/river CEs depend for maintaining water levels or base flows. Additionally, the data provide crucial information on agricultural and municipal water use from these aquifers -- information crucial to assessing the potential impacts of future water resource development associated with land development or other forms of development (as a CA).	These will probably be our primary datasets for assessing which aquifers support which aquatic ecosystem CEs; and our primary means for assessing the potential impacts of water resource or land development (as CAs) on these CEs.
CE Class IV Aquatic/Wetland Coarse Filter	USGS-Nevada State joint study of Nevada alluvial aquifers	in review	Three data sets were created as part of a U.S. Geological Survey study, done in cooperation with the Nevada Division of Environmental Protection, to evaluate the susceptibility and vulnerability of ground water to anthropogenic contamination.	Lopes, T.J., Buto, S.G., Smith, J.L., and Welborn, T.L., 2006, Water-table levels and gradients, Nevada, 1947-2004: U.S. Geological Survey Scientific Investigations Report 2006-5100		USGS		This is a backup dataset for the delineation of aquifers critical to supporting spring/seep and stream/river CEs. It also provides information on changes in water storage in these aquifers, which supports	This is a backup to using the data generated by the USGS Southwest Principal Aquifers study. It is a state database; we'll use whichever is more precise.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
								the assessment of water use as a potential stressor/CA.	
CE Class IV Aquatic/Wetland Coarse Filter	Utah State University-Western Center for Monitoring & Assessment of Freshwater Ecosystems database system	in review	This is the site of a query tool to download data generated by: the USEPA EMAP Western Streams Assessment project; two USEPA STAR grant projects to Utah State University in support of the Western Streams Assessment; and the BLM. The data are managed by the Western Monitoring Center at Utah State University (see link in Dataset Filename entry).			USEPA and BLM via Utah State University		These data provide primary information on the biological condition of stream/river ecosystem CEs.	Because the EMAP and STAR data were developed through a statistically robust geographic sampling design, they provide the statistically most reliable basis for assessing stream/river CE condition without concern for spatial sampling biases at the regional scale.
CE Class IV Aquatic/Wetland Coarse Filter	Stream baseflow index-Western US-Hill & Olson	in review	Calculation of the percentage of flow attributed to groundwater. Index was calculated for each of 9,941 USGS gaging stations in the western USA and values for unmeasured locations were interpolated using inverse-distance-squared weighting of the 12 closest gaging stations within 100 kilometers. Each interpolated value represents a 4 x 4 kilometer cell.			Utah State University, via authors (Hill & Olson)		This would be used as a backup database to assess the contribution of groundwater discharge to the hydrologic regime of stream/river ecosystem CEs; we would use the Hill & Olson findings if we encounter problems with the USGS (Wolcock) model of baseflow.	Highly suitable for assessing the baseflow component of stream/river hydrologic regimes.
CE Class IV Aquatic/Wetland Coarse Filter	Utah Department of Environmental Quality, comprehensive assessment of stream ecosystems (UCASE)	in review	A database generated by the state of Utah's comprehensive stream biomonitoring program, containing data on stream biotic and habitat condition. The 2008-9 strategic plan and the data we potentially need can be accessed through the following links: <a href="http://www.waterquality.utah.gov/Monitoring/index.htm">http://www.waterquality.utah.gov/Monitoring/index.htm</a> The same location lists a contact for questions about the data: Jim Harris at 801-536-4360 or e-mail <a href="mailto:jamesharris@utah.gov">jamesharris@utah.gov</a>			Utah Department of Environmental Quality		These data provide information on biotic and habitat conditions in stream/river ecosystem CEs sampled at the state scale, to supplement and complement the Western Streams Assessment information (see listing under Utah State University). The database may also include information on aquatic nuisance species.	High, as complement to the regional (EMAP) database.
CE Class IV Aquatic/Wetland Coarse Filter	USEPA National Database of State Water Quality Status Listings	in review	This is the Reach Address Database (RAD) Download website. It is the source for the most recent EPA-approved state listings of: 303(d) Listed Impaired Waters; 305(b) Assessed Waters; Clean Watersheds Needs Survey; Fish Consumption Advisories; Nonpoint Source Projects; STORET Water Monitoring Locations; Facilities that Discharge to Water; Impaired Waters with TMDLs; and State Water Quality Standards. We would need to download each, for UT, NV, CA, and AZ; it may be possible select by HUC rather than by State.			USEPA, compiled from EPA-approved state water quality assessments	Yes	These data will allow us to assess current condition of all freshwater ecosystem and community CEs in terms of whether waters meet state water quality standards and what actions the states have identified as necessary to address both point and nonpoint source pollution and other stressors.	Short of our doing our own assessment of masses of water quality monitoring and watershed data, this is the best way for us to conduct a rapid assessment of whether individual water-body CEs are recognized by each state as "impaired" and, if so, the likely causes of that impairment.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class IV Aquatic/Wetland Coarse Filter	Groundwater_Climate_Response_Network	in review	Supplied data is incomplete. Shape file provided has 5 wells for NV, while on the web site they list data from 854 wells.				Yes		
CE Class IV Aquatic/Wetland Coarse Filter	BLM-Utah State University National Aquatic Monitoring Center Data	in review	The BLM National Aquatic Monitoring Center at Utah State University, aka the "Buglab" ( <a href="http://www.usu.edu/buglab/">http://www.usu.edu/buglab/</a> ) has built a large database of stream bioassessment data, mostly on benthic macroinvertebrates from stream sites on public lands throughout the western US. Some habitat data are also included. The data were not collected under a single spatial sampling design, but aside from their spatial unevenness they are among the best available and complement those maintained by the USU Western Monitoring Center (see separate entry).			BLM		The Buglab data will supplement the data from the Western Monitoring Center, for assessing the biotic condition of aquatic (stream/river) ecosystem CEs, and will help map the distribution of aquatic nuisance species (a CA). Dr. Miller has also developed regional benthic macroinvertebrate IBI metrics, and his regional classification may help identify distinct stream ecosystem types within the two ecoregions.	Very high. Potentially spatially uneven, so best if used in conjunction with the WMC data (see separate entry).
CE Class IV Aquatic/Wetland Coarse Filter	USEPA National Wadeable Streams Assessment	in review	The Wadeable Streams Assessment (WSA) is a first-ever statistically-valid survey of the biological condition of small streams throughout the U.S. The U.S. Environmental Protection Agency (EPA) worked with the states to conduct the assessment in 2004-2005.			USEPA		These data will supplement those obtained from the Western Monitoring Center, BLM "Buglab," and state bioassessment programs, for the assessment of (a) stream ecosystem CE condition and (b) aquatic nuisance species distributions (relates to a CA).	Limited sample size but wide breadth of data and sophisticated field and laboratory methods make this a dataset of limited use but highly suitable for that use.
CE Class IV Aquatic/Wetland Coarse Filter	Nevada 2006 303(d)/305(b) Impaired Waters List	in review	This presents the entire state of Nevada database on its "Impaired Waters" as required under the federal Clean Water Act. The data provide information on the status (degree of impairment) of all waters of the state, tagged by NHD designation. This is a backup dataset to the USEPA national integration of all states' Impaired Waters data for the last full reporting cycle (2006). We will use whichever is the more current.			Nevada DEP		This is a backup to using the USEPA national database on 303d/305b impaired waters, TMDLs, etc. If the USEPA database is current (and it should be) then we won't also need to use the Nevada state data layers. Either way, the data provide a means for assessing overall ecological integrity of aquatic ecosystem CEs based on state assessment of whether they meet "aquatic life use" standards. The "causes of impairment" listed for some waters may also include invasive species (aquatic nuisance species), so	Highly suitable, either as obtained from the state or from the USEPA.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
								the data will help with that CA assessment as well.	
CE Class IV Aquatic/Wetland Coarse Filter	California Surface Water Ambient Monitoring Program (SWAMP) Bioassessment Data	need to review	Stream/river bioassessment data for the state of California, collected according to rigorous state data collection and analysis standards.	E.g., Ode, P.R. and A.C. Rehn. 2005 Probabilistic assessment of the biotic condition of perennial streams and rivers in California. Report to the State Water Resources Control Board. California Department of Fish and Game Aquatic Bioassessment Laboratory, Rancho Cordova, California. Ode, P.R. 2007 Ecological condition assessment of California's perennial wadeable streams. Report to the State Water Resources Control Board's Non-Point Source Program. California Department of Fish and Game Aquatic Bioassessment Laboratory, Rancho Cordova, California.		California EPA, State Water Resources Control Board		As with the other state bioassessment datasets, these will be used to supplement the data from the regional stream bioassessment monitoring programs, the data for which will come from the two datasets at Utah State University, either from the BLM "Buglab" or the Western Monitoring Center. The data provide information on the biotic condition of stream ecosystem CEs; and on the distribution of aquatic nuisance species for that CA assessment.	Very high, although limited to wadeable, perennial streams -- which in both the CBR and MBR have very limited spatial distributions.
CE Class IV Aquatic/Wetland Coarse Filter	Arizona DEQ Bioassessment Program Data	need to review	Freshwater bioassessment data collected by ADEQ in support of the state water quality monitoring program. Data will include information on the biotic condition of probably both streams and lakes aquatic ecosystem CEs; and on aquatic nuisance species for CA assessment.			Arizona DEQ		As with the other state bioassessment databases, this database will provide information on the biotic condition of aquatic ecosystem CEs; and will contain information on aquatic nuisance species for CA assessment.	Highly suitable as supplement to the regional data, but limited to perennial waters.
CE Class IV Aquatic/Wetland Coarse Filter	U.S. Army Corps of Engineers National Inventory of Dams (NID)	review not needed	The NID contains 60 fields of data (identification, location, characteristics) for all dams that meet at least one of four criteria: 1) High hazard classification - loss of one human life is likely if the dam fails; 2) Significant hazard classification - possible loss of human life and likely significant property or environmental destruction; 3) Equal or exceed 25 feet in height and exceed 15 acre-feet in storage; or 4) Equal or exceed 50 acre-feet storage and exceed 6 feet in height. Data gaps are possible due to lags or inaccuracies in what states/tribes/territories report to NID.	Documentation on the NID is available at the website noted above, e.g., origins, update procedures, data fields, etc.		US Army Corps of Engineers		To help characterize aquatic coarse-filter CE condition w/r/t connectivity; and to help identify artificial water bodies to address surface water MQs	Very high
CE Class IV Aquatic/Wetland Coarse Filter	USGS Arid Western US runoff and recharge potential	need to review	These are data developed by Flint and Flint (2007 -- see Citation) to estimate watershed runoff and recharge potential using a 270m grid across most of the arid and semiarid western US as part of a study to investigate the interactions of climate and other controlling factors for runoff and recharge.	Flint, Lorraine E. and Alan L. Flint, 2007, Regional analysis of groundwater recharge. Chapter B, pages 29-60, in Stonestrom, D.A., Constantz, J., Ferré, T.P.A., and Leake, S.A., eds., Ground-water recharge in the arid and		USGS		The data will provide crucial information on (a) the surface-runoff driven component of stream hydrologic regimes for coarse-filter aquatic CEs; and (b) the likely recharge zones	Superb.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
				semi-arid southwestern United States: U.S. Geological Survey Professional Paper 1703.				(and rates of recharge) for basin fill aquifers, the eventual discharges from which support baseflow in these same CEs. The data are also necessary because the authors have also modeled the same hydrologic variables on the same grid based on downscaled climate projections. So we will be able to compare their "current conditions" model (described here) with the forecast conditions model (described in a separate entry) to assess potential impacts of climate change on stream hydrology and groundwater recharge.	
CA Class IV Climate Change	USGS Arid Western US future runoff and recharge potential under climate change	need to review	This dataset rests on the work by Flint and Flint (2007) described in the entry for "USGS Arid Western US runoff and recharge potential," but provides estimates of future hydrologic conditions based on climate change modeling. In May 2010, in an email message to Marni Koopman, the authors described the dataset as follows: "We are currently in the process of publishing finely downscaled climate change scenarios for the desert southwest and California. These are at 270-m spatial resolution for GFDL and PCM A2 and B1, with A1Fi to follow this summer.	The publication for the new data is not yet available; see Citation for the separate entry, "USGS Arid Western US runoff and recharge potential," for the original model description (Flint and Flint 2007)		USGS		These data are the primary tool we will use to assess the potential impacts of climate change on stream hydrology for coarse-filter aquatic CEs. As described above and in the entry for "USGS Arid Western US runoff and recharge potential," we will compare the model output for current versus projected runoff and recharge, aggregated to the scale of the watershed for each coarse-filter aquatic CE occurrence. This will allow a comparison of the ways in which mean annual total discharge, mean annual baseflow, and mean monthly discharge potentially will change under different climate change scenarios. We need to find out what time-steps the authors used, to know how we may be able to line up this assessment with that for terrestrial change.	Superb

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CE Class V Aquatic/Wetland Fine Filter	NatureServe Aquatic Element Occurrence Data for CA, NV & UT		NatureServe, in collaboration with its member Natural Heritage Programs and Conservation Data Centres, maintains a database of rare and imperiled species and plant communities across the United States and Canada. The Element Occurrence (EO) records that form the core of the NatureServe database include information on the location, status, characteristics, numbers, condition, and distribution of elements of biological diversity using established Natural Heritage Methodology developed by NatureServe and The Nature Conservancy (TNC). An Element Occurrence (EO) is an area of land and/or water in which a species or natural community is, or was, present. An EO should have practical conservation value for the Element as evidenced by potential continued (or historical) presence and/or regular recurrence at a given location.			NatureServe	Yes		
CE Class III Physical Feature (e.g., erodible soils)	GEOSS USA Moisture Class	in review	Assignment of flow accumulation models to specific moisture categories.  Class 1: Wetlands - $CTI \geq 18.5$ Class 2: Mesic Uplands - $12 \leq CTI < 18.5$ Class 3: Dry Uplands - $CTI < 12$ & not satisfy the aspect and slopes thresholds that identify very dry uplands (below) Class 4: Very Dry Uplands - $CTI < 12$ & $91 \leq \text{aspect} \leq 314$ & $\text{slopes} < 24 \text{degrees}$ (44.5%)			USGS	Yes		
CE Class IV Aquatic/Wetland Coarse Filter	Western Riparian Threats Assessment	need to review	Coarse-scale quantitative assessment of threats to riparian ecosystems using available spatial data applicable across western conterminous U.S.	Theobald, D.M., D.M., Merritt, and J.B. Norman, III. 2010. Assessment of threats to riparian ecosystems in the western U.S.		Forest Service		Calibration of aquatic CE condition assessment; considerations for reporting options.	TBD
CE Class V Aquatic/Wetland Fine Filter	Critical Habitat	need to review	These datasets identify the areas (in general) where final critical habitat for a variety of threatened and endangered plant and animal species occurs		accepted	USFWS	Yes		
CE Class V Aquatic/Wetland Fine Filter	EcoAnalysts Inc macroinvertebrate databases	need to review	EcoAnalysts Inc., Moscow ID has conducted taxonomic identification of aquatic macroinvertebrates, including natives and invasives, for hundreds of projects and hundreds of clients in the Western USA.			multiple agencies, compiled by EcoAnalysts, Moscow, ID	Unknown		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class I Wildfire	MTBS Burn Severity	need to review	The Monitoring Trends in Burn Severity (MTBS) project assesses the frequency, extent, and magnitude (size and severity) of all large wildland fires (includes wildfire, wildland fire use, and prescribed fire) in the conterminous United States (CONUS).			MTBS	No		
CA Class I Wildfire	MTBS Fire Occurrence	need to review	The Monitoring Trends in Burn Severity (MTBS) project assesses the frequency, extent, and magnitude (size and severity) of all large wildland fires (includes wildfire, wildland fire use, and prescribed fire) in the conterminous United States (CONUS), Alas			MTBS	Yes		
CA Class I Wildfire	MTBS Fire Perimeters	need to review	The Monitoring Trends in Burn Severity (MTBS) project assesses the frequency, extent, and magnitude (size and severity) of all large wildland fires (includes wildfire, wildland fire use, and prescribed fire) in the conterminous United States (CONUS), Alas			MTBS	Yes		
CA Class I Wildfire	GeoMAC - Geospatial Multi-Agency Coordination		This is a data set to represent the existing condition of a fire incident at the time data edit.		accepted	USGS	Yes		
CA Class I Wildfire	LANDFIRE Fire Behavior Models		13 Anderson (1982) Fire Behavior Fuel models; 40 Scott and Burgan (2005) Fire Behavior Models <a href="http://www.landfire.gov/products_national.php">http://www.landfire.gov/products_national.php</a>		accepted	LANDFIRE	Yes		
CA Class I Wildfire	NLDN (National Lightning Detection Network)	need to review	The National Lightning Detection Network, NLDN, consists of over 100 remote, ground-based sensing stations located across the United States that instantaneously detect the electromagnetic signals given off when lightning strikes the earth's surface.			NLDN, BLM	No		
CA Class I Wildfire	LANDFIRE FRCC Departure Index	in review	The Fire Regime Condition Class (FRCC) Departure Index data product uses a range from 0 to 100 to depict the amount that current vegetation has departed from simulated historical vegetation reference conditions.			LANDFIRE		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose. The data are, by and large, not suitable in isolation.
CA Class I Wildfire	LANDFIRE Mean Fire Return Interval	in review	The Mean Fire Return Interval layer quantifies the average period between fires under the presumed historical fire regime. This frequency is derived from vegetation and disturbance dynamics simulations using LANDSUM (Keane and others 2002, Hann and others 2004).			LANDFIRE			
CA Class I Wildfire	LANDFIRE Percent of Low-severity Fire	in review	The Percent of Low-severity Fire layer quantifies the amount of low-severity fires relative to mixed- and replacement-severity fires under the presumed historical fire regime.			LANDFIRE		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose. The data are, by and large, not suitable in isolation.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class I Wildfire	LANDFIRE Percent of Mixed-severity Fire	in review	The Percent of Mixed-severity Fire layer quantifies the amount of mixed-severity fires relative to low- and replacement-severity fires under the presumed historical fire regime.			LANDFIRE		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose. The data are, by and large, not suitable in isolation.
CA Class I Wildfire	LANDFIRE Percent of Replacement-severity Fire	review finished	The Percent of Replacement-severity Fire layer quantifies the amount of replacement-severity fires relative to low- and mixed-severity fires under the presumed historical fire regime.		accepted	LANDFIRE		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose. The data are, by and large, not suitable in isolation.
CA Class I Wildfire	LANDFIRE Environmental Site Potential (ESP)	review not needed	The LANDFIRE Environmental Site Potential (ESP) layer represents the vegetation that could be supported at a given site based on the biophysical environment, regardless of natural disturbance regime.			LANDFIRE		Give suitability of BpS maps for related purpose, this map will not be needed.	
CA Class I Wildfire	LANDFIRE Existing Vegetation Height (EVH)	need to review	Vegetation height represents the average height of the dominant vegetation for a 30-m grid cell.			LANDFIRE			
CA Class I Wildfire	LANDFIRE Existing Vegetation Cover (EVC)	review finished	Vegetation cover represents the average percent cover of existing vegetation for a 30-m grid cell.		rejected	LANDFIRE		These data provide seamless coverage of vegetation coverage by class.	NatureServe has more recent and more relevant data on vegetation coverage. Those will be used for spatial modelling / assessments.
CA Class I Wildfire	LANDFIRE Fire Regime Condition Class (FRCC)	review not needed	Fire regime condition class (FRCC) is a discrete metric that quantifies the amount that current vegetation has departed from the simulated historical vegetation reference conditions			LANDFIRE			
CA Class I Wildfire	USGS Land Treatment Digital Library	in review	The LTDL is a centralized digital library hosted by the USGS for federal managers and scientists. The LTDL stores and displays data from previously established land treatments or what often are called legacy data.			USGS			
CA Class I Wildfire	Fire Effects Information System	review not needed	The Fire Effects Information System is a compendium of research reports and other publications relating the effects of fire on native plant and animal species, invasive species, ecological communities, and soils.			USFS		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose.
CA Class I Wildfire	National Interagency Fuels, Fire, and Vegetation Technology Transfer (NIFTT)	review not needed	The NIFTT provides a suite of tools and documents on fire effects, fire and fuels management, and fire ecology.			USGS		These data will be used to inform the fire frequencies and extent parameters in the quantitative terrestrial (VDDT) models.	These data are suitable, in association with other data, for their intended purpose.
CA Class I Wildfire	National LANDFIRE Vegetation Dynamics Models	review finished	These are VDDT models for all terrestrial systems as BpS units with some natural fire regime.		accepted	LANDFIRE		These models, and their supporting data, are intended to provide foundational information for the quantitative terrestrial models produced for	These models, and their supporting data, are suitable for foundational information. The models were created to study historic vegetation patterns and dynamics. As a result,

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
								the CBR and MBR ecoregions.	they do not include unique anthropogenic ecological states, and thus are not suitable for inclusion into the model library without review and revision.
CA Class I Wildfire	TNC Updated Landfire Vegetation Dynamics models	review finished	these are models created for the Great Basin by TNC science staff.		accepted	TNC			
CA Class II Development	2009 Cropland Data Layer	review not needed	See Common Land Unit. This data is produced by the Farm Service Agency they are call CLU files (Common Land Units). Now done for the entire US. Check the USDA Geospatial Gateway website for download. Another option is to contact the FSA coordinator for the state you are interested in and request the statewide shapefile. This data is UNAVAILABLE per BLM and Farm Services.		(empty)	Farm Service Agency & NASS	No	Not intended for use.	
CA Class II Development	Agriculture Census of the United States	review not needed	This map layer portrays a selected set of information that was collected for the 2002 Census of Agriculture by the National Agricultural Statistics Service, U.S. Department of Agriculture.	USDA. 2007. Agriculture Census of the United States. US Department of Agriculture, National Agricultural Statistics Service. <a href="http://www.agcensus.usda.gov/">http://www.agcensus.usda.gov/</a>	rejected	USDA	Yes	Not intended for use.	
CA Class II Development	Alternative Fuels Stations	review not needed	The Alternative Fuels database is a geographic point database of fueling facilities that offer fuels other than gasoline in the United States.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	AM (zip) (07-31-2009)	in review	Extract of AM Radio Station Transmitter sites.		(empty)	FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Amtrak Stations	review not needed	This database is a geographic data set containing Amtrak intercity railroad passenger terminals in the United States and Canada.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Antenna Structure Registration (ASR) (zip) (07-26-2009)	in review	Extract of FCC Antenna Structure Registration database.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Automatic Traffic Recorder Stations	review not needed	The data included in the GIS Traffic Stations Version database have been assimilated from station description files provided by FHWA for Weigh-in-Motion (WIM), and Automatic Traffic Counters (ATR).			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Biomass Potential (2005)	review not needed	Biomass resource potential for the lower 48 states of the United States of America.	NREL. 2005. A Geographic Perspective on the Current Biomass Resource Availability in the United States. <a href="http://www.nrel.gov/docs/fy06osti/39181.pdf">http://www.nrel.gov/docs/fy06osti/39181.pdf</a>	rejected	NREL	No	Not suitable for use.	
CA Class II Development	Biomass Potential (2008)	review finished	Biomass Resources in the United States	NREL. 2008. Biomass Resources in the United States. <a href="http://www.nrel.gov/docs/fy06osti/39181.pdf">http://www.nrel.gov/docs/fy06osti/39181.pdf</a>	rejected	NREL	Yes	Not intended for use.	Not suitable.
CA Class II Development	BLM Linear Disturbance Maps	need to review	Linear disturbance (Roads, Trails)		accepted	BLM	No		
CA Class II Development	Cellular (zip) (07-26-2009)	in review	Extract of Cellular Radiotelephone Service sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Cities and Towns of the United States	review not needed	This map layer includes cities in the United States, Puerto Rico and the U.S. Virgin Islands.			USGS	Yes	Intended as reference only	Suitable for reference only.
CA Class II Development	Housing Density Change co_pbg00 (Colorado Dataset)	review not needed	The overarching goal of this analysis was to create a long-term dataset on housing density change that is accurate, spatially detailed, and consistent across the United States.			USDA, et al	Yes	Not intended for use.	
CA Class II Development	Census Block Attributes coblk00 (Colorado Dataset)	review not needed	U.S. Census blocks with selected attribute information.			Center for International Earth Science Information Network (CIESIN)	Yes	Not intended for use.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Developable Area and Strata Unit Area	in review	This dataset represents the "most geologically prospective" area for oil shale and allowable leasing footprints for tar sand extraction in Special Tar Sands Areas.		(empty)	Argonne National Laboratory	Yes		
CA Class II Development	Photovoltaic Solar Resource	review finished	Monthly and annual average solar resource potential for 48 Contiguous United States utilizing a Direct Normal collection method.	NREL. 2008. Photovoltaic Solar Resource Map of the United States. National Renewable Energy Laboratory. <a href="http://www.nrel.gov/gis/data_analysis.html">http://www.nrel.gov/gis/data_analysis.html</a>	rejected	NREL	Yes	Not intended for use.	
CA Class II Development	Dumps and landfills	review finished	Locations of landfills and waste transfer stations in 11 western states. Data was obtained from state and federal agencies in GIS, tabular, and map format.		accepted	USGS	Yes	This data set will be further evaluated in task three and compared against LU/LC data for accuracy and other proxy data sets.	While the data confidence rating for this data set is low, it represents the only data set of its kind. This data was created as part of the USGS Sagemap effort which helps add credibility despite the lack of documentation.
CA Class II Development	Energy Distribution Control Facilities	review finished	The Energy Distribution Control Facilities layer depicts the facilities which are responsible for balancing the load within their respective control areas. The proper functioning of these facilities is integral to the stability of the North American Elec		accepted	Global Energy Maps	Yes	Data not intended for use.	
CA Class II Development	Oil_Gas Potential EPCA 3	in review	inventory of all onshore Federal lands to identify: "the United States Geological Survey estimates of the oil and gas resources underlying these lands; and "the extent and nature of any restrictions or impediments to the development of the resources..."	DOI. 2008. Inventory of Onshore Federal Oil and Natural Gas Resources and Restrictions to Their Development. Prepared by the U.S. Departments of the Interior, Agriculture and Energy. <a href="http://www.blm.gov/wo/st/en/prog/energy/oil_and_gas/EPCA_III.html">http://www.blm.gov/wo/st/en/prog/energy/oil_and_gas/EPCA_III.html</a>	(empty)	BLM	No		
CA Class II Development	FEMA Transmission Line Connectivity	review finished	NREL received this data from the Federal Emergency Management Agency (FEMA) sometime around 1993. It is our understanding that the data represents a schematic of transmission line connectivity.		rejected	FEMA	No	Not intended or applicable for use.	
CA Class II Development	Fixed-Guideway Transit Facilities (Line)	review not needed	Version 2004 of the Fixed-Guideway Transit Network is a network database of the nation's fixed-guideway transit systems.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Fixed-Guideway Transit Facilities (Stations)	review not needed	Version 2004 of the Fixed-Guideway Transit Network is a network database of the nation's fixed-guideway transit systems.			Bureau of Transportation Statistics	Yes	Not intended for use.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	FM (zip) (07-31-2009)	in review	Extract of FM Radio Station Transmitter sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Freight Analysis Network	review not needed	"Freight Analysis Framework 2.2 Network Machine Readable Data Files" are distributed by the Federal Highway Administration Office of Freight Management and Operations, Operations Core Business Unit, Washington DC, 2007 and contains National Highway System			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Gas pipelines	need to review	The U.S. Department of Transportation (U.S. DOT), Pipeline and Hazardous Materials Safety Administration (PHMSA) is working with other federal and state agencies and the pipeline industry to create a National Pipeline Mapping System (NPMS).		(empty)	U.S. Dept. of Transportation - Pipeline and Hazardous Materials Safety Administration	Yes		
CA Class II Development	Groundwater well locations	need to review	groundwater well locations for residential houses to get at growth trends and patterns of rural development		(empty)				
CA Class II Development	Hazardous Material Routes	review not needed	The Federal Motor Carrier Safety Administration (FMCSA) Hazardous Material Routes were developed using the 2004 First Edition TIGER/Line files.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Highway Performance Monitoring System	review not needed	The Federal Highway Administration (FHWA) has the responsibility to assure that adequate highway transportation information is available to support its functions and responsibilities, including those of the Administration and the Congress.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Highway-Rail Grade Crossings	review not needed	FRA Grade Crossings is a spatial file that originates from the National Highway-Rail Crossing Inventory Program.			Bureau of Transportation Statistics	Yes	Not intended for use.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Intermodal Terminal Facilities	review not needed	This is a public dataset for the Department of Transportation, Research and Innovative Technology Administration's Bureau of Transportation Statistics.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Land Mobile - Broadcast (zip) (07-26-2009)	in review	Extract of Land Mobile Broadcast Service Transmitter sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Land Mobile - Commercial (zip) (07-26-2009)	in review	Extract of Land Mobile Commercial Service Transmitter sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Land Mobile - Private (zip) (07-26-2009)	in review	Extract of Land Mobile Private Service Transmitter sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	LATITL	review finished	Monthly and annual average solar resource potential for 48 Contiguous United States utilizing a Flat Plate Tilted South at Latitude collection method.		rejected	NREL	Yes	Not intended for use.	
CA Class II Development	Market significant transmission lines in North America.	review finished	The Transmission Lines layer is a comprehensive layer consisting of market significant transmission lines in North America. Depicted lines are generally greater than 115 kV and tie major power plants to the electrical grid. Transmission lines are located		accepted	Global Energy Maps	Yes	This layer is intended to represent market significant electricity transmission lines.	This layer is suitable for use however additional transmission line data is being sought.
CA Class II Development	Microwave (zip) (07-26-2009)	in review	Extract of Microwave Service sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	National Bridge Inventory	review not needed	The NBI is a collection of information (database) covering the more than 600,000 bridges located on public roads, including Interstate Highways, U.S. highways, State and county roads, as well as publicly-accessible bridges on Federal lands.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	National Highway Planning Network	review not needed	The National Highway Planning Network is a comprehensive network database of the nation's major highway system.			Bureau of Transportation Statistics	Yes	Not intended for use.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	National Land Cover Dataset (NLCD)	review not needed				MRLC	Yes		
CA Class II Development	Natural Landscapes (Theobald 2010)	review finished		Theobald, D.M. 2010. Estimating changes in natural landscapes from 1992 to 2030 for the conterminous United States. Landscape Ecology 25(7): 999-1011.	accepted		Yes	These data are intended to be used for broad-scale assessments of ecological integrity and as an indication of human modification of landscapes.	Natural landscapes (Theobald 2010) is a multi-scale, integrated metric that incorporate national datasets on land cover, housing density, road existence, and highway traffic volume to measure the dynamics of natural landscapes in the conterminous US. The NL metric is similar to other approaches that evaluate the effect of humans on natural landscapes such as the human footprint (Leu et al. 2008) in that it uses surrogate spatial data on land cover, population, and roads, as well as relying on heuristically derived estimates of human-dominated cover types. NL differs in that it is a simpler metric that has a direct physical interpretation related to proportion of natural cover at a location, examines the broader, landscape-scale pattern to differentiate the spatial context, and assumes that impacts decline continuously as a function of distance, rather than using abrupt buffers. NL also does not rely on pre-established critical scales and avoids the persistent problem of the arbitrariness of defining a patch. As such, this database is recommended as a summary or overview measure of human modification of landscapes, for the Development Change Agent.
CA Class II Development	NCEP Climate Datasets	need to review	geopotential height, u-wind,v-wind, vector wind, omega,air temperature, potential temperature, SST, specific hum, rel humidity, slp, surface pressure, precipitable water, precipitation rate,runoff, soil mositure, streamfunction, velocity potential, diverg			NCEP, NCAR			

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Nighttime Lights of North America	review not needed	This map layer is an image of nighttime lights for North America, including the Caribbean and most of Mexico.			Defense Meteorological Satellite Program (DMSP)	Yes	At a national scale, Nighttime Lights is an adequate for representing urban areas on the US landscape. However its resolution is too coarse for ecoregional use.	Not intended for use.
CA Class II Development	North American Atlas - Populated Places	review not needed	The North American Atlas - Populated Places data set shows a selection of named populated places suitable for use at a scale of 1:10,000,000.			USGS	Yes	Not intended for use.	
CA Class II Development	Oil and Gas Leases and Agreements	review finished	Shows federal current oil and gas leases, agreements, and lease sale parcels in the U.S on federal lands or where lands have been pooled with non-federal lands in the case of an agreement.		rejected	BLM	No	Need to reevaluate after metadata is obtained.	
CA Class II Development	Paging (zip) (07-26-2009)	in review	Extract of Paging Service Transmitter sites.			FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	Potential Geothermal Area	review finished	This coverage shows the regions favorable for the discovery and shallow depth (less than 1000m) of thermal water of sufficient temperature for direct-heat applications.		rejected	Idaho National Engineering & Environmental Laboratory	Yes	Not intended for use.	
CA Class II Development	Public Use Airport Runways	review not needed	The Airport Runways database is a geographic dataset of runways in the United States and US territories containing information on the physical characteristics of the runways.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Public-Use Airports	review not needed	The Airports database is a geographic point database of aircraft landing facilities in the United States and U.S. Territories.			Bureau of Transportation Statistics	Yes	Not intended for use.	
CA Class II Development	Railroads	review finished	The North American Atlas - Railroads data set shows the railroads of North America at 1:10,000,000 scale.		rejected	USGS	Yes	Not intended for use.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Railway Network (Line)	review finished	The Rail Network is a comprehensive database of the nation's railway system at the 1:100,000 scale.		accepted	Bureau of Transportation Statistics	Yes	This layer adequately represents the railway network at an ecoregional scale.	This layer is suitable for use.
CA Class II Development	Railway Network (Node)	review not needed	The Rail Network is a comprehensive database of the nation's railway system at the 1:100,000 scale.		(empty)	Bureau of Transportation Statistics	Yes	Not intended for use.	Not intended for use.
CA Class II Development	Section 368 Energy Corridors	review finished	Represents areas which have been proposed as West-wide energy corridors.	DOE & BLM. 2008. Programmatic Environmental Impact Statement, Designation of Energy Corridors on Federal Land in the 11 Western States (DOE/EIS-0386). <a href="http://corridoreis.anl.gov/documents/fpeis/index.cfm">http://corridoreis.anl.gov/documents/fpeis/index.cfm</a>	accepted	Argonne National Laboratory	Yes	This data belongs to a larger category of development change agents, specifically planned areas of electrical transmission. It will be used to represent areas of likely land use change and investment in energy infrastructure.	This data set is suitable for its intended purpose.
CA Class II Development	Significant Electric Power Generation Plants	need to review	The Electric Plants layer is a comprehensive representation of significant power plants within the North American power grid. The majority of plants shown are greater than three megawatts. Power plants are located using a mixture of sources from regional		(empty)	Global Energy Maps	Yes	Not intended for use.	Not intended for use.
CA Class II Development	Spatially Explicit Regional Growth Model (SERGoM) v1.2	review finished	SERGoM data uses US Census block housing units, protected lands, groundwater well density, and road accessibility to estimate housing density	U.S. Environmental Protection Agency (EPA; Bierwagen, B., D.M. Theobald, C.R. Pyke, A. Choate, P. Groth, J.V. Thomas, and P. Morefield). 2009 Land-Use Scenarios: National-Scale Housing-Density Scenarios Consistent with Climate Change Storylines. Global Change Research Program, National Center for Environmental Assessment, Washington, DC; EPA/600/R-08/076F. Bierwagen, B., D.M. Theobald, C.R. Pyke, A. Choate, P. Groth, J.V. Thomas, and P. Morefield. (In press, accepted 12 October 2010). Land-Use Scenarios: National-Scale Housing-Density Scenarios Consistent with Climate Change Storylines. Proceedings of the National Academy of Sciences. Theobald, D.M. 2005. Landscape patterns of exurban growth in the USA from 1980 to 2020. Ecology and Society 10(1): 32. [online] URL:	accepted	Theobald and US EPA	Yes	Main layer of urban-to-rural patterns of development for Development Change Agent.	The ICLUS (Integrated Climate Land Use System) project has developed national scenarios of housing density that are logically consistent with IPCC emissions storylines. It uses a cohort-component methodology to represent population growth in the US. Spatial allocation is accomplished using SERGoM (4), a hierarchical (national to state to county), deterministic model that calculates the number of additional housing units needed in each county to meet the demand specified by population projections from the demographic model, based on the ratio of housing units to population (downscaled from census tract to block). Housing units are spatially allocated within a county in response to the spatial pattern of land ownership, previous

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
				<a href="http://www.ecologyandsociety.org/vol10/iss1/art32/">http://www.ecologyandsociety.org/vol10/iss1/art32/</a> .					growth patterns, and travel time accessibility. The model is dynamic in that as new urban core areas emerge, the model re-calculates travel time from these areas. SERGoM used refined land ownership, transportation, and groundwater well density using 2009 data, and by weighting housing units by NLCD 2001 cover types (Theobald 2005; US EPA 2009; Bierwagen et al. in press). Other datasets that are suggested for development change agent include SILVIS housing density and LANDSCAN, but these are not based on open source demographic/population projections and do not include the detailed spatial data on land ownership, accessibility, and groundwater density to allocate housing units. They are based on block-group level allocation, whereas SERGoM is based on modified block-level (a finer grain dataset). The ICLUS/SERGoM layer is adequate for use in the REA.
CA Class II Development	Substations and Taps in North American Power Grid	need to review	The Substations layer is a comprehensive layer of the substations and taps that exist in the North American power grid. Substations are snapped into segments of the Transmission Lines layer and are found at every power plant. Substations are located using		(empty)	Global Energy Maps	Yes	Not intended for use.	
CA Class II Development	TIGER 2009 "edges" and roads	need to review	Comprehensive road layer for the ecoregion					TIGER line files and edges is used to represent linear development features such as roads. This layer may be used in BLM Linear Disturbance or USGS 1:24,000 DLG data is unavailable.	Generally not suitable but may be used as a backup.
CA Class II Development	Trails	review not needed	Have historic trails, Pacific Crest			BLM		Not intended for use. Trails will be represented with BLM Linear Disturbance maps.	

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Travel management, OHV use	need to review			accepted				
CA Class II Development	TV - Digital (zip) (07-31-2009)	in review	Extract of NTSC Television Station Transmitter sites.		(empty)	FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	TV - NTSC (zip) (07-31-2009)	in review	Extract of Digital Television Station Transmitter sites.		(empty)	FCC Media Bureau	No	May be used in conjunction with BLM Linear Features maps, energy transmission and others to represent disturbance features on the landscape.	This data set requires metadata to be thematically and technically suitable for the intended use.
CA Class II Development	U.S. Census Database, 1990	review not needed	This data set includes U.S. Census Bureau 1990 population information for the United States, presented by county.			Census	Yes	Not intended to be used directly. See SERGoM/ICLUS.	
CA Class II Development	U.S. Census Database, 2000	review not needed	This data set includes U.S. Census Bureau population information for the United States and Puerto Rico, presented by county.			Census	Yes	Not intended to be used directly. See SERGoM/ICLUS.	
CA Class II Development	Urban Areas of the United States	review not needed	This data set includes a selection of urban areas in the United States derived from the urban areas layer of the Digital Chart of the World (DCW).			USGS	Yes	Not intended for use.	
CA Class II Development	US Roads	review not needed	This data set portrays the major roads in the United States, Puerto Rico, and the U.S. Virgin Islands		rejected	USGS	Yes	Not intended for use.	
CA Class II Development	USFS National Visitor Use Monitoring	review finished			rejected	USDA Forest Service		These data are useful to understand broad-scale (Forests to regional) understanding of recreation use on Forest Service land, but are limited for the spatial assessments for the REA because similar data are not available on BLM, NPS, and USFWS and other public lands. Also, it is difficult to extrapolate to a finer-scale that would be needed for the REAs.	This data is suitable as a reference source only.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Water Use by County	review not needed	This map layer portrays the estimated use of water in counties in the United States, in the year 2000.			USGS	Yes	Not intended for use.	
CA Class II Development	Wildland Urban Interface	need to review	The Wildland-Urban Interface (WUI) is the area where houses meet or intermingle with undeveloped wildland vegetation.			SILVIS Lab, Department of Forest Ecology and Management, University of Wisconsin-Madison	Yes		
CA Class II Development	Known Geothermal Resource Areas, Geothermal Lease Status, Biomass Development Areas, Concentrating Solar Power, Flat plate collector solar resource data, wind power classes	need to review	Assessing The Potential For Renewable Energy On Public Lands Report (DOE/GO-102003-1704 ) GIS Datasets on CD-ROM available at listed website.			NREL and BLM	Yes		
CA Class II Development	50m Wind Potential	in review	Wind power potential for the states at a 50 meter height. This dataset will be replaced when the southwest region has been completed, and the data may change when this region has been completed.	NREL. 1986. Wind Energy Resource Atlas of the United States. National Renewable Energy Laboratory. <a href="http://rredc.nrel.gov/wind/pubs/atlas/">http://rredc.nrel.gov/wind/pubs/atlas/</a>		TrueWind Solutions/NREL	Yes		
CA Class II Development	Solar Energy Study Areas	review finished	This data represents Solar Energy Study Areas developed by the Bureau of Land Management for use in the Solar Energy Programmatic Environmental Impact Statement (PEIS). The areas have been selected as being free of land use restrictions and for their suitability as sites for utility grade solar power plants. For details see the Solar Energy PEIS at <a href="http://solareis.anl.gov">http://solareis.anl.gov</a> .		accepted	BLM	Yes	This data set represents solar energy areas that are most likely to be developed in the short term.	This data set is suitable for its intended purpose.
CA Class II Development	Mineral Resource Data System	in review	MRDS describes metallic and nonmetallic mineral resources throughout the world. It is a vector point file. Included are deposit name, location, commodity, deposit description, geologic characteristics, production, reserves, resources, and references. It includes the original MRDS and MAS/MILS data.		(empty)	USGS	Yes	This data set will represent relative impact by past mining activity.	Dataset may be the best available. Being point data this dataset lacks a spatial component that reflects the total surface footprint of a mine or mine processing site.

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class II Development	Ruby Pipeline	need to review	Spatial layer representing the 677-mile Ruby natural gas pipeline across Wyoming, Utah, Nevada, Oregon and California						
CA Class II Development	Wind resource map, mean annual wind speed at 80m height	need to review	The Department of Energy's Wind Program and the National Renewable Energy Laboratory (NREL) published a new wind resource map showing the predicted mean annual wind speeds at 80-m height.	AWS Truewind & NREL. 2009. Predicted mean annual wind speeds at 80-m height. AWS Truewind & National Renewable Energy Laboratory. <a href="http://www.windpoweringamerica.gov/index.asp">http://www.windpoweringamerica.gov/index.asp</a>		NREL			
CA Class II Development	Concentrated Solar Power Resource Maps	need to review	These direct-normal solar radiation maps filtered by solar resource, land availability and suitability. Identifies the most economically suitable lands available for deploying of large-scale concentrating solar power plants in the southwestern United States.	NREL. 2010. Concentrating Solar Power Resource Maps. <a href="http://www.nrel.gov/csp/maps.html">http://www.nrel.gov/csp/maps.html</a>		NREL			
CA Class II Development	current locations of private and state land renewable energy facilities	need to review	Current location and footprint of existing renewable energy facilities.						
CA Class II Development	Preliminary Geothermal Potential and Exploration in the Great Basin	review finished	This map provides regional information for assessing the potential for high-temperature (>150 deg. C) geothermal systems in the Great Basin- those most likely to be capable of producing electrical energy.	Zehner, R, M Coolbaugh, L Shevenell. 2009. Preliminary Geothermal Potential and Exploration Activity in the Great Basin. Nevada Bureau of Mines and Geology, University of Nevada, Reno.	accepted	Nevada Bureau of Mines and Geology	Yes	This layer will represent geothermal potential for the Central Great Basin and northern Mojave Basin areas.	The data is suitable for the intended use.
CA Class II Development	Geothermal leases	in review	Includes three sets of data: Geothermal leases closed, producing and nonproducing.			BLM	No		
CA Class II Development	Solar Energy Leases	in review				BLM	No		
CA Class II Development	Solid Mineral Leases	in review				BLM	No		
CA Class II Development	Wind Energy Leases	in review				BLM	No		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class III Invasive Species	Annual Grass Index of Nevada (March 2006)	need to review	Arc Grid. 100% of Nevada plus edges of adjacent states. Currency of data: effectively spring 2004/2005. Scale at which data are believed to meet National Map Accuracy Standards: 1:100,000 in most areas. Recommended that the map presented here be interpreted as an annual grass index (ANGRIN) map, rather than an estimate of actual annual grass cover. Nevertheless, the ANGRIN map clearly reveals the pattern of annual grass invasion across Nevada.	Peterson, E. B. 2006. A map of invasive annual grasses in Nevada derived from multitemporal Landsat 5 TM imagery. Report for the U.S.D.I. Bureau of Land Management, Nevada State Office, Reno, by the Nevada Natural Heritage Program, Carson City, Nevada.	(empty)	Nevada Natural Heritage Program	No		
CA Class III Invasive Species	Wild Horse and Burro Herd Areas	need to review			(empty)	BLM			
CA Class III Invasive Species	Invasive Species Infestation location	in review	Polygon feature data set that depicts noxious weed distribution across the western united states. This data supports the noxious weed monitoring and training within the National Invasive Species Information Management System.		(empty)	BLM	Yes	Without species information, this data set may represent a general infestation level by weed species.	Need to clarify with BLM that this data set does not distinguish between species. There are relevant data fields that get at percent cover, extent, etc but nothing relating to species. Need to determine this before determining the intended use and suitability of the data set.
CA Class III Invasive Species	Invasive Species Survey Area	need to review	We didn't receive the data from BLM due to file corruption issues, so cannot assess. The Data source links lead to Geo-Energy web site, which doesn't make sense.		(empty)	BLM			
CA Class III Invasive Species	Boundaries of Invasive Species Treatment Areas	need to review	Have not yet received the data from BLM due to data corruption issues.		(empty)	BLM			
CA Class III Invasive Species	Weed Management Areas	review finished	This data set represents BLM or perhaps multi-agency weed management areas.		(empty)	BLM	No	Use not clear. May be used as a reporting unit.	No metadata was recieved with this layer. The suitability may be acceptable if used solely as a reporting unit.
CA Class III Invasive Species	Non-Native Aquatic Invasive	need to review				USGS	Yes		
CA Class III Invasive Species	New Zealand Mudsail Sightings Distribution: USGS NAS	need to review	This map layer is a compilation of confirmed New Zealand mudsail sighting reports in the United States and Canada from 1987 through 2010 and is updated daily. It provides geographical and historical information to show distribution over space and time. Although it is updated daily it is dependent of reported confirmed sightings which may not be reported daily		accepted	USGS Nonindigenous Aquatic Species	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
CA Class III Invasive Species	Zebra Mussel Locations: USGS NAS	need to review	Mapsite of reported Zebra Mussel locations in USA including our ecoregions. Although it is reported to be updated daily, it is dependent on those reporting zebra mussels to report to this mapping website.		accepted	USGS Nonindigenous Aquatic Species website	Yes		
CA Class III Invasive Species	Quagga Mussel Distribution Map: USGS NAS	need to review	Map data of reported locations of quagga mussels		accepted	USGS Nonindigenous Aquatic Species website	Yes		
CA Class III Invasive Species	New Zealand mudsnail in the Western USA: Montana State U.	need to review	This db is superior to USGS NAS NZMS db but has not been updated since 2009. There are substantially more point locations than USGS with more detailed descriptions			Montana State University	Yes		
CA Class III Invasive Species	USGS Nonindigenous Species database: USGS NAS bullfrog example	need to review	This is an example from our default aquatic invasive species database at the USGS NAS website. The website database has almost all of the aquatic invasives on our list, but I am not sure how 'up to date' it really is. The page source site links to is for bullfrogs and list occurrences by states and HUCs. It also has a link to specific reported locations. It wont be difficult to access all the vital info when the time comes		accepted	USGS	Yes		
CA Class III Invasive Species	Didymo (Didymosphenia geminata) distribution map: USGS Fort Collins	need to review	This is a generalized map with dots indicating didymo presence. Dr. Sarah Spaulding who is the US leading expert on didymo is providing database coordinates that were used for this map and any updated locations. Dr. Spaulding is requesting funding from BLM to update the didymo database		accepted	USGS Fort Collins Science Center	Unknown		
CA Class III Invasive Species	Zebra mussel, quagga mussel and Asian clam veliger locations: EcoAnalysts, Moscow, ID	need to review	EcoAnalysts has just about completed an analysis of water samples collected from a few hundred sites for and by NVDOW that were examined for invasive mussel and clam veligers (tiny babies). At this time the data is considered 'confidential' without permission for use from NVDOW. If dataset looks promising we will ask for permission to use.			Nevada Department of Wildlife	Unknown		
CA Class III Invasive Species	Nevada Noxious Weeds Data	need to review				Nevada Natural Heritage Program			
CA Class III Invasive Species	Cheatgrass (Bromus tectorum) Estimated Percent Cover (December 2003)	need to review	The mapping method involved developing a statistical model for the estimation of B. tectorum cover at training plots with variables derived from Landsat 7 ETM+ satellite data satellite imagery and matching topographic data.	Peterson, E. B. 2003. Mapping Percent-Cover of the Invasive Species Bromus tectorum (Cheatgrass) over a Large Portion of Nevada from Satellite Imagery. Report for the U.S. Fish and Wildlife Service, Nevada State Office,		Nevada Natural Heritage Program	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
				Reno, by the Nevada Natural Heritage Program, Carson City.					
CA Class III Invasive Species	SWEMP--Southwest Exotic Plant Mapping Project	in review	The database represents the known point locations of non-native invasive plant infestations within Arizona and New Mexico, and adjacent portions of California, Colorado, Nevada and Utah. These data, collected from 1911 to 2006. Data includes all counties in NV, UT, and CO, and the 5 southern counties of CA.	Paxton, E.H., M. Sogge, T. Theimer, J. Girard, & P. Keim. 2008. Relevant Invasive Species Program Goals and Invasive Species Related Highlights & Key Findings and Accomplishments. USGS pub?		Arizona Heritage Program	Yes	Degree of conversion by invasive species to assess the amount of stress on natural ecosystems	High
CA Class III Invasive Species	Nevada Cheatgrass Project	in review	Point location with presence/absence for Bromus tectorum in Central Nevada	Bradley, B.A., and J.F. Mustard, "Characterizing the Landscape Dynamics of an Invasive Plant and Risk of Invasion Using Remote Sensing", Ecological Applications, 16(3), 1132-1147, 2006 1. Brte_NV.shp 2006-11-8 12:14, uploaded by Bethany Bradley on November 8th, 2006 Bradley, B.A., and J.F. Mustard. 2005. Remote Sensing of Environment. 94, 204-213					
CA Class IV Climate Change	DayMet	review not needed			(empty)	Oak Ridge National Lab	Yes		
CA Class IV Climate Change	800 m PRISM Monthly Precipitation	need to review				Oregon State	Yes		
CA Class IV Climate Change	Bioclimate Classes: Thermotype and Ombrotype	review not needed	Isobioclimates were generated by combining the thermotypes (warm/cold) and ombrotype (dry/wet gradients) climate classes produced from the Rivas-Martínez method based on the concept of a quantifiable classification system which would closely relate the di			USGS	Yes		
PL Class I Sites of High Biodiversity	Nevada priority conservation areas	need to review	Areas identified through field inventory by the state Natural Heritage Program			Nevada Natural Heritage Program			
PL Class I Sites of High Biodiversity	TNC Ecoregional Assessment - 2010	review finished	Relative Conservation Value as documented by the 2010 updated Mojave Desert ecoregional assessment of The Nature Conservancy	Randall, J.M. SS. Parker, J. Moore, B. Cohen, L. Crane, B. Christian, D. Cameron, J. MacKenzie, K. Klausmeyer and S. Morrison. 2010. Mojave Desert Ecoregional Assessment. Unpublished Report. The Nature Conservancy, San Francisco,	accepted	The Nature Conservancy (NV, CA, AZ)	Yes	Potential use as assessment units; i.e., current and future conditions relative to these selected landscapes of biodiversity significance.	Suitable for this use. See updated version from Mojave (2010).

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
				California. 106 pages + appendices. Available at <a href="http://conserveonline.org/workspaces/mojave/documents/mojave-desert-ecoregional-2010/@@view.html">http://conserveonline.org/workspaces/mojave/documents/mojave-desert-ecoregional-2010/@@view.html</a>					
PL Class I Sites of High Biodiversity	Audubon Important Bird Areas					Audubon			
PL Class I Sites of High Biodiversity	Important Bird Areas - American Bird Conservancy	need to review				American Bird Conservancy			
PL Class I Sites of High Biodiversity	TNC Portfolio Sites	review finished	Portfolio sites identified through ecoregional plans of TNC from late 1990s-early 2000s.	for CBR: Nachlinger, J., K. Sochi, P. Comer, G. Kittel, and D. Dorfman. 2001. Great Basin: an ecoregion-based conservation blueprint. The Nature Conservancy, Reno, NV. 160 pp. + appendices. For MBR: Moore, J., C. Rumsey, T. Knight, J. Nachlinger, P. Comer, D. Dorfman, and J. Humke. 2001. Mojave Desert: an ecoregion-based conservation blueprint. The Nature Conservancy, Las Vegas, NV. 150 pp. + appendices.	accepted	The Nature Conservancy			
PL Class II Specially Designated Areas of Ecological Value	ACEC		will derived from BLM directly		(empty)	BLM			
PL Class II Specially Designated Areas of Ecological Value	Wild Horse and Burro Herd Management Areas	need to review			accepted	BLM			
PL Class II Specially Designated Areas of Ecological Value	National Inventoried Roadless Areas (IRAs)	need to review	This dataset contains all National Forest Inventoried Roadless Areas (IRAs) for the lower 48 states, including Puerto Rico.			USDA	Yes		

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
PL Class II Specially Designated Areas of Ecological Value	BLM National landscape Conservation System (NLCS)	review finished	The Bureau of Land Management's National Landscape Conservation System (NLCS) contains some of the West's most spectacular landscapes. It includes over 886 federally recognized areas and approximately 27 million acres of National Monuments, National Conservation Areas, Wilderness Areas, Wilderness Study Areas, Wild and Scenic Rivers, National Scenic and Historic Trails, and Conservation Lands of the California Desert.		(empty)	BLM	Yes	Suitable for reference only. Not suitable for analysis.	
PL Class II Specially Designated Areas of Ecological Value	Protected Areas Database (PAD) (BLM version)	review finished	Review BLM PAD. The Protected Areas Database of the United States (PAD-US) is a digital map of steward boundaries that combines attributes of ownership, management, and a measure of intent to manage for biodiversity.		(empty)	USGS	Yes	This data set is intended to identify designated areas of high biodiversity value and other managed lands for the ecoregion.	This data set is recommended for display or reference use only.
PL Class III Other Managed Lands	Livestock Grazing Allotments	in review	Grazing allotments and pastures by ecoregion		(empty)	BLM	No	This data may be linked to additional grazing data provided by the NOC. Otherwise will be treated as a reporting unit only.	The data is suitable as a reporting unit however the AMT has indicated that there are likely spatial errors in the dataset. The NOC may replace this or recommend another data set in the future.
PL Class III Other Managed Lands	BLM Admin Boundaries	review finished			accepted	BLM	Yes	Fine for reference purposes.	
PL Class III Other Managed Lands	Common Land Unit	review not needed	NO LONGER ACCESIBLE SINCE 2008 PER BLM. A Common Land Unit (CLU) is the smallest unit of land that has a permanent, contiguous boundary, a common land cover and land management, a common owner and a common producer in agricultural land associated with USDA farm programs. CLU boundaries are delineated from relatively permanent features such as fence lines, roads, and/or waterways.		(empty)	NRCS			
PL Class III Other Managed Lands	Counties	review not needed	County clip by ecoregion		accepted	BLM	No	This data is intended as reference only	The data is suitable for reference only.
PL Class III Other Managed Lands	Land Use Planning Boundaries	in review			(empty)	BLM	Yes	The data will be used as reporting or reference units.	This data is suitable for the intended use.
PL Class III Other Managed Lands	Land Use Planning Decision Boundaries	in review			accepted	BLM			

Primary Data Class	Dataset Name	Review Status	Data Description	Citation	Data Status After Review	Source Agency	Meta data	Intended Use of Data	Suitability for Intended Uses
PL Class III Other Managed Lands	Military expansion areas - Ft Irwin, 29 Palms	need to review							
PL Class III Other Managed Lands	Military flight areas	need to review							
PL Class III Other Managed Lands	Military Use areas	need to review							
PL Class III Other Managed Lands	Special management designations, recreation areas, boat ramps, etc.	need to review	Can probably use our NS PAD for management distinctions, may need to request recreation site info from BLM		(empty)				
PL Class III Other Managed Lands	State Boundaries	review not needed			accepted	BLM			
PL Class III Other Managed Lands	Surface Management Agency	need to review			accepted	BLM	No		
PL Class III Other Managed Lands	mining claims	need to review	Areas where mining claims have been filed but commercial production has not yet comenced						
Other	CEC Level 3	review not needed	This map layer shows Omernik's Level III ecoregions.			USAPA	Yes		
Other	Public Land Survey System	review not needed			(empty)	BLM	Yes		
Other	Visual resource Inventory								

**Appendix II. Coarse-filter Conservation Elements for the Central Basins and Ranges Ecoregion.**

Ecoregion Model Group	Land Cover Class	Conservation Element Name	Percent of Ecoregion	# of Field Referenced Samples	Vegetation Dynamics Models LANDFIRE	Vegetation Dynamics Models TNC NV	NatureServe Ecological Integrity Criteria 2008	NatureServe Ecological Integrity Criteria 2000
Montane Dry	Evergreen Forest and Woodland	Great Basin Pinyon-Juniper Woodland	13.8%	2200	yes	yes		yes
Montane Dry	Shrub-steppe	Inter-Mountain Basins Montane Sagebrush Steppe	3.9%	928	yes	yes	yes	yes
Montane Dry	Sparsely Vegetated	Inter-Mountain Basins Cliff and Canyon	0.7%	267			yes	
Montane Dry	Mixed Evergreen-Deciduous Forest and Woodland	Inter-Mountain Basins Curl-leaf Mountain Mahogany Woodland and Shrubland	0.6%	327	yes	yes		
Montane Dry	Deciduous Forest and Woodland	Rocky Mountain Aspen Forest and Woodland	0.2%	468	yes		yes	yes
Montane Dry	Evergreen Forest and Woodland	Inter-Mountain Basins Subalpine Limber-Bristlecone Pine Woodland	0.2%	51	yes	yes	yes	yes
Montane Dry	Mixed Evergreen-Deciduous Forest and Woodland	Inter-Mountain Basins Aspen-Mixed Conifer Forest and Woodland	0.0%	101	yes	yes	yes	yes
Basin Dry	Short Shrubland	Inter-Mountain Basins Mixed Salt Desert Scrub	20.0%	1907	yes	yes		yes
Basin Dry	Shrub-Steppe	Inter-Mountain Basins Big Sagebrush Shrubland	19.5%	2278	yes	yes	yes	
Basin Dry	Short Shrubland	Great Basin Xeric Mixed Sagebrush Shrubland	9.6%	1407	yes	yes		yes
Basin Dry	Shrub-steppe	Inter-Mountain Basins Semi-Desert Shrub-Steppe	3.1%	171	yes	yes	yes	yes
Basin Dry	Short Shrubland	Mojave Mid-Elevation Mixed Desert Scrub	2.0%	542	yes			
Basin Dry	Upland Grassland and Herbaceous	Inter-Mountain Basins Semi-Desert Grassland	1.0%	195	yes	yes	yes	yes
Basin Dry	Shrub-steppe	Inter-Mountain Basins Big Sagebrush Steppe	0.3%	549	yes	yes	yes	
Basin Dry	Sparsely Vegetated	Inter-Mountain Basins Active and Stabilized Dune	0.2%	31			yes	yes
Basin Dry	Dwarf-shrubland	Colorado Plateau Mixed Low Sagebrush Shrubland	0.1%	15	yes	yes		
Basin Dry	Tall Shrubland	Great Basin Semi-Desert Chaparral	0.0%	27	yes	yes		
Montane Wet	Woody Wetlands and Riparian	Great Basin Foothill and Lower Montane Riparian Woodland and Shrubland/Stream	1.1%	333		yes	yes	yes
Montane Wet	Woody Wetlands and Riparian	Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland/Stream	0.1%	14			yes	yes
Montane Wet	Woody Wetlands and Riparian	Rocky Mountain Subalpine-Montane Riparian Woodland/Stream	0.0%	46		yes	yes	yes
Montane Wet	Herbaceous Wetlands	Rocky Mountain Alpine-Montane Wet Meadow and Pond	0.0%	14		yes	yes	yes
Montane Wet	Woody Wetlands and Riparian	Rocky Mountain Subalpine-Montane Riparian Shrubland/Stream	0.0%	37		yes	yes	yes
Basin Wet	Sparsely Vegetated/Ephemeral Open Water	Inter-Mountain Basins Playa	5.7%	280			yes	
Basin Wet	Woody Wetlands and Riparian	Inter-Mountain Basins Greasewood Flat	5.1%	1705	yes	yes	yes	yes
Basin Wet	Open Water	Great Basin Lake/Reservoir	2.0%					
Basin Wet	Herbaceous Wetlands	North American Arid West Emergent Marsh and Pond	0.2%	189		yes	yes	yes
Basin Wet	Aquatic Point Location	Great Basin Springs and Seeps	0.0%	130				no

**Appendix III: Current Draft of Fine-Filter Conservation Elements for the Central Basin and Range REA**

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Dry	Amphibians	Inyo Mountains Salamander	Batrachoseps campi	No	No	G2	CA	CA		19	CA	No
Dry	Amphibians	Kern Plateau Salamander	Batrachoseps robustus	No	No	G2	CA			3		No
Dry	Amphibians	Western Toad	Bufo boreas	No	Yes	G4	UT	UT		144	SW, CA	No
Dry	Amphibians	Yosemite Toad	Bufo canorus	Yes	No	G2	CA			59	CA	No
Dry	Amphibians	Great Plains Toad	Bufo cognatus	No	Yes	G5	NV, UT	UT	PS	1	SW, CA	No
Dry	Amphibians	Black Toad	Bufo exsul	No	Yes	G1	CA	CA		6		No
Dry	Amphibians	Arizona Toad	Bufo microscaphus	No	Yes	G3	NV, UT	UT	PS	88	SW, CA	No
Dry	Amphibians	Amargosa Toad	Bufo nelsoni	No	Yes	G2	NV		PS	23	SW	No
Dry	Amphibians	Woodhouse's Toad	Bufo woodhousii	No	No	G5	ID					No
Dry	Amphibians	Mount Lyell Salamander	Hydromantes platycephalus	No	No	G3	CA			12	CA	No
Dry	Amphibians	Owens Valley Web-toed Salamander	Hydromantes sp. 1	No	No	G1	CA			2		No
Dry	Amphibians	Canyon Treefrog	Hyla arenicolor	No	No	G5	UT			3	SW	No
Dry	Amphibians	Pacific Chorus Frog	Pseudacris regilla	No	No	G5	UT			53	SW, CA	No
Dry	Amphibians	Columbia Spotted Frog	Rana luteiventris	Yes	Yes	G4	ID, NV, UT	UT	HV	160	SW	No
Dry	Amphibians	Columbia Spotted Frog - Great Basin	Rana luteiventris pop. 3	Yes	Yes	T2				303		No
Dry	Amphibians	Southern Mountain Yellow-legged Frog	Rana muscosa	Yes	No	G2	CA				CA	No
Dry	Amphibians	Relict Leopard Frog	Rana onca	Yes	Yes	G1	NV, UT		MV	6	SW	No
Dry	Amphibians	Sierra Nevada Yellow-legged Frog	Rana sierrae	No	No	G1	NV		PS	58		No
Dry	Ants, Wasps, & Bees	Lassen Chrysidid Wasp	Argochrysis lassenae	No	No	G1				1		No
Dry	Ants, Wasps, & Bees	A Montane Ant	Formica microphthalma	No	No	G2				2		No
Dry	Ants, Wasps, & Bees	Dune Honey Ant	Myrmecocystus snellingi	No	No	G2				4		No
Dry	Ants, Wasps, & Bees	Borrego Parnopes Chrysidid Wasp	Parnopes borregoensis	No	No	G1				1		No
Dry	Ants, Wasps, & Bees	An Ant	Stenamma wheelerorum	No	No	G1				1		No
Dry	Birds	Cooper's Hawk	Accipiter cooperii	No	Yes	G5	CA			1	SW, CA	No
Dry	Birds	Northern Goshawk	Accipiter gentilis	No	Yes	G5	CA, NV, UT	CA, UT	MV	163	SW, CA	No
Dry	Birds	Sharp-shinned Hawk	Accipiter striatus	No	Yes	G5	CA					No
Dry	Birds	Boreal Owl	Aegolius funereus	No	Yes	G5	ID, UT			2	SW	No
Dry	Birds	White-throated Swift	Aeronautes saxatalis	No	Yes	G5	NV					No
Dry	Birds	Tricolored Blackbird	Agelaius tricolor	No	Yes	G2	CA, NV	CA	PS	2	SW, CA	No
Dry	Birds	Grasshopper Sparrow	Ammodramus savannarum	No	Yes	G5	CA, ID, UT	UT		12	SW, CA	No
Dry	Birds	Sage Sparrow	Amphispiza belli	No	Yes	G5	NV, UT		MV	13	SW, CA	No
Dry	Birds	American Pipit	Anthus rubescens	No	Yes	G5						No
Dry	Birds	Golden Eagle	Aquila chrysaetos	No	Yes	G5	CA	CA, UT		9	SW, CA	No
Dry	Birds	Great Egret	Ardea alba	No	Yes	G5	CA, ID			5	SW	No
Dry	Birds	Short-eared Owl	Asio flammeus	No	Yes	G5	CA, ID, NV, UT	UT	PS	84	SW, CA	No
Dry	Birds	Long-eared Owl	Asio otus	No	Yes	G5	CA			9	SW, CA	No
Dry	Birds	Burrowing Owl	Athene cunicularia	No	Yes	G4	CA, ID, UT	CA, UT		496	SW, CA	No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Birds	Western Burrowing Owl	Athene cunicularia hypugaea	No	Yes	T4	NV	AZ	PS	5		No
Dry	Birds	Juniper Titmouse	Baeolophus ridgwayi	No	Yes	G5	ID				SW	No
Dry	Birds	Ruffed Grouse	Bonasa umbellus	No	Yes	G5	CA					No
Dry	Birds	Cattle Egret	Bubulcus ibis	No	Yes	G5	ID			1	SW	No
Dry	Birds	Ferruginous Hawk	Buteo regalis	No	Yes	G4	CA, ID, NV, UT	UT	PS	680	SW	No
Dry	Birds	Swainson's Hawk	Buteo swainsoni	No	Yes	G5	CA, ID, NV	CA	PS	389	SW, CA	No
Dry	Birds	Common Black-Hawk	Buteogallus anthracinus	No	Yes	G4				3	SW	No
Dry	Birds	Lark Bunting	Calamospiza melanocorys	No	Yes	G5				13	SW	No
Dry	Birds	Gambel's Quail	Callipepla gambelii	No	Yes	G5	UT					No
Dry	Birds	Costa's Hummingbird	Calypte costae	No	Yes	G5	CA, NV		IL	4	SW, CA	No
Dry	Birds	Cassin's Finch	Carpodacus cassinii	No	Yes	G5	NV					No
Dry	Birds	Turkey Vulture	Cathartes aura	No	Yes	G5				7	SW, CA	No
Dry	Birds	Veery	Catharus fuscescens	No	Yes	G5				11	SW	No
Dry	Birds	Swainson's Thrush	Catharus ustulatus	No	Yes	G5						No
Dry	Birds	Greater Sage-Grouse	Centrocercus urophasianus	Yes	Yes	G3	CA, ID, NV, UT	CA, UT	HV	73	SW, CA	Yes
Dry	Birds	Vaux's Swift	Chaetura vauxi	No	Yes	G5	CA					No
Dry	Birds	Snowy Plover	Charadrius alexandrinus	No	Yes	G4	UT	UT		33	SW	No
Dry	Birds	Western Snowy Plover	Charadrius alexandrinus nivosus	No	Yes	T3	CA, NV		MV	28		No
Dry	Birds	Mountain Plover	Charadrius montanus	Yes	Yes	G3	CA, UT	AZ, CA, UT		10	SW	No
Dry	Birds	Lark Sparrow	Chondestes grammacus	No	Yes	G5	CA					No
Dry	Birds	Lesser Nighthawk	Chordeiles acutipennis	No	Yes	G5				10	SW, CA	No
Dry	Birds	Northern Harrier	Circus cyaneus	No	Yes	G5	CA			3	SW, CA	No
Dry	Birds	Marsh Wren	Cistothorus palustris	No	Yes	G5						No
Dry	Birds	Evening Grosbeak	Coccothraustes vespertinus	No	Yes	G5				9	SW, CA	No
Dry	Birds	Yellow-billed Cuckoo	Coccyzus americanus	Yes	Yes	G5	ID, UT	UT		38	SW, CA	No
Dry	Birds	Western Yellow-billed Cuckoo	Coccyzus americanus occidentalis	Yes	Yes	T3	CA, NV	CA	MV	22		No
Dry	Birds	Black-billed Cuckoo	Coccyzus erythrophthalmus	No	Yes	G5				2	SW	No
Dry	Birds	Inca Dove	Columbina inca	No	Yes	G5				1	SW, CA	No
Dry	Birds	Olive-sided Flycatcher	Contopus cooperi	No	Yes	G4	CA, NV					No
Dry	Birds	Black Swift	Cypseloides niger	No	Yes	G4	CA, ID, UT			13	SW, CA	No
Dry	Birds	Dusky Grouse	Dendragapus obscurus	No	Yes	G5	NV					No
Dry	Birds	Grace's Warbler	Dendroica graciae	No	Yes	G5	NV					No
Dry	Birds	Black-throated Gray Warbler	Dendroica nigrescens	No	Yes	G5	UT					No
Dry	Birds	Hermit Warbler	Dendroica occidentalis	No	Yes	G4	CA, NV					No
Dry	Birds	A Yellow Warbler	Dendroica petechia brewsteri	No	No	T3	CA			5		No
Dry	Birds	Bobolink	Dolichonyx oryzivorus	No	Yes	G5	NV, UT	UT	PS	39	SW	No
Dry	Birds	Gray Catbird	Dumetella carolinensis	No	Yes	G5				6	SW	No
Dry	Birds	Snowy Egret	Egretta thula	No	Yes	G5	CA, ID, NV		PS	1	SW	No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Birds	Willow Flycatcher	Empidonax traillii	Yes	Yes	G5	CA			15	SW, CA	Yes
Dry	Birds	A Willow Flycatcher	Empidonax traillii adastus	No	No	T5	NV					No
Dry	Birds	Southwestern Willow Flycatcher	Empidonax traillii extimus	Yes	Yes	T1	CA, NV, UT	CA	PS	21		No
Dry	Birds	Gray Flycatcher	Empidonax wrightii	No	Yes	G5				10	SW, CA	No
Dry	Birds	Merlin	Falco columbarius	No	Yes	G5	CA, ID			1	SW	No
Dry	Birds	Prairie Falcon	Falco mexicanus	No	Yes	G5	CA			36	SW, CA	No
Dry	Birds	Peregrine Falcon	Falco peregrinus	No	Yes	G4	ID, NV, UT		PS	163	SW, CA	No
Dry	Birds	American Peregrine Falcon	Falco peregrinus anatum	No	Yes	T4	CA			2		No
Dry	Birds	Common Moorhen	Gallinula chloropus	No	Yes	G5				4	SW	No
Dry	Birds	Greater Roadrunner	Geococcyx californianus	No	Yes	G5				2	SW, CA	No
Dry	Birds	Common Yellowthroat	Geothlypis trichas	No	Yes	G5				40	SW, CA	No
Dry	Birds	Pinyon Jay	Gymnorhinus cyanocephalus	No	Yes	G5	ID, NV		PS	11	SW, CA	No
Dry	Birds	Bald Eagle	Haliaeetus leucocephalus	No	Yes	G5	CA, ID, NV, UT	CA, UT	PS	888	SW, CA	No
Dry	Birds	Yellow-breasted Chat	Icteria virens	No	Yes	G5	CA			7	SW, CA	No
Dry	Birds	Hooded Oriole	Icterus cucullatus	No	Yes	G5				2	SW, CA	No
Dry	Birds	Scott's Oriole	Icterus parisorum	No	Yes	G5	NV		PS	1	SW, CA	No
Dry	Birds	Loggerhead Shrike	Lanius ludovicianus	No	Yes	G4	CA, NV		IL		SW, CA	No
Dry	Birds	California Black Rail	Laterallus jamaicensis coturniculus	No	Yes	T1	CA	CA				No
Dry	Birds	Black Rosy-finch	Leucosticte atrata	No	Yes	G4	ID, NV, UT		HV		SW	No
Dry	Birds	gray-crowned rosy-finch	Leucosticte tephrocotis	No	Yes	G5	NV		HV			No
Dry	Birds	Hooded Merganser	Lophodytes cucullatus	No	Yes	G5	ID					No
Dry	Birds	Red Crossbill	Loxia curvirostra	No	Yes	G5	ID					No
Dry	Birds	White-winged Crossbill	Loxia leucoptera	No	Yes	G5	ID					No
Dry	Birds	Lewis's Woodpecker	Melanerpes lewis	No	Yes	G4	CA, ID, NV, UT	UT	PS	34	SW, CA	No
Dry	Birds	Lincoln's Sparrow	Melospiza lincolni	No	Yes	G5						No
Dry	Birds	Brown-crested Flycatcher	Myiarchus tyrannulus	No	Yes	G5	CA			2	SW, CA	No
Dry	Birds	Painted Redstart	Myioborus pictus	No	Yes	G5				1	SW	No
Dry	Birds	Clark's Nutcracker	Nucifraga columbiana	No	Yes	G5						No
Dry	Birds	Long-billed Curlew	Numenius americanus	No	Yes	G5	CA, ID, NV, UT	UT	PS	216	SW	No
Dry	Birds	MacGillivray's Warbler	Oporornis tolmiei	No	Yes	G5						No
Dry	Birds	Mountain Quail	Oreortyx pictus	No	Yes	G5	ID, NV		PS	9	SW, CA	No
Dry	Birds	Sage Thrasher	Oreoscoptes montanus	No	Yes	G5	UT			1	SW, CA	No
Dry	Birds	Orange-crowned Warbler	Oreothlypis celata	No	Yes	G5						No
Dry	Birds	Flammulated Owl	Otus flammeolus	No	Yes	G4	CA, ID			19	SW, CA	No
Dry	Birds	Osprey	Pandion haliaetus	No	Yes	G5	CA, UT			48	SW, CA	No
Dry	Birds	Savannah Sparrow	Passerculus sandwichensis	No	Yes	G5						No
Dry	Birds	Blue Grosbeak	Passerina caerulea	No	Yes	G5	ID			39	SW, CA	No
Dry	Birds	Indigo Bunting	Passerina cyanea	No	Yes	G5						No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Birds	Band-tailed Pigeon	Patagioenas fasciata	No	Yes	G4	UT			50	SW, CA	No
Dry	Birds	Gray Jay	Perisoreus canadensis	No	Yes	G5						No
Dry	Birds	Phainopepla	Phainopepla nitens	No	Yes	G5	NV		PS	7	SW, CA	No
Dry	Birds	Black-billed Magpie	Pica hudsonia	No	Yes	G5						No
Dry	Birds	White-headed Woodpecker	Picoides albolarvatus	No	Yes	G4	CA, ID, NV					No
Dry	Birds	American Three-toed Woodpecker	Picoides dorsalis	No	Yes	G5	ID, UT	UT		16		No
Dry	Birds	Nuttall's Woodpecker	Picoides nuttallii	No	No	G5	CA					No
Dry	Birds	Downy Woodpecker	Picoides pubescens	No	Yes	G5						No
Dry	Birds	Ladder-backed Woodpecker	Picoides scalaris	No	Yes	G5				2	SW, CA	No
Dry	Birds	Pine Grosbeak	Pinicola enucleator	No	Yes	G5						No
Dry	Birds	Abert's Towhee	Pipilo aberti	No	Yes	G3	CA, NV, UT		IL	9	SW, CA	No
Dry	Birds	Green-tailed Towhee	Pipilo chlorurus	No	Yes	G5						No
Dry	Birds	Summer Tanager	Piranga rubra	No	Yes	G5	CA			3	SW, CA	No
Dry	Birds	Black-capped Chickadee	Poecile atricapillus	No	Yes	G5	CA					No
Dry	Birds	Black-tailed Gnatcatcher	Poliophtila melanura	No	Yes	G5	CA			2	SW, CA	No
Dry	Birds	Purple Martin	Progne subis	No	Yes	G5	CA			10	SW, CA	No
Dry	Birds	Vermilion Flycatcher	Pyrocephalus rubinus	No	Yes	G5	CA			6	SW	No
Dry	Birds	Common Grackle	Quiscalus quiscula	No	Yes	G5				2	SW	No
Dry	Birds	Ruby-crowned Kinglet	Regulus calendula	No	Yes	G5						No
Dry	Birds	Golden-crowned Kinglet	Regulus satrapa	No	Yes	G5						No
Dry	Birds	Bank Swallow	Riparia riparia	No	Yes	G5	CA	CA		9	SW, CA	No
Dry	Birds	Black Phoebe	Sayornis nigricans	No	Yes	G5	NV		IL	4	SW, CA	No
Dry	Birds	Broad-tailed Hummingbird	Selasphorus platycercus	No	Yes	G5	UT			2	SW, CA	No
Dry	Birds	Rufous Hummingbird	Selasphorus rufus	No	Yes	G5	CA, NV					No
Dry	Birds	American Redstart	Setophaga ruticilla	No	Yes	G5				10	SW	No
Dry	Birds	Pygmy Nuthatch	Sitta pygmaea	No	Yes	G5	ID					No
Dry	Birds	Red-naped Sapsucker	Sphyrapicus nuchalis	No	Yes	G5						No
Dry	Birds	Red-breasted Sapsucker	Sphyrapicus ruber	No	Yes	G5	CA, NV					No
Dry	Birds	Williamson's Sapsucker	Sphyrapicus thyroideus	No	Yes	G5	UT			5	SW, CA	No
Dry	Birds	Lawrence's Goldfinch	Spinus lawrencei	No	Yes	G3	CA					No
Dry	Birds	Lesser Goldfinch	Spinus psaltria	No	Yes	G5	ID					No
Dry	Birds	Brewer's Sparrow	Spizella breweri	No	Yes	G5	CA, ID, NV, UT		MV	9	CA	No
Dry	Birds	Chipping Sparrow	Spizella passerina	No	Yes	G5	CA					No
Dry	Birds	Calliope Hummingbird	Stellula calliope	No	Yes	G5					SW, CA	No
Dry	Birds	Great Gray Owl	Strix nebulosa	No	Yes	G5	CA			10	CA	No
Dry	Birds	Spotted Owl	Strix occidentalis	No	Yes	G3				4	SW, CA	Yes
Dry	Birds	California Spotted Owl	Strix occidentalis occidentalis	No	Yes	T3	CA, NV	CA	MV	3		No
Dry	Birds	Tree Swallow	Tachycineta bicolor	No	Yes	G5						No
Dry	Birds	Bendire's Thrasher	Toxostoma bendirei	No	Yes	G4	CA, NV, UT	CA	PS	2	SW, CA	Yes

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Birds	Crissal Thrasher	Toxostoma crissale	No	Yes	G5	CA, NV, UT		IL	3	SW, CA	No
Dry	Birds	Le Conte's Thrasher	Toxostoma lecontei	No	Yes	G4	CA, NV	CA	PS	2	SW, CA	No
Dry	Birds	Winter Wren	Troglodytes troglodytes	No	Yes	G5						No
Dry	Birds	American Robin	Turdus migratorius	No	Yes	G5				1	SW, CA	No
Dry	Birds	Sharp-tailed Grouse	Tympanuchus phasianellus	No	Yes	G4	ID, UT	UT		127		No
Dry	Birds	Columbian Sharp-tailed Grouse	Tympanuchus phasianellus columbianus	No	Yes	T3	CA, NV		MV	74	SW	No
Dry	Birds	Eastern Kingbird	Tyrannus tyrannus	No	Yes	G5				18	SW	No
Dry	Birds	Cassin's Kingbird	Tyrannus vociferans	No	Yes	G5				2	SW, CA	No
Dry	Birds	Virginia's Warbler	Vermivora virginiae	No	Yes	G5	CA, ID, NV, UT		PS	1	SW, CA	No
Dry	Birds	Bell's Vireo	Vireo bellii	Yes	Yes	G5	UT			1	SW, CA	No
Dry	Birds	Least Bell's Vireo	Vireo bellii pusillus	Yes	Yes	T2	CA	CA		3		Yes
Dry	Birds	Gray Vireo	Vireo vicinior	No	Yes	G4	CA, NV, UT	CA	PS	1	SW, CA	No
Dry	Birds	Yellow-headed Blackbird	Xanthocephalus xanthocephalus	No	Yes	G5	CA			1	SW, CA	No
Dry	Birds	White-winged Dove	Zenaida asiatica	No	Yes	G5				1	SW, CA	No
Dry	Birds	White-crowned Sparrow	Zonotrichia leucophrys	No	Yes	G5						No
Dry	Butterflies & Skippers	Desert Green Hairstreak	Callophrys comstocki	No	No	G2				1		No
Dry	Butterflies & Skippers	Mcneill's Saltbush Sootywing	Hesperopsis graciellae	No	No	G2		AZ		1		No
Dry	Butterflies & Skippers	San Emigdio Blue	Plebulina emigdionis	No	No	G2				1		No
Dry	Butterflies & Skippers	Carson Wandering Skipper	Pseudocopaeodes eunus obscurus	Yes	No	T1				22		No
Dry	Butterflies & Skippers	Nokomis Fritillary	Speyeria nokomis	No	No	G3				3		No
Dry	Mammals	Moose	Alces americanus	No	Yes	G5				1	SW	No
Dry	Mammals	Pallid Bat	Antrozous pallidus	No	Yes	G5	CA	CA		46	SW	No
Dry	Mammals	Sewellel	Aplodontia rufa	No	Yes	G5				1	SW, CA	No
Dry	Mammals	Sierra Nevada Mountain Beaver	Aplodontia rufa californica	No	Yes	T3	CA, NV		HV	4		No
Dry	Mammals	Ringtail	Bassariscus astutus	No	No	G5	NV		PS	14	SW, CA	No
Dry	Mammals	Pygmy Rabbit	Brachylagus idahoensis	No	Yes	G4	CA, ID, NV, UT	CA, UT	EV	236	SW, CA	No
Dry	Mammals	Gray Wolf	Canis lupus	Yes	Yes	G4	ID, UT			2	SW	No
Dry	Mammals	Desert Pocket Mouse	Chaetodipus penicillatus	No	No	G5	NV		MV	2	SW, CA	No
Dry	Mammals	Townsend's Big-eared Bat	Corynorhinus townsendii	No	Yes	G4	CA, ID, NV, UT	CA, UT	PS	211	SW	No
Dry	Mammals	White-tailed Prairie Dog	Cynomys leucurus	No	Yes	G4	UT	UT				No
Dry	Mammals	Utah Prairie Dog	Cynomys parvidens	Yes	Yes	G1	UT			502	SW	No
Dry	Mammals	Desert Kangaroo Rat	Dipodomys deserti	No	No	G5	NV, UT		PS	8	SW, CA	No
Dry	Mammals	Merriam's Kangaroo Rat	Dipodomys merriami	Yes	No	G5				5	SW, CA	No
Dry	Mammals	Argus Mountains Kangaroo Rat	Dipodomys panamintinus argusensis	No	No	T2	CA			1		No
Dry	Mammals	Panamint Kangaroo Rat	Dipodomys panamintinus panamintinus	No	No	T3	CA			1		No
Dry	Mammals	Spotted Bat	Euderma maculatum	No	Yes	G4	CA, ID, NV,	CA, UT	PS	41	SW	No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
							UT					
Dry	Mammals	California Bonneted Bat	Eumops perotis californicus	No	Yes	T4		CA		6		No
Dry	Mammals	Northern Flying Squirrel	Glaucomys sabrinus	No	Yes	G5	NV, UT		PS	35	SW, CA	No
Dry	Mammals	Wolverine	Gulo gulo	No	Yes	G4	CA, ID, UT			37	SW, CA	No
Dry	Mammals	North American Wolverine	Gulo gulo luscus	No	Yes	T4				4		No
Dry	Mammals	Allen's Big-eared Bat	Idionycteris phyllotis	No	Yes	G3	NV, UT	AZ, UT	PS	2	SW	No
Dry	Mammals	Silver-haired Bat	Lasionycteris noctivagans	No	No	G5	CA			29	SW	No
Dry	Mammals	Western Red Bat	Lasiurus blossevillii	No	Yes	G5	CA, NV, UT	UT	PS	3	SW	No
Dry	Mammals	Hoary Bat	Lasiurus cinereus	No	No	G5	CA, NV		IL	19	SW	No
Dry	Mammals	Sagebrush Vole	Lemmiscus curtatus	No	No	G5	NV		HV		SW, CA	No
Dry	Mammals	Sierra Nevada Snowshoe Hare	Lepus americanus tahoensis	No	Yes	T3	CA			2		No
Dry	Mammals	White-tailed Jackrabbit	Lepus townsendii	No	Yes	G5				18	SW, CA	No
Dry	Mammals	Canadian Lynx	Lynx canadensis	Yes	Yes	G5	ID, UT			2	SW	No
Dry	Mammals	American Marten	Martes americana	No	Yes	G5	CA, NV, UT		PS	6	SW, CA	No
Dry	Mammals	Sierra Marten	Martes americana sierrae	No	No	T3	CA			20		No
Dry	Mammals	Fisher	Martes pennanti	No	Yes	G5	CA, ID	CA				No
Dry	Mammals	Fisher - West Coast Distinct Population Segment	Martes pennanti pop. 1	Yes	No	T2				7		No
Dry	Mammals	Dark Kangaroo Mouse	Microdipodops megacephalus	No	Yes	G4	NV, UT	UT		27	SW, CA	No
Dry	Mammals	Desert Valley Kangaroo Mouse	Microdipodops megacephalus albiventer	No	Yes	T2	NV		MV	4		No
Dry	Mammals	Fletcher Kangaroo Mouse	Microdipodops megacephalus nasutus	No	Yes	T2			PS	2		No
Dry	Mammals	Pale Kangaroo Mouse	Microdipodops pallidus	No	Yes	G3	NV		PS		SW, CA	No
Dry	Mammals	Owens Valley Vole	Microtus californicus vallicola	No	No	T1	CA	CA		13		No
Dry	Mammals	Pahranagat Valley Vole	Microtus montanus fucosus	No	Yes	T2	NV		PS	6		No
Dry	Mammals	Western Small-footed Myotis	Myotis ciliolabrum	No	No	G5	CA, NV	AZ, CA	PS	67	SW	No
Dry	Mammals	Long-eared Myotis	Myotis evotis	No	No	G5	CA	AZ, CA	IL	53	SW	No
Dry	Mammals	Little Brown Myotis	Myotis lucifugus	No	No	G5	CA, NV	AZ	IL	5	SW	No
Dry	Mammals	Fringed Myotis	Myotis thysanodes	No	Yes	G4	CA, ID, NV, UT	AZ, CA, UT	IL	28	SW	No
Dry	Mammals	Long-legged Myotis	Myotis volans	No	No	G5	CA	AZ		74	SW	No
Dry	Mammals	Yuma Myotis	Myotis yumanensis	No	No	G5	CA, UT	CA		25	SW	No
Dry	Mammals	Yellow-pine Chipmunk	Neotamias amoenus	No	Yes	G5				1	SW, CA	No
Dry	Mammals	Yellow-pine Chipmunk	Neotamias amoenus celeris	No	No	T2	NV		MV	1		No
Dry	Mammals	Cliff Chipmunk	Neotamias dorsalis	No	Yes	G5	ID			1	SW	No
Dry	Mammals	Least Chipmunk	Neotamias minimus	No	Yes	G5						No
Dry	Mammals	Shadow Chipmunk	Neotamias senex	No	No	G5	NV					No
Dry	Mammals	Uinta Chipmunk	Neotamias umbrinus	No	Yes	G5						No
Dry	Mammals	Crawford's Gray Shrew	Notiosorex crawfordi	No	No	G5	UT			2	SW, CA	No
Dry	Mammals	Big Free-tailed Bat	Nyctinomops macrotis	No	Yes	G5	CA, NV, UT	AZ, UT	PS	8	SW	No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Dry	Mammals	American Pika	Ochotona princeps	No	Yes	G5	NV, UT		HV	29	SW, CA	No
Dry	Mammals	White Mountains Pika	Ochotona princeps sheltoni	No	No	T1	CA			11		No
Dry	Mammals	mule deer	Odocoileus hemionus	No	Yes	G5	NV, UT	CBR, MBR	PS		SW, CA	Yes
Dry	Mammals	Common Muskrat	Ondatra zibethicus	No	Yes	G5						No
Dry	Mammals	Bighorn Sheep	Ovis canadensis	Yes	Yes	G4	ID, UT					Yes
Dry	Mammals	Desert Bighorn Sheep	Ovis canadensis nelsoni	No	Yes	T4	CA, NV	CA	PS	5		Yes
Dry	Mammals	Sierra Nevada Bighorn Sheep	Ovis canadensis sierrae	Yes	Yes	T1	CA, NV	CA		4		No
Dry	Mammals	Western Pipistrelle	Parastrellus hesperus	No	Yes	G5				22	SW	No
Dry	Mammals	Brush Deer mouse	Peromyscus boylii	No	No	G5	NV					No
Dry	Mammals	Piñon Deer mouse	Peromyscus truei	No	No	G5	ID					No
Dry	Mammals	Broad-footed Mole	Scapanus latimanus	No	No	G5	NV					No
Dry	Mammals	Western Gray Squirrel	Sciurus griseus griseus	No	Yes	T5				2		No
Dry	Mammals	Mt. Lyell Shrew	Sorex lyelli	No	No	G2	CA			9	CA	No
Dry	Mammals	Merriam's Shrew	Sorex merriami	No	No	G5	ID, UT			1	SW, CA	No
Dry	Mammals	Merriam's Shrew	Sorex merriami leucogenys	No	No	T5	NV		PS	3		No
Dry	Mammals	montane shrew	Sorex monticolus	No	No	G5	NV		MV			No
Dry	Mammals	Dwarf Shrew	Sorex nanus	No	No	G4	UT					No
Dry	Mammals	water shrew	Sorex palustris	No	Yes	G5	NV		MV			No
Dry	Mammals	Preble's Shrew	Sorex preblei	No	Yes	G4	NV, UT	UT	PS	4	SW	No
Dry	Mammals	Inyo Shrew	Sorex tenellus	No	No	G3	NV		PS	3	SW, CA	No
Dry	Mammals	Trowbridge's Shrew	Sorex trowbridgii	No	No	G5	NV		PS	2	SW, CA	No
Dry	Mammals	Vagrant Shrew	Sorex vagrans	No	No	G5	NV					No
Dry	Mammals	Wyoming Ground Squirrel	Spermophilus elegans	No	No	G5	UT					No
Dry	Mammals	Wyoming Ground Squirrel	Spermophilus elegans nevadensis	No	No	T4	NV					No
Dry	Mammals	Mohave Ground Squirrel	Spermophilus mohavensis	No	Yes	G2	CA	CA		5	CA	No
Dry	Mammals	Piute Ground Squirrel	Spermophilus mollis	No	No	G5	ID					No
Dry	Mammals	Rock Squirrel	Spermophilus variegatus	No	Yes	G5	ID			18	SW, CA	No
Dry	Mammals	Brazilian Free-tailed Bat	Tadarida brasiliensis	No	Yes	G5				43	SW	No
Dry	Mammals	American Badger	Taxidea taxus	No	No	G5	CA			14	SW, CA	No
Dry	Mammals	Fish Spring Pocket Gopher	Thomomys bottae abstrusus	No	No	TH	NV		MV	1		No
Dry	Mammals	San Antonio Pocket Gopher	Thomomys bottae curtatus	No	No	TH	NV		MV	1		No
Dry	Mammals	Mountain Pocket Gopher	Thomomys monticola	No	No	G5	NV		PS	1	SW, CA	No
Dry	Mammals	Townsend's Pocket Gopher	Thomomys townsendii	No	No	G4	ID					No
Dry	Mammals	American Black Bear	Ursus americanus	No	Yes	G5				2	SW, CA	No
Dry	Mammals	Brown Bear	Ursus arctos	Yes	Yes	G4	ID, UT			7	SW	No
Dry	Mammals	Kit Fox	Vulpes macrotis	Yes	Yes	G4	NV, UT	UT	PS	308	SW, CA	No
Dry	Mammals	Red Fox	Vulpes vulpes	No	Yes	G5				4	SW	No
Dry	Mammals	Sierra Nevada Red Fox	Vulpes vulpes necator	No	Yes	T2	CA, NV		PS	7		No
Dry	Mammals	Western Jumping Mouse	Zapus princeps	No	No	G5	NV		PS	2	SW, CA	No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Millipedes & Centipedes	A Millipede	Polydesmus cavicola	No	No	G1				1		No
Dry	Other Beetles	Crescent-dune Aegialian Scarab Beetle	Aegialia crescenta	No	No	G1				1		No
Dry	Other Beetles	Hardy's Aegialian Scarab Beetle	Aegialia hardyi	No	No	G1				2		No
Dry	Other Beetles	Utah Chaetarthrian Water Scavenger Beetle	Chaetarthria utahensis	No	No	G1				1		No
Dry	Other Beetles	A Beetle	Coenonycha pygmaea	No	No	G1				2		No
Dry	Other Beetles	Leech's Skyline Diving Beetle	Hydroporus leechi	No	No	G1				1		No
Dry	Other Beetles	Travertine Band-thigh Diving Beetle	Hygrotus fontinalis	No	No	G1				4		No
Dry	Other Beetles	Nelson's Miloderes Weevil	Miloderes nelsoni	No	No	G2				1		No
Dry	Other Beetles	Saline Valley Snow-front Scarab Beetle	Polyphylla anteronivea	No	No	G1				1		No
Dry	Other Beetles	Spotted Warner Valley Dunes Scarab Beetle	Polyphylla avittata	No	No	G2				2		No
Dry	Other Beetles	Crescent Dune Serican Scarab Beetle	Serica ammomenisco	No	No	G1				1		No
Dry	Other Beetles	Humboldt Serican Beetle	Serica humboldti	No	No	G1				1		No
Dry	Other Beetles	Sand Mountain Serican Scarab Beetle	Serica psammobunus	No	No	G1				2		No
Dry	Other Beetles		Stenelmis lariversi	No	No	G1				1		No
Dry	Other Beetles	Moapa Warm Springs Riffle Beetle	Stenelmis moapa	No	No	G1				1		No
Dry	Other Insects	Amargosa Naucorid Bug	Pelocoris shoshone	No	No	G2				1		No
Dry	Reptiles	Glossy Snake	Arizona elegans	No	No	G5	UT			13	SW, CA	No
Dry	Reptiles	Plateau Striped Whiptail	Aspidoscelis velox	No	No	G5	UT			7	SW	No
Dry	Reptiles	Zebra-tailed Lizard	Callisaurus draconoides	No	Yes	G5	UT	UT		60	SW, CA	No
Dry	Reptiles	Northern Rubber Boa	Charina bottae	No	No	G5	UT			64	SW, CA	No
Dry	Reptiles	Western Banded Gecko	Coleonyx variegatus	No	Yes	G5	NV, UT	UT		29	SW, CA	No
Dry	Reptiles	Sidewinder	Crotalus cerastes	No	Yes	G5	UT	UT		17	SW, CA	No
Dry	Reptiles	Speckled Rattlesnake	Crotalus mitchellii	No	Yes	G5	UT	UT		2	SW, CA	No
Dry	Reptiles	Mohave Rattlesnake	Crotalus scutulatus	No	Yes	G5	UT	UT		12	SW, CA	No
Dry	Reptiles	Great Basin Collared Lizard	Crotaphytus bicinctores	No	Yes	G5	ID, NV					No
Dry	Reptiles	Ring-necked Snake	Diadophis punctatus	No	Yes	G5	ID, UT			29	SW, CA	No
Dry	Reptiles	Desert Iguana	Dipsosaurus dorsalis	No	Yes	G5	NV, UT	UT	MV	1	SW, CA	No
Dry	Reptiles	Northern Alligator Lizard	Elgaria coerulea	No	No	G5	ID					No
Dry	Reptiles	Sierra Alligator Lizard	Elgaria coerulea palmeri	No	Yes	T4	NV		PS	2		No
Dry	Reptiles	Shasta alligator lizard	Elgaria coerulea shastensis	No	No	T4			MV			No
Dry	Reptiles	Panamint Alligator Lizard	Elgaria panamintina	No	No	G2	CA	CA	PS	8	CA	No
Dry	Reptiles	Gilbert's Skink	Eumeces gilberti	No	No	G5	NV					No
Dry	Reptiles	Long-nosed Leopard Lizard	Gambelia wislizenii	No	No	G5	NV, UT		PS	7	SW, CA	No
Dry	Reptiles	Gila Monster	Heloderma suspectum	No	Yes	G4	UT	CA, UT		40	SW, CA	No
Dry	Reptiles	Banded Gila Monster	Heloderma suspectum cinctum	No	Yes	T4	CA, NV	AZ	MV	6		No
Dry	Reptiles	Nightsnake	Hypsiglena torquata	No	No	G5	UT					No
Dry	Reptiles	Common Kingsnake	Lampropeltis getula	No	No	G5	UT			12	SW, CA	No
Dry	Reptiles	Sonoran Mountain Kingsnake	Lampropeltis pyromelana	No	Yes	G4	NV, UT		HV	37	SW	No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Dry	Reptiles	Milksnake	Lampropeltis triangulum	No	No	G5	UT			54	SW	No
Dry	Reptiles	Western Threadsnake	Leptotyphlops humilis	No	Yes	G5	UT	UT		5	SW, CA	No
Dry	Reptiles	Coachwhip	Masticophis flagellum	No	No	G5	UT			21	SW, CA	No
Dry	Reptiles	Smooth Greensnake	Opheodrys vernalis	No	Yes	G5	UT	UT		9	SW	No
Dry	Reptiles	Pygmy Horned Lizard	Phrynosoma douglasii	No	No	G5	NV					No
Dry	Reptiles	Short-horned Lizard	Phrynosoma hernandesi	No	No	G5	NV		PS		CA	No
Dry	Reptiles	Desert Horned Lizard	Phrynosoma platyrhinus	No	No	G5	NV		PS	2	SW, CA	No
Dry	Reptiles	Western Skink	Plestiodon skiltonianus	No	No	G5	UT					No
Dry	Reptiles	Coronado Skink	Plestiodon skiltonianus interparietalis	No	No	T5	CA	CA				No
Dry	Reptiles	Long-nosed Snake	Rhinocheilus lecontei	No	Yes	G5	ID, UT			16	SW, CA	No
Dry	Reptiles	Western Patch-nosed Snake	Salvadora hexalepis	No	No	G5	UT			7	SW, CA	No
Dry	Reptiles	Common Chuckwalla	Sauromalus ater	No	Yes	G5	CA, NV, UT	UT		54	SW, CA	No
Dry	Reptiles	Northern Sagebrush Lizard	Sceloporus graciosus graciosus	No	No	T5	CA	AZ, CA		2		No
Dry	Reptiles	Groundsnake	Sonora semiannulata	No	Yes	G5	ID, UT			8	SW, CA	No
Dry	Reptiles	Smith's Black-headed Snake	Tantilla hobartsmithi	No	No	G5	UT			9	SW, CA	No
Dry	Reptiles	Common Gartersnake	Thamnophis sirtalis	No	No	G5	UT			45	SW, CA	No
Dry	Reptiles	Western Lyresnake	Trimorphodon biscutatus	No	No	G5	UT					No
Dry	Reptiles	Sonoran Lyresnake	Trimorphodon lambda	No	No	G5	NV		FOR SPECIES/SUB	3		No
Dry	Reptiles	long-tailed brush lizard	Urosaurus graciosus	No	No	G5	NV		MV			No
Dry	Reptiles	Desert Night Lizard	Xantusia vigilis	No	Yes	G5	UT	UT	FOR SSP	10	SW, CA	No
Dry	Reptiles	desert night lizard	Xantusia vigilis vigilis	No	No	T5	NV		MV			No
Dry	Spiders & other Chelicerates	A Cave Obligate Harvestman	Hesperonemastoma packardi	No	No	G1				1		No
Dry	Terrestrial Snails	Sierra Ambersnail	Catinella stretchiana	No	No	G3				1		No
Dry	Terrestrial Snails	Cross Snaggletooth	Gastrocopta quadridens	No	No	G2				1		No
Dry	Terrestrial Snails	Southern Tightcoil	Ogaridiscus subrupicola	No	Yes	G1				1		No
Dry	Terrestrial Snails	Eureka Mountainsnail	Oreohelix eurekensis	No	Yes	G1				3		No
Dry	Terrestrial Snails	Lyrate Mountainsnail	Oreohelix haydeni	No	Yes	G2				19		No
Dry	Terrestrial Snails	Whitepine Mountainsnail	Oreohelix hemphilli	No	No	G2						No
Dry	Terrestrial Snails	Mill Creek Mountainsnail	Oreohelix howardi	No	No	G1				3		No
Dry	Terrestrial Snails	Goshute Mountainsnail	Oreohelix loisae	No	No	G2				3		No
Dry	Terrestrial Snails	Schell Creek Mountainsnail	Oreohelix nevadensis	No	No	G1				5		No
Dry	Terrestrial Snails	Brian Head Mountainsnail	Oreohelix parawanensis	No	Yes	G1				4		No
Dry	Terrestrial Snails	Deseret Mountainsnail	Oreohelix peripherica	No	Yes	G2				11		No
Dry	Terrestrial Snails	Ogden Rocky Mountainsnail	Oreohelix peripherica wasatchensis	No	Yes	T1				1		No
Dry	Terrestrial Snails	Santa Rita Ambersnail	Succinea grosvenori	No	No	G5		AZ		4		No
Dry	Terrestrial Snails	Rustic Ambersnail	Succinea rusticana	No	No	G2		AZ		3		No
Dry	Tiger Beetles	Mojave Giant Tiger Beetle	Amblycheila schwarzi	No	No	G3				1		No
Dry	Tiger Beetles	Maricopa Tiger Beetle	Cicindela oregona maricopa	No	No	T3		AZ		12		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Tiger Beetles	Riparian Tiger Beetle	Cicindela praetextata	No	No	G2				1		No
Dry	Turtles	Desert Tortoise	Gopherus agassizii	Yes	Yes	G4	CA, NV, UT	CA	PS	632	SW, CA	Yes
Dry	Conifers & relatives	Washoe Pine	Pinus washoensis	No	Yes	G3		NV		5		No
Dry	Ferns & relatives	Upward-lobed Moonwort	Botrychium ascendens	No	No	G2				4		No
Dry	Ferns & relatives	Crenulate Moonwort	Botrychium crenulatum	No	No	G3				15		No
Dry	Ferns & relatives	Narrowleaf Grapefern	Botrychium lineare	No	No	G2				1		No
Dry	Ferns & relatives	Utah Spike-moss	Selaginella utahensis	No	No	G2				5		No
Dry	Flowering Plants	Passey's Onion	Allium passeyi	No	No	G1				12		No
Dry	Flowering Plants	Wheeler's Angelica	Angelica wheeleri	No	No	G2				11		No
Dry	Flowering Plants	Meadow Pussytoes	Antennaria arcuata	No	No	G2		NV		4		No
Dry	Flowering Plants	Beckwith's Rockcress	Arabis beckwithii	No	No	G2				3		No
Dry	Flowering Plants	Bodie Hills Rockcress	Arabis bodiensis	No	No	G2		CA, NV		29		No
Dry	Flowering Plants	Unequal Rockcress	Arabis dispar	No	No	G3				20		No
Dry	Flowering Plants	Grouse Creek Rockcress	Arabis falcatoria	No	No	G1				10		No
Dry	Flowering Plants	Elko Rockcress	Arabis falcifruca	No	No	G1		NV		1		No
Dry	Flowering Plants	Wasatch Range Rockcress	Arabis lasiocarpa	No	No	G3				19		No
Dry	Flowering Plants	Ophir Rockcress	Arabis ophira	No	No	G1				15		No
Dry	Flowering Plants	Pinzl's Rockcress	Arabis pinzliae	No	No	G2				11		No
Dry	Flowering Plants	Darwin Rock Cress	Arabis pulchra var. munciensis	No	No	T4		CA		4		No
Dry	Flowering Plants	Shockley's Rockcress	Arabis shockleyi	No	No	G3				30		No
Dry	Flowering Plants	Tiehm's Rockcress	Arabis tiehmii	No	No	G2				14		No
Dry	Flowering Plants	Dwarf Bear-poppy	Arctomecon humilis	Yes	No	G1				170		No
Dry	Flowering Plants	White Bear-poppy	Arctomecon merriamii	No	No	G3				9		No
Dry	Flowering Plants	Packard's Wormwood	Artemisia packardiae	No	No	G3				3		No
Dry	Flowering Plants	Eastwood's Milkweed	Asclepias eastwoodiana	No	No	G2		NV		32		No
Dry	Flowering Plants	Ackerman's Milkvetch	Astragalus ackermanii	No	No	G2				3		No
Dry	Flowering Plants	Purple Milkvetch	Astragalus agrestis	No	No	G5		CA		1		No
Dry	Flowering Plants		Astragalus ampullarioides	Yes	No	G1				5		No
Dry	Flowering Plants	Silverleaf Milkvetch	Astragalus argophyllus var. argophyllus	No	No	T4		CA		12		No
Dry	Flowering Plants		Astragalus avonensis	No	No	G1				1		No
Dry	Flowering Plants	Beatley's Milkvetch	Astragalus beatleyae	No	No	G2				40		No
Dry	Flowering Plants	Callaway Milkvetch	Astragalus callithrix	No	No	G3				20		No
Dry	Flowering Plants	Ground-crescent Milkvetch	Astragalus chamaemeniscus	No	No	G2				2		No
Dry	Flowering Plants	Cima Milkvetch	Astragalus cimae var. cimae	No	No	T2		NV		3		No
Dry	Flowering Plants	Margaret's Rushy Milkvetch	Astragalus convallarius var. margaretae	No	No	T2		NV		11		No
Dry	Flowering Plants	Mesic Milkvetch	Astragalus diversifolius	No	No	G2				3		No
Dry	Flowering Plants	Pagumpa Milkvetch	Astragalus ensiformis var. gracilior	No	No	T1		NV		1		No
Dry	Flowering Plants	Peck Station Milkvetch	Astragalus eurylobus	No	No	G2		NV		6		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Black Milkvetch	<i>Astragalus funereus</i>	No	No	G2		CA, NV		8		No
Dry	Flowering Plants	Gilman's Milkvetch	<i>Astragalus gilmanii</i>	No	No	G2				3		No
Dry	Flowering Plants	Holmgren's Milkvetch	<i>Astragalus holmgreniorum</i>	Yes	Yes	G1				4		No
Dry	Flowering Plants	Horn's Milkvetch	<i>Astragalus hornii</i> var. <i>hornii</i>	No	No	T2		CA		1		No
Dry	Flowering Plants	Inyo Milkvetch	<i>Astragalus inyoensis</i>	No	No	G3				1		No
Dry	Flowering Plants	Long Valley Milkvetch	<i>Astragalus johannis-howellii</i>	No	Yes	G2		CA, NV		28		No
Dry	Flowering Plants	Lemmon's Milkvetch	<i>Astragalus lemmonii</i>	No	No	G2		CA		8		No
Dry	Flowering Plants	Lens-pod Milkvetch	<i>Astragalus lentiformis</i>	No	No	G2		CA		23		No
Dry	Flowering Plants	Fish Slough Milkvetch	<i>Astragalus lentiginosus</i> var. <i>piscinensis</i>	Yes	No	T1		CA		8		No
Dry	Flowering Plants	Sodaville Milkvetch	<i>Astragalus lentiginosus</i> var. <i>sesquimetralis</i>	No	Yes	T1		NV		3		No
Dry	Flowering Plants	Mottled Milkvetch	<i>Astragalus lentiginosus</i> var. <i>stramineus</i>	No	No	T2		NV		4		No
Dry	Flowering Plants	Heliotrope Milkvetch	<i>Astragalus limnocharis</i> var. <i>montii</i>	Yes	No	T1				11		No
Dry	Flowering Plants	Glenwood Milkvetch	<i>Astragalus loanus</i>	No	No	G1				7		No
Dry	Flowering Plants	Mono Milkvetch	<i>Astragalus monoensis</i>	No	Yes	G2		CA		37		No
Dry	Flowering Plants	Nye Milkvetch	<i>Astragalus nyensis</i>	No	No	G3				2		No
Dry	Flowering Plants	Charleston Milkvetch	<i>Astragalus oophorus</i> var. <i>clokeyanus</i>	No	No	T2		NV		27		No
Dry	Flowering Plants	Lavin's Egg Milkvetch	<i>Astragalus oophorus</i> var. <i>lavinii</i>	No	No	T2		CA, NV		16		No
Dry	Flowering Plants	Pink Egg Milkvetch	<i>Astragalus oophorus</i> var. <i>lonchocalyx</i>	No	No	T2		NV		17		No
Dry	Flowering Plants	Rydberg's Milkvetch	<i>Astragalus perianus</i>	No	No	G3				1		No
Dry	Flowering Plants	Pinyon Milkvetch	<i>Astragalus pinonis</i>	No	No	G2				3		No
Dry	Flowering Plants	Tonopah Milkvetch	<i>Astragalus pseudodanthus</i>	No	No	G2		CA, NV		24		No
Dry	Flowering Plants	Winged Milkvetch	<i>Astragalus pterocarpus</i>	No	No	G3				19		No
Dry	Flowering Plants	Pulsifer's Milk Vetch	<i>Astragalus pulsiferae</i> var. <i>pulsiferae</i>	No	No	T2		CA, NV		51		No
Dry	Flowering Plants	Raven's Milkvetch	<i>Astragalus ravenii</i>	No	No	G1				3		No
Dry	Flowering Plants	Lamoille Canyon Milkvetch	<i>Astragalus robbinsii</i> var. <i>occidentalis</i>	No	No	T2		NV		38		No
Dry	Flowering Plants	Weak Milkvetch	<i>Astragalus solitarius</i>	No	No	G3		NV		3		No
Dry	Flowering Plants	Silver Reef Milkvetch	<i>Astragalus straturensis</i>	No	No	G2				25		No
Dry	Flowering Plants	Tiehm milkvetch	<i>Astragalus tiehmii</i>	No	No	G3		NV				No
Dry	Flowering Plants	Toquima Milkvetch	<i>Astragalus toquimanus</i>	No	No	G2		NV		11		No
Dry	Flowering Plants	Currant Milkvetch	<i>Astragalus uncialis</i>	No	No	G2		NV		79		No
Dry	Flowering Plants	Welsh's Milkvetch	<i>Astragalus welshii</i>	No	No	G2				1		No
Dry	Flowering Plants	Mud-flat Milkvetch	<i>Astragalus yoder-williamsii</i>	No	Yes	G3		NV		1		No
Dry	Flowering Plants	Bonneville Saltbush	<i>Atriplex bonnevillensis</i>	No	No	G2						No
Dry	Flowering Plants	Last Chance Rock Cress	<i>Boechera yorkii</i>	No	No	G1				2		No
Dry	Flowering Plants	Inyo County Mariposa-lily	<i>Calochortus excavatus</i>	No	No	G3		CA		61		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Panamint Mountain Mariposa Lily	Calochortus panamintensis	No	No	G3				1		No
Dry	Flowering Plants	Baird's Camissonia	Camissonia bairdii	No	No	G1				1		No
Dry	Flowering Plants	Diamond Valley Suncup	Camissonia gouldii	No	No	G1				2		No
Dry	Flowering Plants	Nevada Evening-primrose	Camissonia nevadensis	No	No	G3				11		No
Dry	Flowering Plants	Tioga Pass Sedge	Carex tiogana	No	No	G1				4		No
Dry	Flowering Plants	Tushar Paintbrush	Castilleja parvula	No	No	G2				9		No
Dry	Flowering Plants	Reveal's Indian-paintbrush	Castilleja revealii	No	No	G2				2		No
Dry	Flowering Plants	Monte Neva Paintbrush	Castilleja salsuginosa	No	Yes	G1		NV		2		No
Dry	Flowering Plants	Barneby's Caulanthus	Caulanthus barnebyi	No	No	G2				11		No
Dry	Flowering Plants	Jaeger's Caulostramina	Caulostramina jaegeri	No	No	G1		CA		13		No
Dry	Flowering Plants	Pintwater Rabbitbrush	Chrysothamnus eremobius	No	No	G1				1		No
Dry	Flowering Plants	Ownbey's Thistle	Cirsium ownbeyi	No	No	G3				1		No
Dry	Flowering Plants	Virgin Thistle	Cirsium virginense	No	Yes	G2		NV		4		No
Dry	Flowering Plants	Pygmy Pussy-paws	Cistanthe pygmaea	No	No	G2				3		No
Dry	Flowering Plants	Barren Valley Collomia	Collomia renacta	No	No	G1		NV		2		No
Dry	Flowering Plants	Tecopa Bird's-beak	Cordylanthus tecopensis	No	No	G2		CA, NV		2		No
Dry	Flowering Plants	Compact Cat's-eye	Cryptantha compacta	No	No	G2				11		No
Dry	Flowering Plants	Subalpine Cryptantha	Cryptantha crymophila	No	No	G2				5		No
Dry	Flowering Plants	Yellow-white Catseye	Cryptantha ochroleuca	No	No	G1				1		No
Dry	Flowering Plants	Bristle-cone Cryptantha	Cryptantha roosiorum	No	Yes	G1		CA		24		No
Dry	Flowering Plants	Schoolcraft catseye	Cryptantha schoolcraftii	No	No	G3		CA, NV				No
Dry	Flowering Plants	Welsch's Cat's-eye	Cryptantha welshii	No	No	G3				42		No
Dry	Flowering Plants	Bodie Hills Cusickiella	Cusickiella quadricostata	No	No	G2		CA		54		No
Dry	Flowering Plants	Intermountain Wavewing	Cymopterus basalticus	No	No	G2		NV		19		No
Dry	Flowering Plants	Gray Wavewing	Cymopterus cinerarius	No	No	G2				3		No
Dry	Flowering Plants	Coulter's Biscuitroot	Cymopterus coulteri	No	No	G3				27		No
Dry	Flowering Plants	Toiyabe Spring-parsley	Cymopterus goodrichii	No	No	G1		NV		7		No
Dry	Flowering Plants	Jone's Wavewing	Cymopterus jonesii	No	No	G2				15		No
Dry	Flowering Plants	Cedar Breaks Biscuitroot	Cymopterus minimus	No	No	G1				15		No
Dry	Flowering Plants	Sanicle Biscuitroot	Cymopterus ripleyi var. saniculoides	No	No	T3		CA		31		No
Dry	Flowering Plants	Clustered Lady's-slipper	Cypripedium fasciculatum	No	No	G4		CA		9		No
Dry	Flowering Plants	Large Yellow Lady's-slipper	Cypripedium parviflorum var. pubescens	No	Yes	T5				4		No
Dry	Flowering Plants	ornate dalea	Dalea ornata	No	No	G4		CA				No
Dry	Flowering Plants	July Gold	Dedekera eurekaensis	No	Yes	G2		CA		50		No
Dry	Flowering Plants	Desert Whitlow-grass	Draba arida	No	No	G2				19		No
Dry	Flowering Plants	Wasatch Draba	Draba brachystylis	No	No	G1				8		No
Dry	Flowering Plants	White Mountain Draba	Draba californica	No	No	G3				1		No
Dry	Flowering Plants	Rockcross Draba	Draba globosa	No	No	G3				5		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Sweetwater Mountains Draba	<i>Draba incrassata</i>	No	No	G3				17		No
Dry	Flowering Plants	Kass's Rockcress	<i>Draba kassii</i>	No	No	G1				5		No
Dry	Flowering Plants	Maguire's Whitlow-grass	<i>Draba maguirei</i>	No	No	G3				15		No
Dry	Flowering Plants	White Mountains draba	<i>Draba monoensis</i>	No	No	G1				3		No
Dry	Flowering Plants	Pennell's Draba	<i>Draba pennellii</i>	No	No	G2				12		No
Dry	Flowering Plants	Tushar Mountain Whitlow-grass	<i>Draba ramulosa</i>	No	No	G1				5		No
Dry	Flowering Plants	Mt. Whitney Draba	<i>Draba sharsmithii</i>	No	No	G1				4		No
Dry	Flowering Plants	Sierra Nevada Draba	<i>Draba sierrae</i>	No	No	G2				12		No
Dry	Flowering Plants	Stolon Whitlow-grass	<i>Draba sobolifera</i>	No	No	G2				9		No
Dry	Flowering Plants	Mountain Whitlow-grass	<i>Draba sphaeroides</i>	No	No	G2				10		No
Dry	Flowering Plants	White Mountain Draba	<i>Draba subumbellata</i>	No	No	G3				2		No
Dry	Flowering Plants	Engelmann's Hedgehog Cactus	<i>Echinocereus engelmannii</i> var. <i>armatus</i>	No	Yes	T2				2		No
Dry	Flowering Plants	Nevada Willowherb	<i>Epilobium nevadense</i>	No	No	G2		NV		16		No
Dry	Flowering Plants	Deer Goldenweed	<i>Ericameria cervina</i>	No	No	G3		NV		12		No
Dry	Flowering Plants	Pine Valley Goldenbush	<i>Ericameria crispa</i>	No	No	G2				4		No
Dry	Flowering Plants	Gilman Goldenweed	<i>Ericameria gilmanii</i>	No	No	G1		CA		1		No
Dry	Flowering Plants	Greenwood's Heath-goldenrod	<i>Ericameria lignumviridis</i>	No	No	G1				1		No
Dry	Flowering Plants	Cedar Breaks Goldenbush	<i>Ericameria zionis</i>	No	No	G2				1		No
Dry	Flowering Plants	Bald Daisy	<i>Erigeron calvus</i>	No	No	G1				1		No
Dry	Flowering Plants	Carrington's Daisy	<i>Erigeron carringtoniae</i>	No	No	G1				1		No
Dry	Flowering Plants	Cave Mountain Fleabane	<i>Erigeron cavernensis</i>	No	No	G2				4		No
Dry	Flowering Plants	Mound Daisy	<i>Erigeron compactus</i>	No	No	G2				35		No
Dry	Flowering Plants	Cronquist's Daisy	<i>Erigeron cronquistii</i>	No	No	G2				12		No
Dry	Flowering Plants	Garrett's Daisy	<i>Erigeron garrettii</i>	No	No	G2				20		No
Dry	Flowering Plants	Broad Fleabane	<i>Erigeron latus</i>	No	No	G3		NV		5		No
Dry	Flowering Plants	Starved Daisy	<i>Erigeron miser</i>	No	No	G2				1		No
Dry	Flowering Plants	Sheep Fleabane	<i>Erigeron ovinus</i>	No	No	G2		NV		7		No
Dry	Flowering Plants	Professor Daisy	<i>Erigeron proselyticus</i>	No	No	G2				9		No
Dry	Flowering Plants	Ibex Buckwheat	<i>Eriogonum ammophilum</i>	No	No	G1				18		No
Dry	Flowering Plants	Mono Buckwheat	<i>Eriogonum ampullaceum</i>	No	No	G3				3		No
Dry	Flowering Plants	Wind-loving Buckwheat	<i>Eriogonum anemophilum</i>	No	No	G2		NV		35		No
Dry	Flowering Plants	Ruby Valley Buckwheat	<i>Eriogonum argophyllum</i>	No	Yes	G1				1		No
Dry	Flowering Plants	Beatley's Buckwheat	<i>Eriogonum beatleyae</i>	No	No	G2		NV		40		No
Dry	Flowering Plants	Darin Buckwheat	<i>Eriogonum concinnum</i>	No	No	G2		NV		16		No
Dry	Flowering Plants	Reveal's Buckwheat	<i>Eriogonum contiguum</i>	No	No	G2		CA		1		No
Dry	Flowering Plants	Crosby's Buckwheat	<i>Eriogonum crosbyae</i>	No	No	G3		CA, NV		2		No
Dry	Flowering Plants	Darrow's Buckwheat	<i>Eriogonum darrovii</i>	No	No	G2				8		No
Dry	Flowering Plants	Churchill Narrows Buckwheat	<i>Eriogonum diatomaceum</i>	Yes	Yes	G1		NV		31		No
Dry	Flowering Plants	Wildrose Canyon Buckwheat	<i>Eriogonum eremicola</i>	No	No	G1		CA		2		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Limestone Buckwheat	Eriogonum eremicum	No	No	G2				18		No
Dry	Flowering Plants	Gilman's Buckwheat	Eriogonum gilmanii	No	No	G2				5		No
Dry	Flowering Plants	Holmgren's Buckwheat	Eriogonum holmgrenii	No	No	G1				4		No
Dry	Flowering Plants	Lewis' Buckwheat	Eriogonum lewisii	No	No	G2		NV		30		No
Dry	Flowering Plants	Logan Buckwheat	Eriogonum loganum	No	No	G2				9		No
Dry	Flowering Plants	Panamint Mountains Buckwheat	Eriogonum microthecum var. panamintense	No	No	T2		CA		4		No
Dry	Flowering Plants	Slender Buckwheat	Eriogonum microthecum var. schoolcraftii	No	No	T2		CA, NV		7		No
Dry	Flowering Plants	Lost Creek Buckwheat	Eriogonum mitophyllum	No	No	G1				4		No
Dry	Flowering Plants	Son's Buckwheat	Eriogonum natum	No	No	G2				10		No
Dry	Flowering Plants	Death buckwheat	Eriogonum nutans var. glabratum	No	No	T2		NV				No
Dry	Flowering Plants	Steamboat Buckwheat	Eriogonum ovalifolium var. williamsiae	Yes	Yes	T1		NV		2		No
Dry	Flowering Plants	Wire-stem Buckwheat	Eriogonum pharnaceoides var. cervinum	No	No	T2		NV		8		No
Dry	Flowering Plants	A Buckwheat	Eriogonum phoeniceum	No	No	G1				3		No
Dry	Flowering Plants	Prostrate Buckwheat	Eriogonum prociduum	No	No	G3		CA, NV		1		No
Dry	Flowering Plants	Altered Andesite Buckwheat	Eriogonum robustum	No	No	G2		NV		156		No
Dry	Flowering Plants	Lahontan Basin Buckwheat	Eriogonum rubricaula	No	No	G3				6		No
Dry	Flowering Plants	Frisco Buckwheat	Eriogonum soledium	No	No	G1				17		No
Dry	Flowering Plants	Tiehm's Buckwheat	Eriogonum tiehmii	No	No	G1		NV		6		No
Dry	Flowering Plants	Viviparous Foxtail Cactus	Escobaria vivipara var. rosea	No	Yes	T3				54		No
Dry	Flowering Plants	Sunnyside Green-gentian	Frasera gypsicola	No	Yes	G1		NV		29		No
Dry	Flowering Plants	Kingston Bedstraw	Galium hilendiae ssp. kingstonense	No	No	T2		CA		4		No
Dry	Flowering Plants	Nye Gilia	Gilia nyensis	No	No	G3				32		No
Dry	Flowering Plants	Ripley's Gilia	Gilia ripleyi	No	No	G3				6		No
Dry	Flowering Plants	Goldenrod Snakeweed	Gutierrezia petradoria	No	No	G3				19		No
Dry	Flowering Plants	Poison Canyon Stickseed	Hackelia brevicula	No	No	G2				8		No
Dry	Flowering Plants	Deep Creek Stickseed	Hackelia ibapensis	No	No	G1				2		No
Dry	Flowering Plants	Three Forks Stickseed	Hackelia ophiobia	No	No	G3				1		No
Dry	Flowering Plants	Sharsmith's Stickseed	Hackelia sharsmithii	No	No	G3				18		No
Dry	Flowering Plants	Utah Sunflower	Helianthus deserticola	No	No	G2				16		No
Dry	Flowering Plants	White Mountains Horkelia	Horkelia hispidula	No	No	G2				21		No
Dry	Flowering Plants	Sanderson's Cheesebush	Hymenoclea sandersonii	No	No	G1				1		No
Dry	Flowering Plants	California Satintail	Imperata brevifolia	No	No	G2		NV		1		No
Dry	Flowering Plants	Sierra Valley Ivesia	Ivesia aperta var. aperta	No	No	T2		CA, NV		79		No
Dry	Flowering Plants	Rock Purpusia	Ivesia arizonica var. saxosa	No	No	T1		NV		5		No
Dry	Flowering Plants	Field Ivesia	Ivesia campestris	No	No	G3				1		No
Dry	Flowering Plants	King's Ivesia	Ivesia kingii	Yes	No	G3				2		No
Dry	Flowering Plants	King's Ivesia	Ivesia kingii var. kingii	No	No	T2		CA		15		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Pine Nut Ivesia	<i>Ivesia pityocharis</i>	No	No	G2		NV		14		No
Dry	Flowering Plants	Grimy mousetails	<i>Ivesia rhypara</i> var. <i>rhypara</i>	No	No	T2		CA, NV				No
Dry	Flowering Plants	Plumas Ivesia	<i>Ivesia sericoleuca</i>	No	No	G2		CA		68		No
Dry	Flowering Plants	Utah Ivesia	<i>Ivesia utahensis</i>	No	No	G2				4		No
Dry	Flowering Plants	Webber Ivesia	<i>Ivesia webberi</i>	Yes	Yes	G2		CA, NV		27		No
Dry	Flowering Plants	Waxflower	<i>Jamesia tetrapetala</i>	No	No	G2		NV		12		No
Dry	Flowering Plants	Grime's Vetchling	<i>Lathyrus grimesii</i>	No	No	G2				3		No
Dry	Flowering Plants	Bullfrog Hills Sweetpea	<i>Lathyrus hitchcockianus</i>	No	No	G2		NV		13		No
Dry	Flowering Plants	Mountain Pepper-grass	<i>Lepidium montanum</i> var. <i>nevadense</i>	No	No	T1		NV		6		No
Dry	Flowering Plants	Southwestern Pepper-grass	<i>Lepidium nanum</i>	No	No	G3				4		No
Dry	Flowering Plants	Ostler's Pepper-grass	<i>Lepidium ostleri</i>	No	No	G1				4		No
Dry	Flowering Plants	Owyhee Prickly-phlox	<i>Leptodactylon glabrum</i>	No	No	G2		NV		3		No
Dry	Flowering Plants	Garrett's Bladderpod	<i>Lesquerella garrettii</i>	No	No	G2				58		No
Dry	Flowering Plants	Tunnel Springs Mountain Bladderpod	<i>Lesquerella goodrichii</i>	No	No	G2				2		No
Dry	Flowering Plants	Hitchcock's Bladderpod	<i>Lesquerella hitchcockii</i>	No	No	G3						No
Dry	Flowering Plants	Snake Range Bladderpod	<i>Lesquerella pendula</i>	No	No	G2						No
Dry	Flowering Plants	Bryce Bladderpod	<i>Lesquerella rubicundula</i>	No	No	G3				1		No
Dry	Flowering Plants	Utah Bladderpod	<i>Lesquerella utahensis</i>	No	No	G3				1		No
Dry	Flowering Plants	Maguire's Bitterroot	<i>Lewisia maguirei</i>	No	No	G1				8		No
Dry	Flowering Plants	Sage-like Loefflingia	<i>Loefflingia squarrosa</i> ssp. <i>artemisiarum</i>	No	No	T2		NV		17		No
Dry	Flowering Plants	Packard's Desert-parsley	<i>Lomatium packardiae</i>	No	No	G2		NV		2		No
Dry	Flowering Plants	Rose-flower Desert-parsley	<i>Lomatium roseanum</i>	No	No	G2		CA		4		No
Dry	Flowering Plants	Mono Lake Lupine	<i>Lupinus duranii</i>	No	No	G2		CA		45		No
Dry	Flowering Plants	Slender Lupine	<i>Lupinus gracilentus</i>	No	No	G3				2		No
Dry	Flowering Plants	Holmgren Lupine	<i>Lupinus holmgrenianus</i>	No	No	G2		NV		5		No
Dry	Flowering Plants	Mcgee Meadows Lupine	<i>Lupinus magnificus</i> var. <i>hesperius</i>	No	No	T2		CA		1		No
Dry	Flowering Plants	Father Crowley's Lupine	<i>Lupinus padre-crowleyi</i>	No	Yes	G2				18		No
Dry	Flowering Plants	lilliput lupine	<i>Lupinus uncialis</i>	No	No	G4		CA				No
Dry	Flowering Plants	Pioche Blazingstar	<i>Mentzelia argillicola</i>	No	No	G1		NV		5		No
Dry	Flowering Plants	Arapien Stickleleaf	<i>Mentzelia argillosa</i>	No	No	G2				81		No
Dry	Flowering Plants	Inyo blazingstar	<i>Mentzelia inyoensis</i>	No	No	G2		CA		6		No
Dry	Flowering Plants	Smooth Stickleleaf	<i>Mentzelia mollis</i>	No	No	G2		NV		3		No
Dry	Flowering Plants		<i>Mentzelia tiehmii</i>	No	No	G1		NV		7		No
Dry	Flowering Plants	Three-tooth Blazingstar	<i>Mentzelia tridentata</i>	No	No	G2		CA		1		No
Dry	Flowering Plants	Eggleaf Monkeyflower	<i>Mimulus ovatus</i>	No	No	G1				9		No
Dry	Flowering Plants	Bashful Four-o'clock	<i>Mirabilis pudica</i>	No	No	G3				1		No
Dry	Flowering Plants	sweet-smelling monardella	<i>Monardella beneolens</i>	No	No	G1		CA		5		No
Dry	Flowering Plants	Rydberg's Musineon	<i>Musineon lineare</i>	No	No	G2				24		No
Dry	Flowering Plants	Eureka Dunes Evening-primrose	<i>Oenothera californica</i> ssp.	Yes	Yes	T1				3		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
			eurekensis									
Dry	Flowering Plants	Sand Cholla	Opuntia pulchella	No	Yes	G4		NV		54		No
Dry	Flowering Plants	Plumas Mountaintop	Oreostemma elatum	No	No	G2		CA		1		No
Dry	Flowering Plants	Nevada Oryctes	Oryctes nevadensis	No	No	G2		NV		111		No
Dry	Flowering Plants	Beaver Mountain Groundsel	Packera castoreus	No	No	G1				2		No
Dry	Flowering Plants	Podunk Groundsel	Packera malmstenii	No	No	G1				1		No
Dry	Flowering Plants	Ligulate Feverfew	Parthenium ligulatum	No	No	G3		NV		1		No
Dry	Flowering Plants	dwarf lousewort	Pedicularis centranthera	No	No	G4		CA				No
Dry	Flowering Plants	Siler Pincushion Cactus	Pediocactus sileri	Yes	Yes	G3				2		No
Dry	Flowering Plants	Simpson's Hedgehog Cactus	Pediocactus simpsonii	No	Yes	G4				7		No
Dry	Flowering Plants	Firleaf Beardtongue	Penstemon abietinus	No	No	G2				12		No
Dry	Flowering Plants	Dune Beardtongue	Penstemon arenarius	No	No	G2		NV		32		No
Dry	Flowering Plants	Red Canyon Beardtongue	Penstemon bracteatus	No	No	G2				2		No
Dry	Flowering Plants	Limestone Beardtongue	Penstemon calcareus	No	No	G2				8		No
Dry	Flowering Plants	Bear River Range Beardtongue	Penstemon compactus	No	No	G2				19		No
Dry	Flowering Plants	Tunnel Springs Beardtongue	Penstemon concinnus	No	No	G3		NV		22		No
Dry	Flowering Plants	Cordelia's Penstemon	Penstemon floribundus	No	No	G1		NV		8		No
Dry	Flowering Plants	Ben Franklin's Beardtongue	Penstemon franklinii	No	No	G1				7		No
Dry	Flowering Plants	Death Valley Beardtongue	Penstemon fruticiformis ssp. amargosae	No	No	T3		NV		1		No
Dry	Flowering Plants	Janish's beardtongue	Penstemon janishiae	No	No	G4		CA				No
Dry	Flowering Plants	Charleston Beardtongue	Penstemon leiophyllus var. francisci-pennellii	No	No	T2		NV		8		No
Dry	Flowering Plants	Mt. Moriah Beardtongue	Penstemon moriahensis	No	No	G1				8		No
Dry	Flowering Plants	Low Beardtongue	Penstemon nanus	No	No	G3				31		No
Dry	Flowering Plants	Pahute Mesa Beardtongue	Penstemon pahutensis	No	No	G3		NV		48		No
Dry	Flowering Plants	Lahontan Beardtongue	Penstemon palmeri var. macranthus	No	No	T2		NV		26		No
Dry	Flowering Plants	Petiolate Beardtongue	Penstemon petiolatus	No	No	G2		AZ		9		No
Dry	Flowering Plants	Pinyon Penstemon	Penstemon pinorum	No	No	G1				36		No
Dry	Flowering Plants	Broadleaf Beardtongue	Penstemon platyphyllus	No	No	G2				35		No
Dry	Flowering Plants	Kawich Range Beardtongue	Penstemon pudicus	No	No	G1		NV		6		No
Dry	Flowering Plants	Rhizome Beardtongue	Penstemon rhizomatosus	No	No	G1				6		No
Dry	Flowering Plants	Wassuk Beardtongue	Penstemon rubicundus	No	No	G2		NV		22		No
Dry	Flowering Plants	Susanville Beardtongue	Penstemon sudans	No	No	G3		CA, NV		5		No
Dry	Flowering Plants	Jaeger's Beardtongue	Penstemon thompsoniae ssp. jaegeri	No	No	T2		NV		1		No
Dry	Flowering Plants	Tidestrom Beardtongue	Penstemon tidestromii	No	No	G2				14		No
Dry	Flowering Plants	Shoshone Beardtongue	Penstemon tiehmii	No	No	G1		NV		3		No
Dry	Flowering Plants	Tushar Range Beardtongue	Penstemon tusharensis	No	No	G2				4		No
Dry	Flowering Plants	Ward Beardtongue	Penstemon wardii	No	No	G2				33		No
Dry	Flowering Plants	Inyo Rock Daisy	Perityle inyoensis	No	No	G2		CA		6		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Hanaupah rock daisy	Perityle villosa	No	No	G1		CA		2		No
Dry	Flowering Plants	Death Valley Sandpaper-plant	Petalonyx thurberi ssp. gilmanii	No	No	T2		CA		4		No
Dry	Flowering Plants	marble rockmat	Petrophyton acuminatum	No	No	G1				1		No
Dry	Flowering Plants	Aven Nelson's Phacelia	Phacelia anelsonii	No	No	G2				7		No
Dry	Flowering Plants	Beatley's Phacelia	Phacelia beatleyae	No	No	G3				13		No
Dry	Flowering Plants		Phacelia filiae	No	No	G2		NV		4		No
Dry	Flowering Plants	Inconspicuous Scorpionweed	Phacelia inconspicua	No	Yes	G2		NV		4		No
Dry	Flowering Plants	Playa Phacelia	Phacelia inundata	No	No	G2		CA, NV		3		No
Dry	Flowering Plants	Inyo Phacelia	Phacelia inyoensis	No	No	G3		CA		23		No
Dry	Flowering Plants	Tiny-flower Phacelia	Phacelia minutissima	No	No	G3		NV		30		No
Dry	Flowering Plants	Mono County Phacelia	Phacelia monoensis	No	No	G3		CA		43		No
Dry	Flowering Plants	Death Valley Roundleaf Phacelia	Phacelia mustelina	No	No	G2		CA, NV		9		No
Dry	Flowering Plants	Parish's Phacelia	Phacelia parishii	No	No	G2		AZ, CA, NV		9		Yes
Dry	Flowering Plants	Utah Phacelia	Phacelia utahensis	No	No	G2				190		No
Dry	Flowering Plants	Repand Twinpod	Physaria repanda	No	No	G1				1		No
Dry	Flowering Plants	Clustered Popcorn-flower	Plagiobothrys glomeratus	No	No	G2		NV		11		No
Dry	Flowering Plants	Parish's Popcorn-flower	Plagiobothrys parishii	No	No	G1				6		No
Dry	Flowering Plants	Desert Allocarya	Plagiobothrys salsus	No	No	G2				1		No
Dry	Flowering Plants	Mason's Skypilot	Polemonium chartaceum	No	No	G1				14		No
Dry	Flowering Plants	Spiny Milkwort	Polygala heterorhyncha	No	No	G3				5		No
Dry	Flowering Plants	Pygmy Poreleaf	Porophyllum pygmaeum	No	No	G2				1		No
Dry	Flowering Plants	Soldier Meadows Cinquefoil	Potentilla basaltica	Yes	No	G1		CA, NV		9		No
Dry	Flowering Plants	Cottam's Potentilla	Potentilla cottamii	No	No	G1		NV		6		No
Dry	Flowering Plants	Morefield's Cinquefoil	Potentilla morefieldii	No	No	G1				17		No
Dry	Flowering Plants	Ruby Mountains Primrose	Primula capillaris	No	No	G1				8		No
Dry	Flowering Plants	House Range Primrose	Primula domensis	No	No	G1				5		No
Dry	Flowering Plants	Maguire's Primrose	Primula maguirei	Yes	No	G1				14		No
Dry	Flowering Plants	Nevada Primrose	Primula nevadensis	No	No	G2				10		No
Dry	Flowering Plants	King's Indigo-bush	Psorothamnus kingii	No	No	G3				10		No
Dry	Flowering Plants	Sticky Haplopappus	Pyrrocoma lucida	No	No	G3		CA		83		No
Dry	Flowering Plants	Obscure Buttercup	Ranunculus glaberrimus var. reconditus	No	No	T2		NV		1		No
Dry	Flowering Plants	Blaine's Pincushion	Sclerocactus blainei	No	Yes	G1		NV		4		No
Dry	Flowering Plants	Nye County Fish-hook Cactus	Sclerocactus nyensis	No	Yes	G1		NV		3		No
Dry	Flowering Plants	Mohave Fishhook Cactus	Sclerocactus polyancistrus	No	Yes	G4				23		No
Dry	Flowering Plants	Great Basin Fishhook Cactus	Sclerocactus pubispinus	No	Yes	G4		NV		36		No
Dry	Flowering Plants	Schlesser's Pincushion	Sclerocactus schlesseri	No	Yes	G1		NV		13		No
Dry	Flowering Plants	Desert Valley Fishhook Cactus	Sclerocactus spinosior	No	No	G2				18		No
Dry	Flowering Plants	Musinea Ragwort	Senecio musiniensis	No	No	G1				3		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Dry	Flowering Plants	Mono Ragwort	Senecio pattersonensis	No	No	G2				12		No
Dry	Flowering Plants	Owens Valley Checker-mallow	Sidalcea covillei	No	Yes	G3		CA		52		No
Dry	Flowering Plants	Jan's Catchfly	Silene nachlingerae	No	No	G2		NV		19		No
Dry	Flowering Plants	Peterson's Catchfly	Silene petersonii	No	No	G2				11		No
Dry	Flowering Plants	Funeral Mountain Blue-eyed-grass	Sisyrinchium funereum	No	No	G2				3		No
Dry	Flowering Plants	Big-root Blue-eyed-grass	Sisyrinchium radicatum	No	No	G2		NV		1		No
Dry	Flowering Plants	Nye County Smelowskia	Smelowskia holmgrenii	No	No	G2		NV		18		No
Dry	Flowering Plants	Jone's Globemallow	Sphaeralcea caespitosa	No	No	G2				13		No
Dry	Flowering Plants	Jone's Globemallow	Sphaeralcea caespitosa var. williamsiae	No	No	T2		NV		6		No
Dry	Flowering Plants	Ute Ladies'-tresses	Spiranthes diluvialis	Yes	Yes	G2		NV		18		No
Dry	Flowering Plants	Hooded Ladies'-tresses	Spiranthes romanzoffiana	No	Yes	G5				1		No
Dry	Flowering Plants		Stipa shoshoneana	No	No	G2				1		No
Dry	Flowering Plants	Alpine Jewelflower	Streptanthus gracilis	No	No	G3				3		No
Dry	Flowering Plants	Masonic Mountain Jewelflower	Streptanthus oliganthus	No	No	G2		CA		32		No
Dry	Flowering Plants	Tiehm's Stroganowia	Stroganowia tiehmii	No	No	G2		NV		43		No
Dry	Flowering Plants	Eureka Dunes Grass	Swallenia alexandrae	Yes	Yes	G1				5		No
Dry	Flowering Plants	Welsh's American-aster	Symphyotrichum welshii	No	No	G2				5		No
Dry	Flowering Plants	Alpine Goldenweed	Tonestus alpinus	No	No	G2				11		No
Dry	Flowering Plants	Granite Haplopappus	Tonestus graniticus	No	No	G1		NV		2		No
Dry	Flowering Plants	Currant Summit Clover	Trifolium andinum var. podocephalum	No	No	T1		NV		3		No
Dry	Flowering Plants	Dedecker's Clover	Trifolium dedeckerae	No	No	G2		CA		13		No
Dry	Flowering Plants	Frisco Clover	Trifolium friscanum	No	No	G1				6		No
Dry	Flowering Plants	Leiberg's Clover	Trifolium leibergii	No	No	G2				13		No
Dry	Flowering Plants	Rollins Clover	Trifolium rollinsii	No	No	G2				13		No
Dry	Flowering Plants	Frank Smith's Violet	Viola frank-smithii	No	No	G1				31		No
Dry	Flowering Plants	Rock Violet	Viola lithion	No	No	G1		NV		6		No
Dry	Mosses		Bruchia bolanderi	No	No	G3				1		No
Dry	Mosses		Orthotrichum shevockii	No	No	G1		CA, NV		4		No
Dry	Mosses		Orthotrichum spjutii	No	No	G1				1		No
Dry	Mosses		Pohlia tundrae	No	No	G2				1		No
Wet	Amphibians	Mountain Yellow-legged Frog	Rana muscosa	Yes	No	G2					SW, CA	No
Wet	Amphibians	Northern Leopard Frog	Rana pipiens	No	Yes	G5	CA, ID, NV, UT	UT	PS	164	SW, CA	No
Wet	Amphibians	Great Basin Spadefoot	Spea intermontana	No	No	G5		CA			SW, CA	No
Wet	Birds	Clark's Grebe	Aechmophorus clarkii	No	Yes	G5	ID, NV					No
Wet	Birds	Western Grebe	Aechmophorus occidentalis	No	Yes	G5	ID, NV					No
Wet	Birds	Wood Duck	Aix sponsa	No	Yes	G5						No
Wet	Birds	Northern Pintail	Anas acuta	No	Yes	G5	ID, NV					No
Wet	Birds	American Wigeon	Anas americana	No	Yes	G5						No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Wet	Birds	Northern Shoveler	Anas clypeata	No	Yes	G5						No
Wet	Birds	Cinnamon Teal	Anas cyanoptera	No	Yes	G5	NV					No
Wet	Birds	Blue-winged Teal	Anas discors	No	Yes	G5						No
Wet	Birds	Great Blue Heron	Ardea herodias	No	Yes	G5	CA					No
Wet	Birds	Lesser Scaup	Aythya affinis	No	Yes	G5	ID					No
Wet	Birds	Redhead	Aythya americana	No	Yes	G5	NV		PS		SW	No
Wet	Birds	Canvasback	Aythya valisineria	No	Yes	G5	CA, NV					No
Wet	Birds	American Bittern	Botaurus lentiginosus	No	Yes	G4	CA					No
Wet	Birds	Canada Goose	Branta canadensis	No	Yes	G5						No
Wet	Birds	Barrow's Goldeneye	Bucephala islandica	No	Yes	G5	CA					No
Wet	Birds	Green Heron	Butorides virescens	No	Yes	G5				3	SW	No
Wet	Birds	Least Sandpiper	Calidris minutilla	No	Yes	G5	NV					No
Wet	Birds	Black Tern	Chlidonias niger	No	Yes	G4	CA, ID, NV		PS	7	SW	No
Wet	Birds	American Dipper	Cinclus mexicanus	No	Yes	G5						No
Wet	Birds	Trumpeter Swan	Cygnus buccinator	No	Yes	G4	ID			10	SW	No
Wet	Birds	Wilson's Snipe	Gallinago delicata	No	Yes	G5						No
Wet	Birds	Common Loon	Gavia immer	No	Yes	G5	CA, ID, NV		PS	2	SW	No
Wet	Birds	Whooping Crane	Grus americana	Yes	Yes	G1	UT			1	SW	No
Wet	Birds	Sandhill Crane	Grus canadensis	No	Yes	G5	ID			3	SW	No
Wet	Birds	Greater Sandhill Crane	Grus canadensis tabida	No	Yes	T4	CA, NV	CA	PS	23		No
Wet	Birds	Black-necked Stilt	Himantopus mexicanus	No	Yes	G5	ID, NV, UT		PS	10	SW	No
Wet	Birds	Harlequin Duck	Histrionicus histrionicus	No	Yes	G4	CA, ID			4		No
Wet	Birds	Caspian Tern	Hydroprogne caspia	No	Yes	G5	CA, ID, UT			12	SW	No
Wet	Birds	Least Bittern	Ixobrychus exilis	No	Yes	G5	CA			5	SW	No
Wet	Birds	Western Least Bittern	Ixobrychus exilis hesperis	No	Yes	T3	NV		PS	3		No
Wet	Birds	California Gull	Larus californicus	No	Yes	G5	CA, ID			2	SW	No
Wet	Birds	Franklin's Gull	Leucophaeus pipixcan	No	Yes	G4	ID, NV			1	SW	No
Wet	Birds	Long-billed Dowitcher	Limnodromus scolopaceus	No	Yes	G5	NV					No
Wet	Birds	Common Merganser	Mergus merganser	No	Yes	G5				1	SW	No
Wet	Birds	Wood Stork	Mycteria americana	No	Yes	G4	CA			2		No
Wet	Birds	Black-crowned Night-Heron	Nycticorax nycticorax	No	Yes	G5	CA, ID			2	SW	No
Wet	Birds	American White Pelican	Pelecanus erythrorhynchos	No	Yes	G4	CA, ID, NV, UT		MV	79	SW	No
Wet	Birds	Brown Pelican	Pelecanus occidentalis	No	Yes	G4				1		No
Wet	Birds	Double-crested Cormorant	Phalacrocorax auritus	No	Yes	G5	CA					No
Wet	Birds	red-necked phalarope	Phalaropus lobatus	No	Yes	G4	NV		MV			No
Wet	Birds	Wilson's Phalarope	Phalaropus tricolor	No	Yes	G5	ID				SW	No
Wet	Birds	White-faced Ibis	Plegadis chihi	No	Yes	G5	CA, ID, NV		PS	5	SW	No
Wet	Birds	Horned Grebe	Podiceps auritus	No	Yes	G5					SW	No
Wet	Birds	Eared Grebe	Podiceps nigricollis	No	Yes	G5	NV		PS	1	SW	No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
Wet	Birds	American Avocet	Recurvirostra americana	No	Yes	G5	ID, NV, UT		PS	40	SW	No
Wet	Birds	Forster's Tern	Sterna forsteri	No	Yes	G5	CA, ID, NV		PS	1	SW	No
Wet	Birds	Willet	Tringa semipalmata	No	Yes	G5	NV					No
Wet	Caddisflies	Denning's Cryptic Caddisfly	Cryptochia denningi	No	No	G1				1		No
Wet	Fairy, Clam, & Tadpole Shrimps	Mono Lake Brine Shrimp	Artemia monica	No	No	G1				1		No
Wet	Freshwater & Anadromous Fishes	Desert Sucker	Catostomus clarkii	No	Yes	G3		AZ, UT		172		No
Wet	Freshwater & Anadromous Fishes	White River Desert Sucker	Catostomus clarkii intermedius	No	Yes	T1			HV	10		No
Wet	Freshwater & Anadromous Fishes	Meadow Valley Wash Desert Sucker	Catostomus clarkii ssp. 2	No	Yes	T2				12		No
Wet	Freshwater & Anadromous Fishes	Bluehead Sucker	Catostomus discobolus	No	Yes	G4		UT		7		No
Wet	Freshwater & Anadromous Fishes	Owens Sucker	Catostomus fumeiventris	No	No	G3				19		No
Wet	Freshwater & Anadromous Fishes	Flannelmouth Sucker	Catostomus latipinnis	No	Yes	G3		AZ, UT	PS	45		No
Wet	Freshwater & Anadromous Fishes	Wall Canyon sucker	Catostomus sp. 1	No	No	G1			MV			No
Wet	Freshwater & Anadromous Fishes	Cui-ui	Chasmistes cujus	Yes	Yes	G1			MV	1		No
Wet	Freshwater & Anadromous Fishes	June Sucker	Chasmistes liorus	Yes	Yes	G1				10		No
Wet	Freshwater & Anadromous Fishes	White River Sculpin	Cottus sp. 3	No	No	G1				1		No
Wet	Freshwater & Anadromous Fishes	Preston White River Springfish	Crenichthys baileyi albivallis	No	Yes	T1			PS	6		No
Wet	Freshwater & Anadromous Fishes	White River Springfish	Crenichthys baileyi baileyi	Yes	Yes	T1			PS	2		Yes
Wet	Freshwater & Anadromous Fishes	Hiko White River Springfish	Crenichthys baileyi grandis	Yes	Yes	T1			PS	3		Yes
Wet	Freshwater & Anadromous Fishes	Moorman White River Springfish	Crenichthys baileyi thermophilus	No	Yes	T1			PS	3		No
Wet	Freshwater & Anadromous Fishes	Railroad Valley Springfish	Crenichthys nevadae	Yes	Yes	G2			PS	18		Yes
Wet	Freshwater & Anadromous Fishes	Amargosa Pupfish	Cyprinodon nevadensis amargosae	No	No	T1		CA		1		No
Wet	Freshwater & Anadromous Fishes	Owens River Pupfish	Cyprinodon radiosus	Yes	Yes	G1		CA		17		No
Wet	Freshwater & Anadromous Fishes	Pahrump Poolfish	Empetrichthys latos latos	Yes	Yes	T1			MV	1		No
Wet	Freshwater & Anadromous Fishes	Desert Dace	Eremichthys acros	Yes	Yes	G1			MV	11		Yes
Wet	Freshwater & Anadromous Fishes	Alvord Chub	Gila alvordensis	No	No	G2				2		No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Wet	Freshwater & Anadromous Fishes	Fish Creek Springs Tui Chub	<i>Gila bicolor euchila</i>	No	Yes	T1				1		No
Wet	Freshwater & Anadromous Fishes	Independence Valley Tui Chub	<i>Gila bicolor isolata</i>	No	Yes	T1			PS	1		No
Wet	Freshwater & Anadromous Fishes	Newark Valley Tui Chub	<i>Gila bicolor newarkensis</i>	No	Yes	T1				21		No
Wet	Freshwater & Anadromous Fishes	Lahontan Creek Tui Chub	<i>Gila bicolor obesa</i>	No	Yes	T4				6		No
Wet	Freshwater & Anadromous Fishes	Owens Tui Chub	<i>Gila bicolor snyderi</i>	Yes	Yes	T1		CA		16		Yes
Wet	Freshwater & Anadromous Fishes	Fish Lake Valley Tui Chub	<i>Gila bicolor</i> ssp. 4	No	Yes	T1			PS	1		No
Wet	Freshwater & Anadromous Fishes	Hot Creek Valley Tui Chub	<i>Gila bicolor</i> ssp. 5	No	Yes	T1				4		No
Wet	Freshwater & Anadromous Fishes	Little Fish Lake Valley Tui Chub	<i>Gila bicolor</i> ssp. 6	No	Yes	T1				1		No
Wet	Freshwater & Anadromous Fishes	Railroad Valley Tui Chub	<i>Gila bicolor</i> ssp. 7	No	Yes	T1			MV	7		No
Wet	Freshwater & Anadromous Fishes	Big Smokey Valley Tui Chub	<i>Gila bicolor</i> ssp. 8	No	Yes	T1			HV	5		No
Wet	Freshwater & Anadromous Fishes	Bonytail	<i>Gila elegans</i>	Yes	Yes	G1			PS	2		No
Wet	Freshwater & Anadromous Fishes	Roundtail Chub	<i>Gila robusta</i>	Yes	Yes	G3		UT		10		No
Wet	Freshwater & Anadromous Fishes	A Roundtail Chub	<i>Gila robusta jordani</i>	Yes	Yes	T1			PS	5		No
Wet	Freshwater & Anadromous Fishes	Virgin River Chub	<i>Gila seminuda</i>	Yes	Yes	G1			PS	31		Yes
Wet	Freshwater & Anadromous Fishes	Least Chub	<i>Iotichthys phlegethontis</i>	Yes	Yes	G1		UT		53		No
Wet	Freshwater & Anadromous Fishes	Red Hills Roach	<i>Lavinia symmetricus</i> ssp. 3	No	No	T1		CA				No
Wet	Freshwater & Anadromous Fishes	White River Spinedace	<i>Lepidomeda albivallis</i>	Yes	Yes	G1			PS	8		Yes
Wet	Freshwater & Anadromous Fishes	Southern Leatherside Chub	<i>Lepidomeda aliciae</i>	No	Yes	G2		UT		61		No
Wet	Freshwater & Anadromous Fishes	Northern Leatherside Chub	<i>Lepidomeda copei</i>	No	Yes	G1		UT		1		No
Wet	Freshwater & Anadromous Fishes	Virgin Spinedace	<i>Lepidomeda mollispinis</i>	Yes	Yes	G1				109		No
Wet	Freshwater & Anadromous Fishes	Virgin River Spinedace	<i>Lepidomeda mollispinis mollispinis</i>	No	Yes	T1		UT	PS	2		No
Wet	Freshwater & Anadromous Fishes	Big Spring Spinedace	<i>Lepidomeda mollispinis pratensis</i>	Yes	Yes	T1			MV	3		Yes
Wet	Freshwater & Anadromous Fishes	Moapa Dace	<i>Moapa coriacea</i>	Yes	Yes	G1			PS	2		No
Wet	Freshwater &	Lahontan Cutthroat Trout	<i>Oncorhynchus clarkii henshawi</i>	Yes	Yes	T3			MV	149		No

Ecoregion Model Group	Taxonomic Group	Common Name	Scientific Name	Federally Listed	State Protected	Rounded Global Rank	Relevant SWAPs	Relevant BLM Special Status	NatureServe Climate Vulnerability Index	# of Natural Heritage Locations	Available GAP Habitat Models	Other Spatial Data
	Anadromous Fishes											
Wet	Freshwater & Anadromous Fishes	Paiute Cutthroat Trout	Oncorhynchus clarkii seleniris	Yes	No	T1				10		No
Wet	Freshwater & Anadromous Fishes	Bonneville Cutthroat Trout	Oncorhynchus clarkii utah	No	Yes	T4		UT		197		No
Wet	Freshwater & Anadromous Fishes	Inland Redband Trout & Redband Steelhead	Oncorhynchus mykiss gairdneri	No	Yes	T4				1		No
Wet	Freshwater & Anadromous Fishes	Woundfin	Plagopterus argentissimus	Yes	Yes	G1			PS	29		Yes
Wet	Freshwater & Anadromous Fishes	Relict Dace	Relictus solitarius	No	Yes	G2				49		No
Wet	Freshwater & Anadromous Fishes	Speckled Dace	Rhinichthys osculus	Yes	No	G5		AZ		189		No
Wet	Freshwater & Anadromous Fishes	Big Smokey Valley Speckled Dace	Rhinichthys osculus lariversi	No	Yes	T1			HV	4		No
Wet	Freshwater & Anadromous Fishes	Independence Valley Speckled Dace	Rhinichthys osculus lethoporus	Yes	Yes	T1			HV	1		No
Wet	Freshwater & Anadromous Fishes	Clover Valley Speckled Dace	Rhinichthys osculus oligoporus	Yes	Yes	T1			HV	4		No
Wet	Freshwater & Anadromous Fishes	Lahontan Speckled Dace	Rhinichthys osculus robustus	No	Yes	T5						No
Wet	Freshwater & Anadromous Fishes	Diamond Valley Speckled Dace	Rhinichthys osculus ssp. 10	No	No	TH			HV	1		No
Wet	Freshwater & Anadromous Fishes	Owens Speckled Dace	Rhinichthys osculus ssp. 2	No	No	T1		CA		24		No
Wet	Freshwater & Anadromous Fishes	Monitor Valley Speckled Dace	Rhinichthys osculus ssp. 5	No	No	T1			HV	2		No
Wet	Freshwater & Anadromous Fishes	White River Speckled Dace	Rhinichthys osculus ssp. 7	No	No	T2			MV	20		No
Wet	Freshwater & Anadromous Fishes	Pahranagat Speckled Dace	Rhinichthys osculus velifer	No	Yes	T1			PS	6		No
Wet	Freshwater & Anadromous Fishes	A Speckled Dace	Rhinichthys sp. 3	No	No	G1				3		No
Wet	Freshwater & Anadromous Fishes	Bull Trout	Salvelinus confluentus	Yes	Yes	G3			HV	1		No
Wet	Freshwater & Anadromous Fishes	bull trout	Salvelinus confluentus pop. 4	Yes	Yes	T2			HV			No
Wet	Freshwater Mussels	California Floater	Anodonta californiensis	No	Yes	G3			MV	16		No
Wet	Freshwater Mussels	Western Pearlshell	Margaritifera falcata	No	Yes	G4				3		No
Wet	Freshwater Snails	Badwater Snail	Assiminea infima	No	No	G1			PS	1		No
Wet	Freshwater Snails	Steptoe Hydrobe	Eremopyrgus eganensis	No	No	G1			PS	4		No
Wet	Freshwater Snails	Green River Pebblesnail	Fluminicola coloradoensis	No	No	G2				5		No
Wet	Freshwater Snails	Pyramid Lake Pebblesnail	Fluminicola dalli	No	No	G1			HV	2		No
Wet	Freshwater Snails	Pinhead Pebblesnail	Fluminicola sp. 21	No	No	G1						No
Wet	Freshwater Snails	turban pebblesnail	Fluminicola turbiniformis	No	No	G3			HV			No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Wet	Freshwater Snails	Virginia Mountains Pebblesnail	Fluminicola virginius	No	No	G1			HV	1		No
Wet	Freshwater Snails	Deep Springs Snail	Fontelicella sp. 6	No	No	G1				1		No
Wet	Freshwater Snails	Great Basin Rams-horn	Helisoma newberryi	No	No	G1				1		No
Wet	Freshwater Snails	smooth juga	Juga interioris	No	No	G1			EV			No
Wet	Freshwater Snails	Utah Physa	Physa gyrina utahensis	No	Yes	T2				6		No
Wet	Freshwater Snails	Cloaked Physa	Physa megalochlamys	No	Yes	G3				1		No
Wet	Freshwater Snails	Lamb Rams-horn	Planorbella oregonensis	No	No	G1				1		No
Wet	Freshwater Snails	Benton Valley Springsnail	Pyrgulopsis aardahli	No	No	G1				1		No
Wet	Freshwater Snails	Duckwater Pyrg	Pyrgulopsis aloba	No	No	G1			PS	2		No
Wet	Freshwater Snails	Southern Duckwater Pyrg	Pyrgulopsis anatina	No	No	G1			PS	1		No
Wet	Freshwater Snails	Longitudinal Gland Pyrg	Pyrgulopsis anguina	No	Yes	G1			EV	3		No
Wet	Freshwater Snails	Elongate Cain Spring Pyrg	Pyrgulopsis augustae	No	No	G1			EV	1		No
Wet	Freshwater Snails	Pleasant Valley Pyrg	Pyrgulopsis aurata	No	No	G1			EV	1		No
Wet	Freshwater Snails	Large Gland Carico Pyrg	Pyrgulopsis basiglans	No	No	G1			EV	2		No
Wet	Freshwater Snails	Small Gland Carico Pyrg	Pyrgulopsis bifurcata	No	No	G1			EV	1		No
Wet	Freshwater Snails	Flat Pyrg	Pyrgulopsis breviloba	No	No	G1			EV	3		No
Wet	Freshwater Snails	Fly Ranch Pyrg	Pyrgulopsis bruesi	No	No	G1			HV	1		No
Wet	Freshwater Snails	Cortez Hills Pebblesnail	Pyrgulopsis bryantwalkerii	No	No	G1			EV	1		No
Wet	Freshwater Snails	Smooth Glenwood Pyrg	Pyrgulopsis chamberlini	No	Yes	G1				1		No
Wet	Freshwater Snails	Transverse Gland Pyrg	Pyrgulopsis cruciglans	No	No	G1			EV	4		No
Wet	Freshwater Snails	Desert Springsnail	Pyrgulopsis deserta	No	Yes	G2		AZ		4		No
Wet	Freshwater Snails	Dixie Valley Pyrg	Pyrgulopsis dixensis	No	No	G1			MV	1		No
Wet	Freshwater Snails	Smoke Creek Pyrg	Pyrgulopsis eremica	No	No	G2				5		No
Wet	Freshwater Snails	Otter Creek Pyrg	Pyrgulopsis fusca	No	Yes	G1				1		No
Wet	Freshwater Snails	Emigrant Pyrg	Pyrgulopsis gracilis	No	No	G1			PS	2		No
Wet	Freshwater Snails	Hamlin Valley Pyrg	Pyrgulopsis hamlinensis	No	Yes	G1				1		No
Wet	Freshwater Snails	Upper Thousand Spring Pyrg	Pyrgulopsis hovinghi	No	No	G1			EV			No
Wet	Freshwater Snails	Hubbs Pyrg	Pyrgulopsis hubbsi	No	No	G1		AZ	PS	2		No
Wet	Freshwater Snails	Humboldt Pyrg	Pyrgulopsis humboldtensis	No	No	G1			EV	4		No
Wet	Freshwater Snails	Kings River Pyrg	Pyrgulopsis imperialis	No	No	G1			EV	2		No
Wet	Freshwater Snails	Carinate Glenwood Pyrg	Pyrgulopsis inopinata	No	Yes	G1				2		No
Wet	Freshwater Snails	Toquerville Springsnail	Pyrgulopsis kolobensis	No	No	G5		AZ		81		No
Wet	Freshwater Snails	Landyes Pyrg	Pyrgulopsis landyei	No	No	G1			PS	1		No
Wet	Freshwater Snails	Butterfield Pyrg	Pyrgulopsis lata	No	No	G1			EV	1		No
Wet	Freshwater Snails	Crittenden springsnail	Pyrgulopsis lentiglans	No	No	G1			EV			No
Wet	Freshwater Snails	Elko Pyrg	Pyrgulopsis leporina	No	No	G1			EV	2		No
Wet	Freshwater Snails	Squat Mud Meadows Pyrg	Pyrgulopsis limaria	No	No	G1			HV	5		No
Wet	Freshwater Snails	Lockes Pyrg	Pyrgulopsis lockensis	No	No	G1			PS	1		No
Wet	Freshwater Snails	Long Valley Pyrg	Pyrgulopsis longae	No	No	G1				1		No
Wet	Freshwater Snails	Western Lahontan Pyrg	Pyrgulopsis longiglans	No	No	G2				13		No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Wet	Freshwater Snails	Hardy Pyrg	Pyrgulopsis marcida	No	No	G1			EV	7		No
Wet	Freshwater Snails	Pahrnagat Pebblesnail	Pyrgulopsis merriami	No	No	G1		AZ	PS	6		No
Wet	Freshwater Snails	Oasis Valley Springsnail	Pyrgulopsis micrococcus	No	No	G3		AZ	MV	4		No
Wet	Freshwater Snails	Northern Soldier Meadow Pyrg	Pyrgulopsis militaris	No	No	G1			HV	1		No
Wet	Freshwater Snails	Twentyone Mile Pyrg	Pyrgulopsis millenaria	No	No	G1			EV			No
Wet	Freshwater Snails	Camp Valley Pyrg	Pyrgulopsis montana	No	No	G1			EV	1		No
Wet	Freshwater Snails	Neritiform Steptoe Ranch Pyrg	Pyrgulopsis neritella	No	No	G1			PS	1		No
Wet	Freshwater Snails	Ninemile Pyrg	Pyrgulopsis nonaria	No	Yes	G1				2		No
Wet	Freshwater Snails	Elongate Mud Meadows Pyrg	Pyrgulopsis notidicola	Yes	No	G1			HV	2		No
Wet	Freshwater Snails	Sub-globose Steptoe Ranch Pyrg	Pyrgulopsis orbiculata	No	No	G1			PS	2		No
Wet	Freshwater Snails	Owens Valley Springsnail	Pyrgulopsis owensensis	No	No	G1				11		No
Wet	Freshwater Snails	Big Warm Spring Pyrg	Pyrgulopsis papillata	No	No	G1			PS	2		No
Wet	Freshwater Snails	Bifid Duct Pyrg	Pyrgulopsis peculiaris	No	Yes	G2			EV	8		No
Wet	Freshwater Snails	Antelope Valley Pyrg	Pyrgulopsis pellita	No	No	G1			EV	1		No
Wet	Freshwater Snails	Fish Slough Springsnail	Pyrgulopsis perturbata	No	No	G1				3		No
Wet	Freshwater Snails	Ovate Cain Spring Pyrg	Pyrgulopsis pictilis	No	No	G1			EV	1		No
Wet	Freshwater Snails	Flat-topped Steptoe Pyrg	Pyrgulopsis planulata	No	No	G1			PS	1		No
Wet	Freshwater Snails	Fish Lake Pyrg	Pyrgulopsis ruinosa	No	No	GX			MV	1		No
Wet	Freshwater Snails	Sada's Pyrg	Pyrgulopsis sadai	No	No	G1			EV	6		No
Wet	Freshwater Snails	White River Valley Pyrg	Pyrgulopsis sathos	No	No	G1			EV	6		No
Wet	Freshwater Snails	Sub-globose Snake Pyrg	Pyrgulopsis saxatilis	No	Yes	G1				1		No
Wet	Freshwater Snails	Northern Steptoe Pyrg	Pyrgulopsis serrata	No	No	G1			EV	3		No
Wet	Freshwater Snails	Sterile Basin Pyrg	Pyrgulopsis sterilis	No	No	G1			EV	3		No
Wet	Freshwater Snails	Lake Valley Pyrg	Pyrgulopsis sublata	No	No	G1			EV	1		No
Wet	Freshwater Snails	Southern Steptoe Pyrg	Pyrgulopsis sulcata	No	No	G1			PS	2		No
Wet	Freshwater Snails	Southern Bonneville Pyrg	Pyrgulopsis transversa	No	Yes	G2				4		No
Wet	Freshwater Snails	Southern Soldier Meadow Pyrg	Pyrgulopsis umbilicata	No	No	G1			HV	5		No
Wet	Freshwater Snails	Northwest Bonneville Pyrg	Pyrgulopsis variegata	No	Yes	G2			EV	10		No
Wet	Freshwater Snails	Duckwater Warm Springs Pyrg	Pyrgulopsis villacampae	No	No	G1			PS	2		No
Wet	Freshwater Snails	Vineyards Pyrg	Pyrgulopsis vinyardi	No	No	G1			EV	2		No
Wet	Freshwater Snails	Wong's Springsnail	Pyrgulopsis wongi	No	No	G2		AZ	MV	49		No
Wet	Freshwater Snails	Fat-whorled Pondsnaail	Stagnicola bonnevillensis	No	Yes	G1				5		No
Wet	Freshwater Snails	Mountain Marshsnail	Stagnicola montanensis	No	No	G3				4		No
Wet	Freshwater Snails	Widelip Pondsnaail	Stagnicola traski	No	No	G3				2		No
Wet	Freshwater Snails	Grated Tryonia	Tryonia clathrata	No	No	G2			PS	3		No
Wet	Freshwater Snails	Grapevine Springs Elongate Tryonia	Tryonia margae	No	No	G1				2		No
Wet	Freshwater Snails	Monitor Tryonia	Tryonia monitorae	No	No	G1			PS	2		No
Wet	Freshwater Snails	Desert Tryonia	Tryonia porrecta	No	No	G3				9		No
Wet	Freshwater Snails	Grapevine Springs Squat Tryonia	Tryonia rowlandsi	No	No	G1				1		No
Wet	Freshwater Snails	Desert Valvata	Valvata utahensis	No	Yes	G2				1		No

<b>Ecoregion Model Group</b>	<b>Taxonomic Group</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Federally Listed</b>	<b>State Protected</b>	<b>Rounded Global Rank</b>	<b>Relevant SWAPs</b>	<b>Relevant BLM Special Status</b>	<b>NatureServe Climate Vulnerability Index</b>	<b># of Natural Heritage Locations</b>	<b>Available GAP Habitat Models</b>	<b>Other Spatial Data</b>
Wet	Mammals	American Beaver	Castor canadensis	No	Yes	G5						No
Wet	Mammals	North American River Otter	Lontra canadensis	No	Yes	G5	NV, UT		PS	19	SW	No
Wet	Mayflies	A Mayfly	Ameletus edmundsi	No	No	G1				2		No
Wet	Mayflies	A Mayfly	Cinygmula gartrelli	No	No	G2				1		No
Wet	Mayflies	A Mayfly	Paraleptophlebia packii	No	No	G2				2		No
Wet	Mayflies	A Mayfly	Parameletus columbiae	No	No	G2				1		No
Wet	Mayflies	A Mayfly	Susperatus tuberculatus	No	No	G1				1		No
Wet	Stoneflies	A Stonefly	Capnia hornigi	No	No	G3						No
Wet	Stoneflies	A Stonefly	Capnia mono	No	No	G2						No
Wet	Stoneflies	Tiny Forestfly	Malenka tina	No	No	G3				1		No
Wet	Stoneflies	Utah Needlefly	Perlomyia utahensis	No	No	G3				16		No
Wet	Stoneflies	Utah Sallfly	Sweltsa gaufini	No	No	G3				4		No
Wet	Turtles	Western Pond Turtle	Actinemys marmorata	No	No	G3	CA	CA		10	SW, CA	No
Wet	Turtles	Northern Pacific Pond Turtle	Actinemys marmorata marmorata	No	No	T3	CA, NV		PS			No
Wet	Flowering Plants	Davis peppercress	Lepidium davisii	No	No	G3		NV				No
Wet	Flowering Plants	Williams combleaf	Polyctenium williamsiae	No	Yes	G2		NV		35		No

**Appendix IV. Management Questions: Implications from Data Evaluation**

Following are management questions forwarded from Task 1. In the last column we identify the relevant data sources and indicate any need for change or possible removal due to inadequate data.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
<b>Species</b>				
What is the current distribution of occupied habitat for each CE, including seasonal habitat, and movement corridors?	Each CE			<b>Terrestrial Coarse Filter CEs:</b> NatureServe map (ReGAP and LANDFIRE EVT); with add'l refinement. <b>Aquatic Coarse Filter CEs:</b> NatureServe map plus NHD Plus, and NWI. <b>Fine-filter CEs:</b> Natural Heritage, FWS, SWAP, and Misc. sources data. <b>Data for Movement Corridors not yet identified.</b>
Where are current CE populations potentially affected by change agents (and potentially at risk)?	Each CE crossed with CAs	All CAs		Criteria for evaluating ecological integrity exist in some form for most Coarse Filter CEs. These finer-grain conceptual models enable us to state assumptions about effects of Change agents. It will be feasible to complete review and refinement of these criteria for subsequent application to spatial modeling.
What is the current distribution of suitable habitat for each CE?	Each CE			The same data sets from the first two questions apply to answer these questions.
Where are change agents potentially affecting this habitat and/or movement corridors?	Each CE crossed with CAs	All CAs		We do NOT yet have all corridor-related data identified.
Where are CEs whose habitats are systematically threatened by CAs (other than climate change)?	Subset of CEs with restricted habitats	All CAs	During Task 3, select CE subset	The same data sets from the first two questions apply to answer these questions.
What areas have been surveyed and what areas have not been surveyed (i.e., data gap locations)?	Each CE			This is a Task 3 activity once species CEs are finalized.
Given current and anticipated future locations of change agents, which habitat areas remain as opportunities for habitat enhancement/restoration?	Subset of CEs		During Task 3, select CE subset or specific habitats.	In addition to the same data sets referenced in the first two questions, SSURGO and LANDFIRE BpS data sets will be useful for this application.
Where are potential areas to restore connectivity?	Selected subset of habitats and locations.		Determine which CEs have connectivity as a relevant concern. Select subset of habitats or locations.	This will be explored and documented as methodology in Task 3. We will answer remaining data input questions at that point.
Where will CEs experience climate outside their current climate envelope?	Each CE	Climate Change	Standard climate envelope analysis	We are reasonably well positioned to address this for major CEs using climate effects models that build on PRISM (4km data) and downscaled future projects (15 km data). Confidence in outputs will vary depending on natural characteristics of CEs and spatial resolution of climate data.
<b>Native Plant Communities</b>				
Where are intact CE vegetative communities located?	All CEs that are vegetative communities			Terrestrial Coarse Filter CEs: NatureServe map (ReGAP and LANDFIRE EVT); with add'l refinement.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where are the locations that most likely include the highest-integrity examples of each major terrestrial ecological system type?	All CEs that are vegetative communities		Develop metric for Integrity that can be applied to CE communities with available data.	Criteria for evaluating ecological integrity provide conceptual model detail. Spatial information to be derived from various landscape condition models and LANDFIRE spatial outputs (raw and refined).
Where will these current communities be potentially affected by Change Agents?	All CEs that are vegetative communities crossed with CAs	All CAs		Data referenced above for current location of all CEs.
Where will current locations of these communities experience significant and abrupt deviations from normal climate variation?	All CEs that are vegetative communities	Climate Change	TBD: Climate models to use and the definition of "significant". This could evolve into a standard climate envelope analysis.	Georeference sample data (from ReGAP & LANDFIRE LFRDB) represent current distributions of types and dominant species for climate envelope models with PRISM data. These then for source material for analysis of future climate envelopes using USGS 15 km data.
<b>Terrestrial Sites of High Biodiversity</b>				
Where are High Biodiversity sites?	Ecoregion-wide		During Task 3, develop a specific working definition of "high biodiversity". For example, is it just species richness, R? Or richness of CEs?	These have been defined as priority sites identified through previous planning efforts. These can be covered adequately with SWAP locations (not yet acquired) TNC ecoregional portfolio sites, and other selected sources.
Where will these High Biodiversity sites be potentially affected by Change Agents?	All High Biodiversity sites (working definition required) crossed with CAs	All CAs		same as above, in combination with CA data.
Where will current locations of these High Biodiversity sites experience significant and abrupt deviations from normal climate variation?	All High Biodiversity sites (working definition required)	Climate Change, potentially other CAs	TBD: Climate models to use and the definition of "significant". This could evolve into a standard climate envelope analysis.	Same as above, with climate effects model outputs (and inherent limitations based on spatial resolution and uncertainty stemming from climate data).
<b>Aquatic Sites of High Biodiversity</b>				
What areas have been (and have not been) surveyed for spring snails and other species of concern?	All aquatic CEs			To be completed in Task 3.
Where are Aquatic High Biodiversity sites?	All Aquatic High Biodiversity sites (working definition required)		During Task 3, develop a specific working definition of "high biodiversity". For example, is it just species richness, R? Or richness of CEs?	These have been defined as priority sites identified through previous planning efforts. These can be covered adequately with SWAP locations (not yet acquired) TNC ecoregional portfolio sites, and other selected sources.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where will these Aquatic High Biodiversity sites be potentially affected by Change Agents?	All Aquatic High Biodiversity sites (working definition required) crossed with CAs	All CAs		Same as above, in combination with CA data
Where will current locations of these Aquatic High Biodiversity sites experience significant and abrupt deviations from normal climate variation?	All Aquatic High Biodiversity sites (working definition required)	Climate Change	TBD: Climate models to use and the definition of "significant". This could evolve into a standard climate envelope analysis.	Same as above, with climate effects model outputs (and inherent limitations based on spatial resolution and uncertainty stemming from climate data).
<b>Specially Designated Areas of Ecological Value</b>				
Where are specially designated areas of ecological value?	Ecoregion-wide		Define subset from the list of CEs or other designated locations.	The 2010 Protected Areas Database provides a foundation for this. Additional selected data sets can fill this out.
<b>Grazing, Wild Horses and Burros</b>				
Where are the current herds of Wild Horses?	Wild horses			These are shown in the BLM herd and herd management area maps
Where are the current herds of Burros?	Burros			Same as above
Where are the current Herd Management Areas (HMAs)?	Wild horses, Burros			Same as above
Which HMAs are exceeding AML?	Wild horses, Burros	Grazing		Additional data on herd numbers and range conditions are required. AMT indicated that this will be very difficult to answer given current uncertainties about the data.
Which current MHA will experience significant effects of Change Agents?	HMAs, Grazing	All CAs		This will be addressed further as change agent datasets are identified and compared against HMAs.
Which current Allotments will experience significant effects of Change Agents?	Allotments, Grazing	All CAs		This will be addressed further as change agent datasets are identified and compared against allotment areas
Which Allotments and HMA will experience climate outside their current climate envelope?	HMAs, Allotments, Grazing	Climate Change, Grazing	Standard climate envelope analysis	This will be addressed further as climate change data is developed and compared against those target areas
<b>Soils</b>				
Where are target soil types within the ecoregion?	Ecoregion-wide		Develop list of relevant soil types.	SSURGO, with gap-filling using STATSGO and 10m DEM-derived landforms. BLM has provided a key to identifying sensitive soils types.
Where will these target soil types be potentially affected by Change Agents?	All target soil types (working definition)	All CAs		same as above, in combination with CA data.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
	required) crossed with CAs			
Surface and Subsurface Water Availability				
Where are current water resources, both natural and man-made?	All surface water bodies		Note: coordinate with a related question in Groundwater Extraction.	NHD, NHDPlus, NID (the latter to help identify artificial impoundments). Data on small man-made sources (stock tanks & wildlife guzzlers) is unavailable.
Of these water resources, which are perennial, ephemeral, etc?	All surface water bodies			NHDPlus. Data on small man-made sources (stock tanks & wildlife guzzlers) is unavailable.
Of these water resources, what is their surface water/groundwater connectivity?	All surface water bodies			Not directly measurable at regional scale; surrogate for streams will be: (a) USGS-SWPA data to identify basin fill aquifers surrounding water bodies; (b) USGS baseflow index data, either organized by grid (bfi48grd) or for NHDPlus (nhd_bfi) to assess the relative contribution of groundwater discharge to coarse-filter aquatic CE stream hydrology. For springs/seeps, we will use the source identified in spring/seep site assessment data if available.
What is the natural range of variation in high and low water levels or flows (e.g., frequency, timing, duration of high and low water levels or flows)?	All surface water bodies			Not directly measurable at regional scale; surrogate will be: (a) catchment runoff estimate from USGS Flint & Flint (2007) data; (b) catchment runoff estimate from the NHDPlus attribute layer for overland flow (nhd_ieof); and (c) baseflow estimation from the NHDPlus attribute layer for USGS Baseflow Index (nhd_bfi) or gridded bfi values (USGS bfi48grd) depending on which we find most easily manipulable
Where are the aquifers and their recharge areas?	All relevant areas			USGS SWPA and Flint & Flint 2007
Where will these water resources be potentially affected by Change Agents?	All surface water bodies crossed with CAs	Many CAs		(see discussion of CAs)
Aquatic Ecological Function and Structure				

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
What is the condition of target aquatic systems? OR What is the condition of target aquatic systems in terms of PFC?	All surface water bodies (may require a subset)	Hydrologic alteration, Invasive species, Development	Many may not have "PFC" defined, especially if they are not riparian. Need to look beyond "function and structure" to look at factors that may contribute to resistance and resilience in the face of disturbances and change agents. This requires a conceptual model: What are the ecological and environmental factors that contribute the most to ecological structure and function, including resistance and resilience in the face of disturbances and change agents? To be developed further during Task 3.	<ul style="list-style-type: none"> <li>• Biotic condition: aquatic bioassessment data from federal and state monitoring programs (EMAP-WSA and other data from Utah State University Western Monitoring Center and Utah State University-BLM National Monitoring Center [aka BLM "Buglab"]); and data on native aquatic species distributions (from Heritage pgms) and aquatic non-native (nuisance) species distributions (see Invasives CA discussion)</li> <li>• Abiotic condition: data on the proportion of annual stream flow resulting from groundwater discharge (baseflow) via USGS bfi datasets (see above); the spatial extent of perennial versus intermittent flow via NHDPlus (see above); the intensity of runoff across associated watershed catchment via Flint &amp; Flint (2007) data and via NHDPlus (nhd_ieof); water quality via USEPA database on USEPA State Impaired Waters data (linked to NHD); the distribution of dams (Army Corps NID); and habitat quality (from Utah State University Western Monitoring Center data and BLM "Buglab" data).</li> <li>• Landscape context: data on near-stream and watershed land use (same as source of Landscape Condition data for terrestrial CEs), water use in the surrounding surface watershed and contributing groundwater zone (from USGS SWPA and state publications), atmospheric deposition of N (a representative potential acidification agent as well as a nutrient) and Hg (a representative potential bioaccumulative pollutant) (from NADP data. To support the analysis of landscape context, we have also identified sources of data with which to identify the basin fill aquifers potentially responsible for sustaining base flow or base water elevations in aquatic CEs, and the watershed zones within each HUC potentially most responsible for generating surface runoff to streams and recharge to basin fill aquifers (USGS SWPA; Flint &amp; Flint 2007 data).</li> </ul>
Where are the degraded aquatic systems (e.g., water quality)?	All surface water bodies	Hydrologic alteration, Invasive species, Development	Requires a working definition of degraded. TBD in a conceptual model.	See notes above on biotic, abiotic condition; landscape context for hydrologic and water quality degradation; see Invasives for the latter.
<b>Fire History</b>				
What areas have experienced significant fire?	Ecoregion-wide	Wildfire (increased and/or decreased frequency)		GeoMac, Fire Perimeters, Fire Occurrence, and Burn Severity data sets
In places that have experience fire, where does the resulting vegetative structure and composition differ from the desired state?	Among locations that have experience significant fire	Wildfire (increased and/or decreased frequency)	Requires, for each location, a definition of what constitutes "desired state". TBD in Task 3.	LANDFIRE FRCC and subsequent spatial model outputs.
<b>Fire Potential</b>				

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where are current areas with high potential for fire?	Ecoregion-wide	Wildfire (increased and/or decreased frequency)	Devise a working definition of "potential for fire". TBD in Task 3.	LANDFIRE FRCC and subsequent spatial model outputs; National Lightning Detection Network.
Where are areas that in the future will have high potential for fire?	Ecoregion-wide	Wildfire (increased and/or decreased frequency)	Devise a working definition of "potential for fire". TBD in Task 3. Based on climate changes and potential changes in vegetation. Coordinate with other relevant MQs.	LANDFIRE FRCC and subsequent spatial model outputs, in combination with Climate Change effects models; severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
<b>Invasive Species</b>				
What is the current distribution of invasive species included as CAs?	Ecoregion-wide	All invasive species CAs		A very diverse selection of datasets are available, most of which are highly localized or state-level. will like require modeling for many species. Aquatics: USGS Nonindigenous Aquatic Species Program, supplemental datasets, supplemental datasets from Montana State University, USGS Ft Collins, Desert Research Institute
What areas are significantly ecologically affected by invasive species?	Ecoregion-wide	All invasive species CAs	Requires a working definition of "significantly ecologically affected". Various definitions are possible (e.g., dominance, alterations of ecological function, in some cases mere presence). AMT should discuss possible definitions.	Conservation element databases and the resulting models, invasive species locations and resulting models. Some existing models will be further reviewed for use. Species may best be approached as ecologically-based groupings.
Where are areas (significantly affected by invasives) that have restoration potential?	Areas identified as significantly affected by invasives.	All invasive species CAs	Requires working definition of "restoration potential. There should be specific definitions for each invasive species under consideration.	Data and model development will reveal areas where restoration is possible however guidance and further development of "restoration potential" is required to target and refine this MQ.
Given current patterns of occurrence and expansion, what is the potential future distribution of invasive species included as CAs?	Ecoregion-wide	All invasive species CAs	Based on climate changes and recent patterns of occurrence and expansion.	Data and model development will suggest where future distribution will take place.
<b>Development</b>				
Where are current locations of relevant development types?	Ecoregion-wide	Development, Transportation and Energy Infrastructure		Spatially explicit datasets of different development types are available for most development CAs. Raster datasets of LU/LC may needed to fill in data gaps.
Where are areas of planned or potential development (outside of current urban areas)(e.g., under lease, plans of operation, governmental planning), including transmission corridors?	Ecoregion-wide	Development, Transportation and Energy Infrastructure	Based on available planning documents.	Some planned development areas are thoroughly documented and available (proposed energy transmission corridors, planned pipelines, etc). Off-the-shelf models (SURGoM, ICLUS) can be customized for ecoregion. Many development plans put forth by private industry will be unavailable unless in NEPA process and recorded by state authorities in a spatially enabled database.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where are the areas of significant ecological change from these anthropogenic activities?	Ecoregion-wide	Development, Transportation and Energy Infrastructure	Based on areas thought to be the targets of development. Develop a working definition of "potential development" that incorporates proximity to existing urban areas, roads, or power lines. Develop a working definition of "significant ecological changed". TBD in Task 3.	Need to clarify several terms, this will likely be answered later in the process. Focus on identifying ecological areas most vulnerable to change and their relative contribution to overall system(s).
Where do locations of current CEs overlap with areas of potential change from anthropogenic activities?	All CEs	Development, Transportation and Energy Infrastructure	Coordinate with Species and other CE-related MQs. This MQ may obviate the MQ "Where are the areas of significant ecological change from these anthropogenic activities?"	Urban growth models can be intersected with CEs to identify locations where resource and development conflicts are likely to occur.
Where are ecological areas with significant recreational use?	Ecoregion-wide	Recreation (land-based, water-based)		Recreation data from BLM is still pending. We have recreational use data for USFS lands.
<b>Groundwater Extraction and Transportation</b>				
Where are aquifers and their recharge zones?	Ecoregion-wide			USGS SWPA, Flint & Flint 2007 and nhd_recharge data; backup datasets include USGS Great Basin 1:1,000,000 aquifer study and USGS-Nevada joint aquifer study (2006)
Where will change agents be more powerful if groundwater is extracted?	Ecoregion-wide	All CAs		(see discussion of CAs)
Where are areas with groundwater resources available to sustain renewable energy projects that would not degrade aquatic ecosystems that also depend on these groundwater resources.	Ecoregion-wide	Hydrologic Alteration, Renewable Energy Development	Coordinate with Renewable Energy MQs	This may be too fine-detailed a question to be answered with a REA, because the groundwater zones contributing to a surface aquatic feature may be quite localized or identifiable only via detailed hydrogeologic field investigations. All we can do is overlay aquatic CE locations with aquifer locations (from USGS SWPA), filtered for aquatic CE occurrences with perennial water (from NHDPlus, including via nhd_bfi) to identify principal aquifers that potentially support perennial water levels/flows in these CE occurrences.
Where are areas under leases of water rights?	Ecoregion-wide		Assume this refers to leases of water rights, or of lands with groundwater rights.	We have not identified a consistent set of data with which to assess the spatial distribution of either surface or ground-water use rights, and will need to clarify with the BLM what they need here. Water use rights are not identified to "areas" and are not "leases," unless a rights holder has leased those rights to another party, in which latter case the lease would be a contract between two private parties and not visible to regulatory agencies. Instead, we will use USGS SWPA (see above) for data on municipal water extraction and agricultural extraction from the principal aquifers, and use various USGS and state reports (publications) to extract more general information on water use and its geography.
Where are the areas showing effects from existing groundwater extraction?	Ecoregion-wide	Hydrologic Alteration	Requires a working definition of "effects".	NWIS for water level declines, but more importantly USGS SWPA, and state water atlas publications for water level declines and ground collapses

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where are artificial water bodies including evaporation ponds, etc.?	Ecoregion-wide		Note: Coordinate with an MQ in Surface Water.	Not sure how we would distinguish "artificial" except as impoundments behind dams (US Army Corps NID). Have located statewide data for evaporation ponds, slurry lagoons for NV only.
Where are the areas with groundwater basins in an overdraft condition?	Ecoregion-wide	Hydrologic Alteration	This is not a question about areas where existing groundwater extraction is having ecological effects (already addressed elsewhere) but a question of where groundwater extraction exceeds the long-term potential for recharge.	This is essentially the same question as the one about "areas showing effects from existing groundwater extraction" with the same answer as above.
Surface Water Consumption and Diversion				
Where are the areas of potential future change in surface water consumption and diversion?	Ecoregion-wide	Hydrologic alteration, Climate change, Development	This should show up in any analysis of where "development" growth is most likely; and in the mapping of where water-intensive energy development is most likely.	This will be an output of the analysis of development/urbanization CA
Where are the areas with surface water resources available to sustain solar power, and other forms of development without degrading aquatic ecosystems that also depend on these groundwater resources?	Ecoregion-wide	Renewable energy development	Coordinate with Renewable Energy MQs. This is an extension of the mapping of where surface waters exist that depend on groundwater levels or discharges for their hydrology, combined with the mapping of development potential.	Question should be about aquatic resources that depend on SURFACE WATER resources. Answer: Since this is the arid west, we can safely assume that every surface water body in CBR is fully appropriated for water rights. In fact, they may potentially be over-appropriated, i.e., some rights can be exercised only during wet years when all other senior rights are fully served. For this reason, we would argue that no surface waters are available for such development without transfer or private lease from an existing rights holder.
Where are the areas showing ecological effects from existing surface water exploitation?	Relevant CEs	Hydrologic alteration, Development	Generate this information by coupling map information on density of surface water use (diversions as well as consumption) from state and USGS reports, with information on degree of degradation of aquatic ecological integrity.	We have to rely on comparisons of historic <u>published</u> records (rather than GIS data) on the distribution of perennial flows and perennial water levels in springs, to records of their distribution today; we have not identified GIS data layers for this purpose.
Where are artificial water bodies including evaporation ponds, etc.?	Ecoregion-wide		Coordinate with an MQ in Surface Water.	We will see what we can get from NHD, but this may simply be too fine-detailed a question for a REA.
Where are the areas with existing surface water extraction that has caused natural aquatic communities to become entirely dry, either seasonally or perennially?	Relevant CEs	Hydrologic alteration, Development	Generate this information by coupling map information on existence of formerly perennial streams with where they don't exist anymore, and overlay information on intensity of upstream and adjacent surface water extraction.	This is essentially the same question as the one about "areas showing effects from existing surface water exploitation" with the same answer as above.
Climate Change: Terrestrial Resource Issues				

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where will changes in climate be greatest relative to normal climate variability?	Ecoregion-wide	Climate Change	Climate change will affect every location, but affect different locations in different ways. So the issue is not where any effects will occur, but where these effects will potentially cause significant ecological change affecting priority conservation elements. Exact climate models are TBD.	Current climate envelopes for CEs based on 4 km PRISM data and change measured through 15 km downscaled data. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Given anticipated climate shifts and the direction shifts in distributions, where are areas of potential habitat fragmentation?	Ecoregion-wide	Climate Change	Fragmentation may be difficult to assess. Consider species-specific responses/perceptions of fragmentation.	Current CA data, project CA data, and Projected CE distribution models. Confidence decreases rapidly with future projections as both spatial resolution gets coarser and confidence in predicted patterns decreases approaching 2060. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Which native plant communities will experience climate completely outside their normal range?	CEs that are plant communities.	Climate Change	Climate envelope studies are complicated by the likelihood that assemblages will not move intact, but shift and reform based on the movements of individual species. This MQ needs further refinement during Task 3 and the analysis. Coordinate with MQ in "Native Plant Communities".	Current climate envelopes for CEs based on 4 km PRISM data and change measured through 15 km downscaled data. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Where will wildlife habitat experience climate completely outside its normal range?	Select relevant wildlife species	Climate Change	Requires a working definition of "wildlife habitat". Coordinate with the "plant communities and climate change MQ".	Current climate envelopes for CEs based on 4 km PRISM data and change measured through 15 km downscaled data. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Where are wildlife species ranges (on the element list) that will experience significant and abrupt deviations from normal climate variation?	Select relevant wildlife species	Climate Change	Consider further reframe as standard climate envelope analysis.	Current climate envelopes for CEs based on 4 km PRISM data and change measured through 15 km downscaled data. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Based on recent distributions and expansion patterns of insect pests and disease, what are expected distributions in the future?	Select relevant pest species	Climate Change, Invasive species	This is a research question that possibly requires speculation beyond the scope of the REA. This MQ remains provisional, and be dropped and listed as a gap in research.	Current climate envelopes for CAs based on 4 km PRISM data and change measured through 15 km downscaled data. Climate Change effects models are severely limited by spatial resolution and uncertainty inherent with use of future climate projections.
Climate Change: Aquatic Resource Issues				

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
Where are aquatic resources that will experience significant and abrupt deviations from normal climate variation?	Ecoregion-wide	Climate Change, Hydrologic alteration	Climate change will affect every location, but affect different locations in different ways. So the issue is not where any effects will occur, but where these effects will potentially cause significant ecological change affecting priority conservation elements.	Does this question refer to aquatic CE occurrences or "resources" for human use, or both? Going by our "Notes" from Memo 1C, my comments are: We will use the Flint & Flint climate-impact data associated with the model they developed for their 2007 USGS publication (USGS Flint & Flint Climate Impact data requested) to assess where and to what extent major changes are forecast for runoff, recharge, and snowmelt patterns. As a backup, we can use NHDPlus attributes from the USGS (nhd_bfi; nhd_ieof; nhd_recharge; nhd_ppt30yr; nhd_tmax30yr; nhd_tmin30yr) to develop a rough empirical model of how runoff and recharge hydrology (the first three of these NHDPlus attribute sets) vary in relation to climate (the last three of these NHDPlus attribute sets). This empirical model would allow us to plug in forecast future climate estimates for the latter three, to produce rough estimates of future conditions for the former three, if we found strong empirical relationships are present. In either case, we won't be able to identify "abrupt" deviations unless we work with large numbers of time steps, and that is unlikely.
Where are aquatic resources that will experience significant and abrupt deviations from normal flow regime or mean water levels?	Ecoregion-wide	Climate Change, Hydrologic alteration	There will potentially include effects on water levels in wetlands and groundwater-driven systems, and changes in riparian inundation patterns. Plus the changes won't be in simple magnitude but may also be in the timing, duration, and frequency of different hydrologic conditions.	Same as above, but linked to identification of which aquifers support baseflow/base water levels in which water bodies (see above). Note, however, that aquifer recharge/discharge is a process taking decades to centuries (or millennia) to unfold, and so the effects of climate change on aquifer discharge rates will take a long time to become evident.
Where will aquatic resources experience significant and abrupt deviations from normal temperature regime?	Ecoregion-wide	Climate Change, Hydrologic alteration	Both "flow" and "hydrologic change will occur. Includes not just "temperature change" but change in the temperature regime.	Same as above vis Flint & Flint projections
Where are aquatic resources that will experience additional effects on physical habitat such as channel morphology due to significant and abrupt deviations in climate and hydrologic regimes?	Ecoregion-wide	Climate Change, Hydrologic alteration		This is a secondary effect of changes in runoff and recharge, per above
<b>Military Constrained Areas</b>				
Where are military constrained areas?	Ecoregion-wide	Military use areas, conflict of use areas, areas of moratoria, potential military expansion,	No official military expansion areas in CBR. Military flight areas will show areas of potential conflict with other development types (wind). Surface disturbance can be shown with LU/LC classifications.	Will address military constraints in terms of alternative energy development, transmission lines and conflicts with flight areas. DoD document to be released in early 2011 will help identify these areas.

Management Questions: Central Basin & Range				
Management Question	Relevant CEs or other unit	Relevant Change Agents	Memo 1C Notes	Data Sources & Recommendations
		DOE contracted areas, installation boundaries		
Where might these areas change in the future?	Ecoregion-wide	Military use areas, conflict of use areas, areas of moratoria, potential military expansion, DOE contracted areas, installation boundaries	Coordinate with various other MQs on climate change and water resources. Consult INRMP of the relevant installations to determine available data and potential presence of CEs and CAs.	Difficult to predict as the armed forces have no official plans to change or expand land use. Suggest removal of this MQ.
Where are areas of possible expansion of military use?	Ecoregion-wide	Potential military expansion	Based on BRAC or other planning documents.	As above.
<b>Bald Eagles, Golden Eagles</b>				
Where are active Bald Eagle nests?	Bald Eagle CE			over 800 locations from Natural Heritage programs
Where are active Golden Eagle nests?	Golden Eagle CE			9 locations from Natural Heritage programs
<b>Atmospheric Deposition</b>				
Where are areas affected by atmospheric deposition of pollutants (nutrient deposition, acid deposition, mercury deposition)?	Ecoregion-wide	Air and Water Quality: Fugitive dust, air pollution, atmospheric deposition	Atmospheric deposition affects ecosystems via both nutrient enrichment and via acid deposition; and affects some individual species through these effects and through mercury deposition. This is a known problem in the higher elevations of the western US.	We will use NADP data on Nitrogen as a stand-in for all air pollutants that involve acid deposition AND result in nutrient enrichment once buffered. We will use NHDPlus nhd_no3 and USGS-Nitrogen Groundwater Risk (gwrisk) data sets as cross-checks on the NADP regional estimates. We will use NADP data on Mercury as a stand-in for all air pollutants that can bio-accumulate and cause physiological or developmental harm.