

Bureau of Land Management, Buffalo Field Office
Smith Creek, Horseshoe Ranch, & Little Bighorn Ranch Allotments Grazing Lease
Renewals
Environmental Assessment (EA), WY-070-EA12-92

1.0 Introduction

PROJECT TITLE: Smith Creek, Horseshoe Ranch, & Little Bighorn Ranch Allotments 10-Year Term Grazing Lease Renewal

LOCATION: **Smith Creek Allotment (32010):** T57N, R87W: Section 29: SW $\frac{1}{4}$, 160 acres of public land.

Horseshoe Ranch Allotment (02327): T56N, R87W: Sec.23: S $\frac{1}{2}$ S $\frac{1}{2}$; Sec.25: S $\frac{1}{2}$ SW $\frac{1}{4}$, Section 36 (entire). 880 total acres of public land.

Little Bighorn Ranch Allotment (12198): T58N, R89W: Sec.20: NW $\frac{1}{4}$ NW $\frac{1}{4}$. 40 acres of public land.

(See attached maps)

PREPARED BY: Charlotte Darling, Biological Science Technician, BLM, BFO

CASEFILE NUMBERS: 4907019; 4907552; 4907323

APPLICANTS: Smith Canyon Ranch Partnership; CR Energy; Little Bighorn Ranch, Inc.

This site-specific Environmental Assessment (EA) tiers to and incorporates by reference the Buffalo Resource Management Plan (RMP) dated October 4, 1985, and the 2001 amendment. This EA follows the format in Chapter 8 of the BLM National Environmental Policy Act Handbook, H-1790-1.

1.1 Background

Smith Canyon Ranch Partnership owns the base property associated with the Smith Creek Allotment, CR Energy owns the base property associated with the Horseshoe Ranch Allotment, and Little Bighorn Ranch, Inc. owns the base property associated with the Little Bighorn Ranch Allotment, and each party currently holds the grazing authorization for its respective allotment. The previous grazing leases expired on February 28, 2009 and February 28, 2010. Smith Canyon Ranch Partnership, CR Energy, and Little Bighorn Ranch, Inc. applied for renewal of the leases authorizing grazing on the Smith Creek, Horseshoe Ranch, and Little Bighorn Ranch Allotments, respectively. Per 43 CFR 4110, Smith Canyon Ranch Partnership, CR Energy, and Little Bighorn Ranch, Inc. have preference in retaining the grazing privileges attached to each property.

These three allotments are being analyzed in a single document due to their proximity to one another, similar vegetative and topographic characteristics, and because they are all located within the Tongue River and Little Bighorn watersheds. BLM is analyzing these allotments and their associated grazing leases on a watershed scale in order to evaluate the effects of the Proposed Action on the wider environment and to better capture cumulative impacts.

1.2 Purpose and Need for the Proposed Action

The BLM promotes healthy sustainable rangeland ecosystems and provides for the sustainability of the western livestock industry and communities that are dependent upon productive, healthy public rangelands while complying with land use plans and multiple use objectives, including environmental and economic values, as provided in 43 CFR 4100, the Taylor Grazing Act of 1934 and the Federal Land Policy and Management Act of 1976. The proposed action would allow livestock grazing on public land through the exercise of grazing preference attached to controlled base property while considering these multiple use objectives (43 CFR 4110).

There is need for the action due to the requirement that an individual or group desiring to graze livestock on public land must hold a valid grazing authorization in the form of a permit or lease; the BLM is to balance the authorization with other uses of public land. The current grazing lessee has a preference to receive the authorization if grazing is to continue on the associated allotment. The BLM issued the current grazing leases in 2009 and 2010 under Public Law 106-291 allowing for authorization of grazing leases until completion of environmental analysis.

1.3 Decision to be Made

The BLM will decide whether or not to issue a grazing lease with no change in terms and conditions to Smith Canyon Ranch Partnership for the Smith Creek Allotment, to CR Energy for the Horseshoe Ranch allotment, and to Little Bighorn Ranch, Inc. for the Little Bighorn Ranch Allotment, and how to balance the proposed action with multiple public uses.

1.4 Conformance with Land Use Plan and Other Laws, Regulations, and Policies

The Proposed Action is in conformance with the Record of Decision for the Buffalo Resource Management Plan approved October 4, 1985, the 2001 and 2011 amendments, and the Powder River Basin Oil & Gas Project Final Environmental Impact Statement and Resource Management Plan Amendment (PRB FEIS) approved April 30, 2003. The action is also consistent with the land use plan terms and conditions as required by 43 CFR 1610.5-3(a). The Buffalo RMP EIS analyzed the impacts of grazing.

This Environmental Analysis fulfills the 1969 National Environmental Policy Act (NEPA) requirement for site-specific analysis. The Proposed Action is in accordance with the following laws and/or regulations, other plans, and is consistent with Federal, State, and local laws, regulations:

- Taylor Grazing Act of June 30, 1934
- Federal Land Policy and Management Act of 1976 (43 U.S.C. 1701 et seq.)
- Public Rangelands Improvement Act of 1978
- Endangered Species Act of 1973
- 43 CFR § 4100 Grazing Administration-Exclusive of Alaska
- Clean Water Act Section 303d
- National Historic Preservation Act of 1966 Section 106
- National Environmental Policy Act of 1969
- Sikes Act of 1969 (Habitat Improvement on Public Land)
- Fish and Wildlife Improvement Act of 1978

- Executive Order 13186 – Responsibilities of Federal Agencies to Protect Migratory Birds
- Grazing Regulations as codified in 43 CFR § 4100 as amended in 2005
- DOI Secretarial Order No.3310—Protecting Wilderness Characteristics on Lands Managed by the BLM, Dec. 2010
- BLM Instruction Memorandum No. WY-2012-019, Greater Sage-Grouse Habitat Management Policy on Wyoming Bureau of Land Management Administered Public Lands including the Federal Mineral Estate

BLM IM WY-2012-019 requires the BLM to address a reasonable range of alternatives in livestock grazing EAs in order to assess the impacts of livestock grazing on Greater Sage-Grouse habitat and land health. The IM also stipulates that a deferred grazing system alternative should be considered if the size of the allotment warrants it. The size and continuity of the public lands in these allotments make a BLM-administered deferred or rest-rotation grazing system an unreasonable alternative in this specific case. In addition, there is no identified sage-grouse habitat present. These allotments are category “C” allotments, meaning their management is custodial in nature. Intensive management of category “C” grazing allotments is at a public cost which largely outweighs public benefit; see generally, *Ted Lapis v. U.S.*, 178 IBLA 62 (2009).

1.4.1 Wyoming Standards for Rangeland Health

Particularly applicable to livestock grazing management by the BLM are the Wyoming Standards for Healthy Rangelands and Guidelines for Livestock Grazing Management. The Secretary of the Interior developed and approved the Standards and Guidelines on August 12, 1997. They address watersheds, ecological condition, water quality and habitat for special status species. These policies and guidelines are critical to achieving ecologically sustainable range management.

The regulation at 43 CFR 4180.1 details four fundamentals of rangeland health. They are:

1. Watersheds are in or are making progress toward properly functioning physical condition, including their upland, riparian-wetland, and aquatic components; soil and plant conditions support water infiltration, soil moisture storage, and the release of water that are in balance with climate and landform and maintain or improve water quality, water quantity, and timing and duration of flow.
2. Ecological processes including the hydrologic cycle, nutrient cycle, and energy flow are maintained, or there is significant progress toward their attainment, in order to support healthy biotic populations and communities.
3. Water quality complies with state water quality standards and achieves, or is making significant progress toward achieving established BLM management objectives such as meeting wildlife needs.
4. Habitats are, or are making significant progress toward, being restored or maintained for Federal threatened and endangered species, Federal Proposed, Category 1 and 2 Federal Proposed Candidate and other special status species.

The BLM developed the Wyoming Standards for Healthy Public Rangelands and Guidelines for Livestock Grazing Management (S&Gs) to achieve the four fundamentals of rangeland health detailed above. These Standards relate the minimal acceptable conditions for BLM administered public rangelands, including the health, productivity, and sustainability of the land. Observing, measuring, and monitoring field conditions of range sites, on a watershed scale, determine whether a Standard is being achieved. In accordance with the grazing regulations, if livestock grazing practices are found to be contributing to a failure to meet a Standard, corrective action is developed and implemented before the next grazing season. Guidelines provide reasonable, responsible, and cost-effective management practices at the grazing allotment and watershed levels to attain and maintain rangeland Standards. These management practices either maintain existing desirable conditions or move rangelands toward statewide Standards within reasonable timeframes.

The six Standards for Healthy Public Rangelands are:

Standard 1: Within the potential of the ecological site (soil type, landform, climate, and geology), soils are stable and allow for water infiltration to provide for optimal plant growth and minimal surface runoff.

Standard 2: Riparian and wetland vegetation have structural, age, and species diversity characteristic of the state of channel success and is resilient and capable of recovering from natural and human disturbance in order to provide forage and cover, capture sediment, dissipate energy, and provide for ground water recharge.

Standard 3: Upland vegetation on each ecological site consists of plant communities appropriate to the site which are resilient, diverse, and able to recover from natural and human disturbance.

Standard 4: Rangelands are capable of sustaining viable populations and a diversity of native plant and animal species appropriate to the habitat. Habitats that support or could support threatened species, endangered species, species of special concern, or sensitive species will be maintained or enhanced.

Standard 5: Water Quality meets state standards.

Standard 6: Air Quality meets state standards.

The Buffalo RMP has been amended to adopt the Wyoming Standards for Healthy Rangelands. An assessment of the S&Gs has not yet been conducted for the Smith Creek Horseshoe Ranch, and Little Bighorn Ranch Allotments. These allotments are “C” category allotments, which are low priority for evaluation (see Section 3.3). In 1998 the BFO developed a schedule for evaluating S&Gs. The allotments on this list are all in the “I” and “M” categories, which are highest priority for management and evaluation as described in the WY S&Gs Implementation Plan. Active management of category “C” isolated public lands is at a public cost and management effort largely beyond the scope of generating public benefit; see generally, *Ted Lapis v. U.S.*, 178 IBLA 62 (2009).

1.5 Scoping and Issues

The BLM conducts its decision-making in accordance with the requirements of the Council on Environmental Quality (CEQ) regulations implementing NEPA, and the Department of Interior (DOI) and BLM policies and procedures implementing NEPA. NEPA and the associated regulatory and policy framework require federal agencies to involve the interested public in their decision-making.

This EA received internal scoping. The identified issues are:

- How would the proposed action affect current livestock grazing management?
- Would the proposed action impact riparian areas?
- Would the proposed action impact invasive species?
- Would and how would the proposed action affect any special status species, particularly sage-grouse (candidate species)?
- Would the proposed action impact big game habitat?
- Would the proposed action impact migratory bird habitats or populations?
- Would the proposed action impact cultural resources and/or lands with wilderness characteristics?

A draft version of this EA is sent to interested parties of record and is posted on the Buffalo Field Office (BFO) website to solicit public and cooperating agency comments over a 30-day period. The BFO uses received comments to assess whether the draft EA covers the issues raised and adequately addresses their significance. The BFO's response consists of either addressing public comments in the decision record or results in the preparation of a final EA.

2.0 PROPOSED ACTION AND ALTERNATIVES

2.1 Alternative A – No Livestock Grazing

Under this alternative the BLM will not permit livestock grazing on the Smith Creek, Horseshoe Ranch, and Little Bighorn Ranch Allotments. The existing grazing leases will be cancelled in accordance with 43 CFR parts 4100 and 1600 to eliminate grazing on the allotments.

2.2 Alternative B- Proposed Action – Issuance of Leases without Modification

The BLM Buffalo Field Office proposes to maintain and improve land health and enhance habitat conditions on public lands in the BFO stewardship area by maintaining and/or enhancing upland grassland health and sagebrush habitats (species composition and structure) and maintaining riparian, wetland, and aquatic habitats through existing livestock grazing management.

The BLM also proposes to issue new a 10-year term grazing lease to Smith Canyon Ranch Partnership for the Smith Creek Allotment, to CR Energy for the Horseshoe Ranch Allotment, and to Little Bighorn Ranch, Inc. for the Little Bighorn Ranch Allotment. There are no modifications to the terms and conditions outlined in the expiring leases. Table 1 lists the details of these BLM grazing leases.

Table 1

Allotment*	Livestock		Season of Use	% PL	AUMs	Type Use
	Number	Kind				
Smith Creek Allotment (32010)	10	Cattle	5/15 – 6/14	100	10	Active
Horseshoe Ranch Allotment (02327)	2	Cattle	3/01—2/28	100	24	Active
Little Bighorn Ranch Allotment (12198)	10	Horse	5/15—9/30	18	8	Active

*BLM recognizes that these allotments consist primarily of non-federal lands. As such, BLM will not limit the season of use or number of livestock as long as grazing use is not to the detriment of the public lands. The lease schedule shown is primarily for billing purposes. The Smith Creek Allotment is primarily administered and managed by the USDA Forest Service Bighorn National Forest.

The proposed action will issue new 10-year term grazing leases to Smith Canyon Ranch Partnership, CR Energy, and Little Bighorn Ranch, Inc. All applicants are currently in good standing with the BLM and meet all mandatory qualifications for obtaining a grazing lease per 43 CFR 4110.1 and 4110.2. In accordance with Title 43 CFR 4130.2(a), “Grazing permits or leases shall be issued to qualified applicants to authorize use on the public lands and other lands under the administration of the Bureau of Land Management that are designated as available for livestock grazing through land use plans.”

The applicants are not proposing any projects or other surface disturbing activities in connection to these lease issuances. The BLM will analyze any future range improvement projects associated with these allotments under separate, site-specific EAs.

3.0 AFFECTED ENVIRONMENT

3.1 Introduction

3.1.1 Location

The Smith Creek (SC) and Horseshoe Ranch (HR) grazing allotments are about 7 miles west and southwest of Dayton, Wyoming in Sheridan County. The Little Bighorn Ranch (LBR) grazing allotment is about twenty miles northwest of Dayton. All three allotments lie along the face of the northern Bighorn Mountains in an area of prairie-mountain transition. The allotments are mixtures of public, private, and state lands (lands administered by the Wyoming Office of State Lands and Investments). Private lands compose the majority of each allotment, with one 40-acre parcel of BLM land in the LBR allotment, one 160 acre parcel of BLM land in the SC allotment, and two parcels of BLM land in the HR allotment. There is legal public access to the public lands in all three allotments via U.S. Forest Service lands.

3.1.2 General Description

The SC, HR, and LBR Allotments are typical of the land forms, soils, and vegetation in the area of influence for northern Bighorn Mountains foothills. Differences in dominant species within the allotment vary with soil type, aspect, topography, and water availability. Annual precipitation and growing season are the principal factors limiting forage production. Floodplains and lowlands with intermittent streams are the most productive sites and the very steep escarpments, ridges, and slopes are the least productive.

The public land in these allotments is clearly lacking in wilderness characteristics due to its small size (less than 5,000 acres) and the presence of mechanically maintained roads.

The soils within the SC, HR, and LBR allotments vary greatly depending on topographic location, slope, elevation, and precipitation. The allotments are at the higher end of the 15-19 inch Northern Plains precipitation zone. The majority of soils on the public lands in these allotments are loams and coarse uplands.

Vegetation in these three allotments varies substantially across the elevation gradient. On the BLM lands, which are at higher elevations in the allotments (around 6,500 feet), the vegetative community is dominated by ponderosa pine, lodgepole pine, limber pine and Douglas fir. The understory consists primarily of creeping juniper, Idaho fescue (*Festuca idahoensis*), green needlegrass (*Nasella viridula*), and rhizomatous wheatgrasses.

At lower elevations of the allotments, on private lands, the plant community is dominated by mixed grasses and forbs, along with a shrub component of Wyoming big sagebrush and shrubby cinquefoil. Grass species in this community include rhizomatous wheatgrasses, prairie junegass, and other cool-season bunchgrasses.

With elevations above 5,000 feet, the growing season in these allotments is short, consisting of the months of April through mid-August.

Historically, native plants in northeastern Wyoming evolved under prehistoric conditions which included grazing and browsing by bison and other native ungulates, and an associated low frequency of fire. This community is well suited to grazing by both domestic livestock and wildlife year round.

3.1.3 Energy Development

The BLM permits federal mineral development (coal bed natural gas, conventional oil, and coal) in the PRB. This includes federal minerals below federal and/or private (split estate) surface. The BLM prepares EAs, as required by NEPA, for this federal mineral development. In general, companies submit proposals in the form of Plans of Development (PODs) that may consist of one to 200 wells. Currently the SC, JR, and LBR allotments do not lie within any mineral development. Any future development will be analyzed in a separate EA.

This grazing lease issuance does not affect the following resources, which receive no further analysis:

Air Quality	Mineral Resources	Visual Resource Management
Areas of Critical Environmental Concern (ACEC)	Native American Religious Concerns	Water Quality and Prime or Sole Source of Drinking Water
Environmental Justice	Paleontology	Wild and Scenic Rivers
Prime or Unique Farmlands	Recreation	Wilderness Values
Flood Plains	Soils	
Hazardous or Solid Wastes	Traditional Cultural Properties	

3.2 Cultural Resources

Class III inventory for cultural resources has not occurred on the Horseshoe Ranch allotment; however one not-eligible site is known to exist. Class III inventory for cultural resources has not occurred on the majority of the Little Bighorn Ranch allotment and no known cultural resources are known to exist. Class III inventory for cultural resources has not occurred on the majority of the Smith Creek allotment, although the Wyoming Cultural Records Office database revealed that inventories related primarily to timber harvest have discovered cultural sites. The Smith Creek allotment contains 6 known cultural sites, 4 of which are eligible for the National Register of Historic Places and 2 are not eligible. There may be many more unrecorded cultural sites, some which may be eligible for listing on the National Register, within the allotments.

3.3 Livestock Grazing

In 1985, BLM established three categories for allotments to identify areas where management was potentially needed, as well as to prioritize workloads and the use of range improvement funds. The categories classify allotments as Improve Existing Resource Conditions (I), Maintain Existing Resource Conditions (M), or Custodial Management (C) (USDI 2008). The SC, HR, and LBR Allotments are category “C” allotments, meaning their management is minimal in nature, due to the small amount of public land within the allotments. The BLM’s rationale for this classification is that there are no identified resource problems, and the size and continuity of the public land is not conducive to more intensive management by the BLM. The allotments have a low potential for yielding a positive return on public investment in management or rangeland project development.

The Smith Creek Allotment consists of 160 acres of BLM land, 320 acres of state land, approximately 5760 acres of USDA Forest Service land, and about 90 acres of deeded land. There are 10 AUMs associated with the BLM lands in the allotment. Grazing of public land parcels is in conjunction with state, forest service and deeded lands. This allotment is primarily managed and administered by the USDA Forest Service Bighorn National Forest. BLM’s role in management is limited primarily to lease issuance.

The Horseshoe Ranch Allotment consists of 880 acres of BLM land, 120 acres of state land, and 3208 acres of deeded land. There are 24 AUMs associated with the federal lands in the allotment. The Little Bighorn Ranch allotment consists of 40 acres of BLM land and 147 acres of deeded land. There are 8 AUMs associated with the BLM land in the allotments. In both allotments, grazing of public land parcels is in conjunction with state and private lands.

3.4 Invasive Species/Noxious Weeds

Invasive species and noxious weeds exist in the affected environment. The primary species in the allotment are downy brome (*Bromus tectorum*) and Japanese brome (*Bromus Japonicus*). These *Bromus* species occur in such high densities and numerous locations throughout Northeast Wyoming that a control program is not considered feasible at this time.

3.5 Wetlands and Riparian Zones

The South Fork of the Little Tongue River flows through public land on the northwestern corner of the Horseshoe Ranch Allotment. The river flows for about 0.25 miles on BLM land. There is also a small perennial stream located on public land in this allotment. Both of these waterways

are located in steep, forested terrain, which livestock do not frequently enter. Due to the short stream length on public land and fragmented land ownership downstream of the allotment, these waterways are not priorities for more intensive BLM monitoring or management.

All other stream channels in the Smith Creek, Horseshoe Ranch, and Little Bighorn Ranch allotments are intermittent streams. This means that water flow generally occurs during the wet season (50% of the year or less) so water typically only flows in these channels during times such as spring runoff. Water ceases to flow in these channels during drier periods but may still continue to run underground. As such, there may or may not be riparian vegetation associated with intermittent stream channels. Also, they are not a reliable source of water for livestock or wildlife.

3.6 Wildlife, Migratory Birds, Threatened & Endangered, Candidate and Sensitive Species

The BLM conducted wildlife evaluations to assess the occurrence of selected wildlife species and their habitats, as well as to evaluate the anticipated effects associated with issuance of these grazing leases on the SC, HR, and LBR Allotments. The evaluations included selected individual species or species groupings that are ecologically, economically, or socially important.

Evaluation methods included comparison of aerial imagery (1994 to 2009) and review of wildlife geospatial datasets (available at BFO). Datasets included occurrence information for big game, raptors, bald eagles, sage-grouse, sharp-tailed grouse, mountain plover, black-tailed prairie dogs, and sagebrush in the project area.

Wildlife habitats occurring on the SC, HR, and LBR Allotments are results of a complex history of natural and man-caused influences. Important natural influences included short- and long-term climate variation, infrequent wildfire, and ungulate grazing; especially by bison (Baker 2006; Mack and Thompson 1982). From about 1880 to 1910, the removal of native bison, and their subsequent replacement with “vast numbers” of cattle and excessive numbers of sheep, greatly influenced the PRB, including these allotments (Cassity 2007; Patterson 1952). The combined impacts of cattle and sheep overstocking and climate may have initiated the ongoing epicycle of gully erosion that is evident throughout the Basin (Leopold and Miller 1954). Enactment of the Taylor Grazing Act of 1934 repaired early range degradation and aided the recoveries of reduced wildlife populations (Patterson 1952).

The following tables summarize the affected environment relative to selected wildlife.

Table 2. Summary of Species Habitat and Project Effects.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Amphibians				
Columbia Spotted frog (<i>Rana pretiosa</i>)	Breeds in the shallows of lakes, ponds, marshes, and small streams (NatureServe).	S	MIIH	Habitat may be present. Individuals or eggs may be trampled.
Northern leopard frog (<i>Rana pipiens</i>)	Beaver ponds, permanent water in plains and foothills (SS Policy). Swampy, cattail marshes on the plains (WGFD CWCS).	S	MIIH	Habitat may be present. Individuals or eggs may be trampled.
Birds				
Baird's sparrow (<i>Ammodramus bairdii</i>)	Grasslands, weedy fields (SS Policy). Un- or lightly grazed mixed-grass prairie, wet meadows, tallgrass prairie. Prairie w/ scattered low bushes and matted vegetation (NatureServe). In dry years, grassy slough bottoms, alkali flats, and depressions in low lying grasslands.	S	NI	Trampling of nests may occur. Negligible impacts from livestock or humans disrupting breeding, dislodging nests, or causing adult to leave eggs or chicks unattended.
Bald eagle (<i>Haliaeetus leucocephalus</i>)	Mature forest cover often within one mile of large water body (SS Policy). Nests near large lakes and rivers in forested habitat where adequate prey and old, large-diameter cottonwood or conifer trees are available for nesting (WGFD CWCS). Migrating and wintering eagles congregate near open water areas where concentrations of prey are available, such as carcasses of ungulate species, and spawning areas for kokanee, trout, and other fish (WGFD CWCS).	K	NI	Roosting and nesting habitat is present within one mile of the allotments. Bald eagles may use the area for foraging. At least 4 individuals have been observed less than one mile away. Activities associated with ongoing livestock grazing operations are not likely to occur to such an extent that foraging behavior will be disrupted.
Brewer's sparrow (<i>Spizella breweri</i>)	Basin-prairie shrub (SS Policy). Closely associated with sagebrush shrublands that have abundant, scattered shrubs and short grass (WGFD CWCS).	S	MIIH	Trampling of nests may occur. Negligible impacts from livestock or humans disrupting breeding, dislodging nests, or causing adult to leave eggs or chicks unattended.
Burrowing owl (<i>Athene cunicularia</i>)	Grasslands, basin-prairie shrub (SS Policy). Prefers open prairie, grassland, desert, and shrub-steppe habitats, and may also inhabit agricultural areas. It depends on mammals that dig burrows, which it uses for nesting, roosting, and escape (WGFD CWCS).	NP	NI	No prairie dog colonies present.
Ferruginous hawk (<i>Buteo regalis</i>)	Basin-prairie shrub, grasslands, rock outcrops (SS Policy). Semi-arid open country, primarily grasslands, basin-prairie shrublands, and badlands (WGFD CWCS). Requires large tracts of relatively undisturbed rangeland and nests in rock outcrops, the ground, cutbanks, cliff ledges, or trees (WGFD CWCS).	NS	NI	Ferruginous hawks may forage in this area. Livestock activity should not affect foraging behavior.
Loggerhead shrike (<i>Lanius ludovicianus</i>)	Basin-prairie shrub, mountain-foothill shrub (SS Policy). Grasslands interspersed with scattered trees and shrubs that provide nesting and perching sites.	S	MIIH	Ongoing livestock operations will not result in reduced shrub cover or habitat fragmentation. Nests may be toppled by livestock.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Long-billed curlew (<i>Numenius americanus</i>)	Grasslands, plains, foothills, wet meadows (SS Policy). Inhabits a variety of grassland types ranging from moist meadow grasslands to agricultural areas to dry prairie upland, usually near water. Prefers a complex of shortgrass prairies, agricultural fields, wet and dry meadows and prairies, and grazed mixed-grass and scrub communities. Nests on the ground in habitat that includes grass <12", bare ground, shade, abundant invertebrate prey, and a minimum on 40 acres of suitable habitat (WGFD CWCS).	S	MIIH	Suitable habitat may be present. Nests may be trampled or toppled.
Migratory bird species (Various)	Multiple vegetation types are used for breeding, foraging and wintering, with habitat types ranging from grasslands and shrub-steppe to woodlands and riparian areas.	K	MIIH	Trampling of nests may occur. Negligible impacts from livestock or humans disrupting breeding, dislodging nests, or causing adult to leave eggs or chicks unattended. Ongoing livestock operations should not create significant additional impacts.
Mountain plover (<i>Charadrius montanus</i>)	Short-grass prairie with slopes < 5% (SS Policy). Low, open habitats such as arid shortgrass and mixed-grass prairies dominated by blue grama and buffalo grass with scattered clumps of cacti and forbs, and saltbush habitats of the shrub-steppe. Prefers to nest in large, flat grassland expanses with sparse, short vegetation (<=4") and bare ground. Adapted to areas that have been disturbed by prairie dogs, heavy grazing, or fire (WGFD CWCS).	NS	MIIH	A small amount of suitable plover habitat is present on private lands in the Smith Creek and Horseshoe Ranch allotments. Birds may prefer grazed areas.
Northern goshawk (<i>Accipiter gentilis</i>)	Conifer and deciduous forests (SS Policy). Mixed coniferous habitat of a wide variety of ages, structural conditions, and successional stages. Nests in mature stands with multilayered canopies with open understory, small openings, and water within 0.25 miles. Nest stands often on slopes with northerly exposures or in drainages or canyon bottoms protected by such slopes. Post-fledging area is a mosaic of forest types that provide hiding cover and abundant prey. Foraging area may include a variety of forest types and structures but most often consists of forests with a high density of large trees, high canopy closure, high basal area, and relatively open understories, interspersed w/ shrublands and openings with perching trees to observe prey. Winter habitat probably includes a variety of vegetation types, such as forests, woodlands, shrublands, and forested riparian strips (WGFD CWCS).	S	NI	Foraging and nesting habitat is present in both allotments. Ongoing livestock operations should not affect foraging or nesting behavior.
Peregrine falcon (<i>Falco peregrinus</i>)	Cliffs (SS Policy). Forages in open woodlands and forests, shrub-steppe, grasslands, marshes, and riparian habitats. Nests in cliffs that are usually proximate to habitats with abundant prey (WGFD CWCS).	S	NI	Nest substrate is present. Ongoing livestock operations should not affect falcons if they are present.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Plains Sharp-Tailed Grouse (<i>Tympanuchus phasianellus jamesi</i>)	Short and mixed-grass prairie, sagebrush shrublands, woodland edges, and river canyons. Common where grasslands are intermixed with other shrublands, especially wooded draws, shrubby riparian area, and wet meadows. Diets include a variety of forbs, grasses and insects. In winter, sharp-tailed grouse also feed on buds and catkins of deciduous trees or shrubs and berries. Birds are also known to feed on the buds of aspen and willow.	NS	MIIH	Properly managed grazing will maintain quality cover and habitat. Nests or chicks may occasionally be trampled. There are several leks within 4 miles of the Smith Creek allotment. Ongoing livestock operations are not likely to change use of this area by Sharp-tailed grouse.
Sage sparrow (<i>Amphispiza billineata</i>)	Basin-prairie shrub, mountain-foothill shrub (SS Policy). Considered a sagebrush obligate. Inhabits prairie and foothills shrubland habitat where sagebrush is present. Prefers shrublands with tall shrubs and low grass cover, where sagebrush is clumped in a patchy landscape. Requires a large block of unfragmented habitat to successfully breed and survive (WGFD CWCS).	S	MIIH	Nests may be trampled. Cover will be affected.
Sage thrasher (<i>Oreoscoptes montanus</i>)	Basin-prairie shrub, mountain-foothill shrub (SS Policy). Considered a sagebrush obligate. Inhabits prairie and foothills shrubland habitat where sagebrush is present. Prefers shrublands with tall shrubs and low grass cover, where sagebrush is clumped in a patchy landscape (WGFD CWCS).	S	MIIH	Nests may be trampled. Uncommon cowbird host, which are associated with cattle. May be more susceptible to higher parasitism pressure.
Trumpeter swan (<i>Cygnus buccinator</i>)	Lakes, ponds, rivers (SS Policy). Inhabits shallow marshes, ponds, lakes, and river oxbows. Prefers stable, quiet, and shallow waters where small islands, muskrat houses, or dense emergent vegetation provide nesting and loafing sites. Nutrient-rich water, with dense aquatic plant and invertebrate growth, provide the most suitable habitat. Winter habitat must provide extensive beds of aquatic plants that remain ice-free. In Wyoming, cold temps and ice restrict trumpeters to sites where geothermal waters, springs, or outflow from dams maintain ice-free areas (WGFD CWCS).	NP	NI	Habitat not present.
White-faced ibis (<i>Plegadis chihi</i>)	Marshes, wet meadows (SS Policy). Inhabits marshes, wet-moist meadows, lakes, and irrigated meadows. Nests on the ground in bulrushes, cattails, or reeds; on a floating mat; or in a low tree.	NS	MIIH	Nests and eggs may be trampled.
Yellow-billed cuckoo (<i>Coccyzus americanus</i>)	Open woodlands, streamside willow and alder groves (SS Policy). Nests primarily in large stands of cottonwood-riparian habitat below 7000 feet, including such habitats that occur in urban areas. It is a riparian obligate species that prefers extensive areas of dense thickets and mature deciduous forests near water, and requires low, dense, shrubby vegetation for nest sites.	S	MIIH	Suitable habitat may be present. Ongoing livestock operations should not create significant additional impacts. Negligible impacts from livestock or humans disrupting breeding, dislodging nests, or causing adult to leave eggs or chicks unattended.
Fish				
Yellowstone cutthroat trout (<i>Oncorhynchus clarki bouvieri</i>)	Mountain streams and rivers in Tongue River drainage	S	MIIH	Suitable habitat is found in the allotments. Presence of individuals is unknown. Livestock operations should not have population-level effects.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Mammals				
Black-tailed prairie dog (<i>Cynomys ludovicianus</i>)	Prairie habitats with deep, firm soils and slopes less than 10 degrees (SS Policy). Inhabits dry, flat, open, shortgrass and mixed-grass grasslands with low, relatively sparse vegetation, including areas overgrazed by cattle. Constructs burrows in fine to medium soils (WGFD CWCS).	NP	NI	No prairie dog colonies present.
Fringed myotis (<i>Myotis thysanodes</i>)	Conifer forests, woodland chaparral, caves and mines (SS Policy). Found in a wide range of habitats, including coniferous forests, woodlands, grasslands, and shrublands. Probably most common in xeric woodlands, such as juniper, ponderosa pine, and Douglas-fir. Typically forages over water, along forest edges, or within forests and woodlands. During summer, uses a variety of roosts, including rock crevices, tree cavities, caves, abandoned mines, and buildings. During winter, it hibernates in caves, abandoned mines, and buildings (WGFD CWCS). Must remain within commuting distance of drinking water. Roosts in rock crevices that typically face southeast or southwest and are in low elevation forests or woodlands (WGFD Bat Conservation Plan).	S	NI	Roosting and foraging habitat is present. Ongoing livestock operations are not likely to affect.
Long-eared myotis (<i>Myotis evotis</i>)	Conifer and deciduous forest, caves and mines (SS Policy). Primarily inhabits coniferous forest and woodland, including juniper, ponderosa pine, and spruce-fir. Typically forages over rivers, streams, and ponds within the forest-woodland environment. During summer, it roosts in a wide variety of structures, including cavities in snags, under loose bark, stumps, buildings, rock crevices, caves, and abandoned mines. During winter, it probably hibernates primarily in caves and abandoned mines (WGFD CWCS). Occasionally found in cottonwood riparian areas, basins, and sagebrush grasslands where roost sites are available (WGFD Bat Conservation Plan). Most likely found in areas close to a water source. May also occur more frequently in suitable habitat near rock outcroppings or cliffs. Primarily forages over rivers, streams, and ponds within the forest-woodland environment. Also forages over open areas such as campgrounds, small forest openings, and edges, although foraging areas are most likely to be close to a water source. Large-diameter conifer snags provide primary roosting habitat (WGFD Bat Conservation Plan).	S	NI	Roosting and foraging habitat is available. Ongoing livestock operations are not likely to affect.
Spotted bat (<i>Euderma maculatum</i>)	Cliffs over perennial water (SS Policy). Occupies a wide variety of habitats, from desert scrub to coniferous forest. Most often observed in low deserts and basins and juniper woodlands. Roosts in cracks and crevices in high cliffs and canyons. May occasionally roost in buildings, caves, or abandoned mines, although cliffs are the only roosting habitat in which reproductive females have been located (WGFD CWCS). Often occurs in association with canyons, prominent rock features, and	S	NI	Cliffs and perennial water sources are available. Ongoing livestock operations should not affect bats.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
	permanent water sources. In desert environments, it forages in canyons, in the open, or over riparian vegetation. All recorded occurrences of spotted bats in WY were close to a permanent water source (WGFD Bat Conservation Plan).			
Swift fox (<i>Vulpes velox</i>)	Grasslands (SS Policy). Inhabits shortgrass and mixed-grass prairies. Often uses highway and railroad ROWs, agricultural areas, and sagebrush-grasslands. Closely associated w/ prairie dog colonies and uses underground dens year-round. Selects habitat with low-growing vegetation, relatively flat terrain, friable soils, and high den availability (WGFD CWCS).	NS	MIIH	If present, inappropriate grazing could reduce hiding cover and increase susceptibility to predation.
Townsend's big-eared bat (<i>Corynorhinus townsendii</i>)	Caves and mines (SS Policy). Occupies a variety of xeric to mesic habitats, including coniferous forests, juniper woodlands, deciduous forests, basins, and desert shrublands, and is absent only from the most extreme deserts and highest elevations. Requires caves or abandoned mines for roost sites during all seasons and stages of its life cycle, and its distribution is strongly correlated with the availability of these features (WGFD CWCS). May be limited to areas with reliable, accessible sources of drinking water. Forages primarily along forest and woodland edges, riparian corridors, and in open areas near wooded habitat. May avoid open, grazed pasture land.	S	NI	Roost sites are available and foraging habitat is present. Ongoing livestock grazing unlikely to affect prey abundance or availability of foraging habitat.
Plants				
Limber Pine (<i>Pinus flexilis</i>)	High-elevation pine, often marking the tree line either on its own, or with Whitebark Pine (<i>Pinus albicaulis</i>), either of the Bristlecone pines, or Lodgepole Pine (<i>Pinus contorta</i>). Found in steeply-sloping, rocky and windswept terrain in the Rocky Mountains.	K	MIIH	Limber pine may be present in association with conifer species. Livestock may forage on young seedlings.
Porter's sagebrush (<i>Artemisia porterii</i>)	Sparsely vegetated badlands of ashly or tufaceous mudstone and clay slopes 5300-6500 ft.	NP	NI	Habitat not present
William's wafer parsnip (<i>Cymopterus williamsii</i>)	Open ridgetops and upper slopes with exposed limestone outcrops or rockslides, 6000-8300 ft.	NP	NI	Habitat not present
<p>Presence K - Known, documented observation within project area. S - Habitat suitable and species suspected, to occur within the project area. NS - Habitat suitable but species is not suspected to occur within the project area. NP - Habitat not present and species unlikely to occur within the project area.</p> <p>Project Effects NI - No Impact. MIIH - May Impact Individuals or Habitat, but will not likely contribute to a trend towards Federal listing or a loss of viability to the population or species. WIPV - Will Impact Individuals or Habitat with a consequence that the action may contribute to a trend towards Federal listing or cause a loss of viability to the population or species. BI - Beneficial Impact</p>				

Table 3. Summary of Threatened and Endangered Species Habitat and Project Effects

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Threatened				
Ute ladies'-tresses orchid (<i>Spiranthes diluvialis</i>)	Riparian areas with permanent water. Sub-irrigated meadows.	NP	NE	Habitat not present
Candidates for listing				
Greater sage-grouse (<i>Centrocercus urophasianus</i>)	Basin-prairie shrub, mountain-foothill shrub (SS Policy). Also includes wet-moist meadows, and alfalfa and irrigated meadows when adjacent to sagebrush (WGFDF CWCS).	S	MIIH	There are no leks or modeled habitat in the allotments. The lower elevations on private land may provide marginally suitable summer habitat for grouse. If the area is used for nesting, incubating female, eggs, and/or chicks may occasionally be trampled. Ongoing livestock operations are not likely to change current use of this area by sage-grouse.
Presence K - Known, documented observation within project area. S - Habitat suitable and species suspected, to occur within the project area. NS - Habitat suitable but species is not suspected to occur within the project area. NP - Habitat not present and species unlikely to occur within the project area.			Project Effects LAA - Likely to adversely affect NE - No Effect NLAA - May Affect, not likely to adversely affect individuals or habitat. NLJ - Not likely to jeopardize continued existence MIIH - May impact individuals and habitat NP - Habitat not present and species unlikely to occur within the project area.	

3.6.1 Candidate Species

This environmental assessment discusses Greater sage-grouse (sage-grouse) in detail because they are classified as a Candidate Species, currently warranted for listing under the Endangered Species Act (USFWS 2010) and are thus of heightened management concern in the BFO. Sage-grouse are also a Wyoming BLM sensitive species and a Wyoming Game & Fish Department (WGFD) Species of Greatest Conservation Need (SGCN).

Sage-grouse habitat is not present on BLM lands in the SC, HR, and LBR allotments. Habitat models indicate that BLM lands within the allotments do not contain nesting or winter habitat (Doherty et al. 2007, Doherty 2008). There are no known leks within the allotments, or within 4 miles of the allotments. Private lands at lower elevations in the allotments may provide suitable sage-grouse habitat, especially in the summer months.

3.6.2 Big Game

Big game species occurring within the SC, HR, and LBR allotments include elk, pronghorn, whitetail deer and mule deer. The following table (Table 4) summarizes WGFD big game seasonal range data for the allotments.

Table 4

Big Game Species	Seasonal habitat provided in Smith Creek Allotment	Seasonal habitat provided in Horseshoe Ranch Allotment	Seasonal habitats provided in Little Bighorn Ranch Allotment
<i>Elk</i>	Crucial winter, Spring, Summer, Fall	Crucial winter, Spring, Summer, Fall	Winter
<i>Mule deer</i>	Winter-yearlong, Spring, Summer, Fall	Winter-yearlong, Spring, Summer, Fall	Winter-yearlong, Spring, Summer, Fall

Winter-yearlong use is when a population or a portion of a population of animals makes general use of the documented suitable habitat sites within this range on a year-round basis, but during the winter months there is a significant influx of additional animals into the area from other seasonal ranges. Crucial ranges are seasonal habitats which are determining factors in a population's ability to maintain itself.

The SC and HR allotments provide crucial winter range for elk. This means that this area is used by elk in the winter months, and availability of this habitat is a determining factor in the population's ability to maintain itself.

As of the most recent available report, populations of mule deer in their respective hunt areas are below their WGFD objective (WGFD 2009a).

4.0 ENVIRONMENTAL EFFECTS

This section describes the environmental consequences of the no action alternative (Alternative A), and those of the proposed action, Alternative B. The effects analysis addresses the direct and indirect effects of implementing the proposed action, the cumulative effects of the proposed action combined with reasonably foreseeable Federal and non-federal actions, identifies mitigation measures, and discloses any residual effects.

4.1 Direct and Indirect Effects

4.1.1 Cultural Resources

Alternative A- No Grazing

The absence of grazing will not result in impacts to cultural resources.

Alternative B- Lease Renewal & Issuance

Any activity that removes vegetation or leads to soil erosion can cause impacts to cultural resources. Livestock concentration areas (such as those that form near water sources, supplemental feeding areas, fence corners, etc.) and livestock trail formation may result in impacts to cultural resources. According to the State Protocol Agreement between the Wyoming BLM and the Wyoming SHPO, grazing lease renewals that do not include seasonal grazing changes or changes in livestock types are exempt from case-by-case review. As per Appendix B item #27 and following section IV(A)(3) of the Wyoming State Protocol, on 02/17/12 the Bureau electronically notified the Wyoming State Historic Preservation Office (SHPO) of this grazing lease renewal.

4.1.2 Livestock Grazing

Alternative A- No Grazing

FLPMA requires the BLM to manage public lands and resources according to the principals of multiple use and sustained yield and recognizes the Nation's need for domestic sources of minerals, food, timber, and fiber. FLPMA also requires the BLM—except in cases of emergency—to give two years' notification when an authorization for domestic livestock grazing is cancelled, in whole or in part, to devote the associated lands to another public purpose, including disposal.

The Buffalo RMP states as a resource management decision that *livestock grazing is allowed on all public lands in the resource area except on about 6,000 acres where it has been determined to be incompatible with other resource uses or values.*

There are no fences or natural barriers separating BLM and non-BLM lands. At this time, fencing out the public lands is not practical or cost effective. If extraordinary circumstances arise, such as the identification of an endangered plant or damageable cultural resource on the site, fencing may be a greater priority, and the BLM will address the matter in a separate EA. If the public lands are not leased, and subsequently not fenced, any livestock use occurring thereon is unauthorized. Selecting this alternative will affect how the adjacent private and state lands are grazed because the operator must keep livestock off public lands through herding or fencing, or else be in violation of federal grazing regulations. The mixed ownership pattern in the BFO resource area makes herding difficult, in addition to the fact that herding does not ensure that public lands are not grazed. A rider needs to remain with livestock at all times. Because it is not economically feasible for the BLM to fence all federal land parcels, fences will likely be constructed on private land, fragmenting the area and making BLM unable to stipulate wire spacing to facilitate wildlife movement. Most four-strand fences on private land have a top wire of 46-48 inches with 10-12 inch wire spacing and all wires are barbed. In the absence of fences, the BLM must constantly supervise the public lands to assure they are not being grazed.

No adverse resource impacts resulting from livestock grazing have been identified which would warrant cancellation of all grazing on these allotments. The Buffalo RMP allows for adjustment of forage allocation based on an evaluation of monitoring, field observations, or other data as needed. Additionally, changes in grazing practices can be effective in mitigating impacts without a corresponding reduction in forage allocation.

Alternative B- Lease Renewal & Issuance

Rangeland vegetation inventory (MRB 1957) data indicates an adequate amount of forage is available to support the proposed number of livestock and for wildlife use and the effects of that use within these allotments. The new grazing leases authorize the same numbers and kind of livestock and season of use as the expiring leases. This action is not proposing any changes to grazing management. The BLM does not expect the renewal and issuance of the grazing lease to have any effects on range management.

4.1.3 Invasive Species/Noxious Weeds

Alternative A- No Grazing

Removing livestock grazing from the public land can promote growth—and potential overgrowth—of perennial grasses and forbs, thus crowding out or reducing the potential for invasion of noxious and/or invasive species. However, the overgrowth of vegetation increases the availability of fine fuels, which also increases the risk of wildfire. These fires would also be more intense, allowing opportunistic noxious and invasive species to colonize the public lands. Cooperative weed control efforts could discourage overgrowth of vegetation and decrease the fire return interval.

Alternative B- Lease Renewal & Issuance

Implementing appropriate grazing use, as described in the Proposed Action, along with ongoing cooperative weed control efforts, benefits the health of the native plant community. A healthy native plant community often provides competition against the establishment and/or spread of noxious weeds. Issuing the grazing leases will not result in any additional impacts in relation to the spread of noxious weeds.

4.1.4 Wetlands and Riparian Zones

Alternative A- No Grazing

Removing livestock grazing from the public land in the Smith Creek and Little Bighorn Ranch allotments will have no effect on wetlands and riparian zones, as none are present on the public lands.

Removing grazing from the Horseshoe Ranch allotment should not affect the South Fork Little Tongue River, because topography already restricts most livestock access to the river.

Alternative B- Lease Renewal & Issuance

Implementation of the proposed action would have no effect on wetlands and riparian zones in the Smith Creek and Little Bighorn Ranch allotments, as these areas do not exist on public land in the allotments.

Continuing to allow grazing in the Horseshoe Ranch allotment would have a minimal effect, if any, on the South Fork Little Tongue River, because topography already restricts livestock access to the water. If individual livestock did access the river, no major change to riparian condition is expected, as livestock grazing has historically occurred in this allotment. Due to the short stream length on BLM land and fragmented land ownership downstream of the allotment, these waterways are not priorities for more intensive BLM monitoring or management.

4.1.5 Wildlife, Migratory Birds, Threatened & Endangered, Candidate and Sensitive Species

The U.S. Fish and Wildlife Service has issued a block clearance for the Powder River Basin for the endangered black-footed ferret and threatened blowout penstemon. These species do not occur in the area.

Alternative A- No Grazing

If grazing is removed from the allotment, there will be “no effect” on Ute ladies’-tresses orchid, because there is no suitable habitat for this species. Cancelling grazing may have a negative impact on mountain plover, burrowing owls, and black-tailed prairie dogs by reducing the number of grazed areas, which provide preferred habitat for these species.

Alternative B- Lease Renewal & Issuance

(See tables in Section 3.6)

The proposed action will have “no effect” on Ute ladies’-tresses orchid, as suitable habitat for this species is not present in the allotment. The proposed action is “not likely to jeopardize”—and may benefit—mountain plover, because the birds prefer areas with little vegetative cover (Derner et al. 2009).

4.1.5.1 Candidate Species

Alternative A- No Grazing

There are no known leks or modeled habitat present in the allotments. No impacts to greater sage-grouse are expected to occur.

Alternative B- Lease Renewal & Issuance

There are no known leks and no modeled habitat present in the allotments. No impacts to greater sage-grouse are expected to occur.

4.1.5.2 Big Game

Alternative A- No Grazing

Under the no grazing alternative, winter browse conditions for big game would not improve. Encroaching herbaceous species may ultimately out-compete shrub species, resulting in a reduction in quality of big game winter range (Smith 1949). Additionally, livestock would not remove unpalatable standing dead material, resulting in unimproved forage.

Alternative B- Lease Renewal & Issuance

By managing land to meet Rangeland Health Standards and improving overall rangeland condition, forage for elk and deer will improve.

Forage resources on winter ranges typically limit mule deer populations (Clements and Young 1997). Livestock grazing tends to favor shrubs over grasses, and thus may provide more desirable winter browse conditions on the allotments (Austin and Urness 1996, Austin et al. 1986, Smith 1949).

Livestock grazing may enhance big game forage by reducing unpalatable standing dead material (Short and Knight 2003). Big game and cattle may compete for forage on a minor level. There is very little dietary overlap between cattle and deer during spring and early summer, since cattle feed primarily on grasses while deer select mostly forbs and some grasses. Cattle begin to use more forbs in late summer and fall, potentially increasing competition. Deer increase the amount of shrubs in their diet in fall and winter, thus reducing competition during those seasons (Anderson and McCuiston 2008).

The fences on the allotments pose a hazard to mule deer. In the BFO resource area, fences have caught and trapped deer and pronghorn. Modifying fence in areas used by cattle to a more wildlife “safe” design with height under 48 inches and the bottom wire 16 inches from the ground may reduce this hazard. Most fences in these allotments are located on private land and are not subject to BLM management

Cattle and elk have more similar dietary preferences, and elk have been shown to prefer sites that have already been grazed by cattle. Proper grazing management can improve winter forage conditions for big game (Anderson and Scherzinger 1975). The small amount of grazing occurring in the SC and HR allotments should not negatively impact wintering elk herds. Livestock grazing is most concentrated in the lower elevations of the allotments, outside the boundaries of elk crucial winter range. Livestock grazing has occurred historically on this allotment and the BLM expects no additional impacts to big game from implementation of the proposed action.

4.2 Cumulative Effects

Cumulative effects are those resulting from the incremental impact of an action when added to other past, present, or reasonably foreseeable actions regardless of what agency or person undertakes such other actions. Identified actions include noxious weed control and sage-grouse protection. If future assessments reveal that rangeland health standards are not being met due to livestock grazing, the BLM will address these issues before the start of the next grazing season as required by 43 CFR 4180.

The BLM will continue managing the Smith Creek, Horseshoe Ranch, and Little Bighorn Ranch Allotments to achieve the Wyoming Standards for Rangeland Health. All elements of the environment will benefit from rangelands in good health. The applicants are not proposing any projects or other surface disturbance in connection to these lease renewals, and the terms and conditions of the leases will remain the same. Thus any cumulative impacts resulting from the proposed action should be minor.

4.2.1 Noxious Weeds


Noxious weeds/invasive non-native plants are present within the assessment area to varying degrees. Livestock grazing may benefit certain weeds by reducing competition with grasses but may also help control other species through defoliation. Currently the BFO is addressing the

situation by mapping weed locations and treating them with herbicides or bio-controls in conjunction with the local weed and pest organizations.

4.2.2 Sage-grouse

The sage-grouse population within northeast Wyoming is exhibiting a steady long term downward trend (WGFD 2008a, USFWS 2010). The figure below illustrates a ten-year cycle of periodic highs and lows. Each subsequent population peak is lower than the previous peak. Long-term harvest trends are similar to that of lek attendance (WGFD 2008b). Habitat fragmentation is the primary attributor to these declines (USFWS 2010). No critical sage-grouse habitat has been identified in the allotments affected by the proposed action. There are no identified leks within 8 miles of BLM land in the allotments.

Figure 1



Average peak number of male sage-grouse per active lek and trend line within the BFO 1967-2009

4.3 Mitigation Measures Considered

The terms and conditions included as part of the term grazing leases will mitigate anticipated impacts. No additional mitigation measures are proposed.

4.4 Residual Effects

There are no residual impacts associated with the proposed action.

5.0 Tribes, Individuals, Organizations or Agencies Consulted

Smith Canyon Ranch Partnership
 CR Energy
 Little Bighorn Ranch, Inc.

6.0 List of Preparers

Charlotte Darling, Biological Science Technician

6.1 List of Reviewers, BLM Buffalo Field Office

Name	Title	Responsibility
Kay Medders	Rangeland Management Specialist	Range, Vegetation, Soils
Bill Ostheimer	Wildlife Biologist	Wildlife, Migratory Birds
Seth Lambert	Archaeologist	Cultural Resources
Janelle Gonzales	Rangeland Management Specialist	Invasive Species
Chris Durham	Assistant Field Manager, Resources	Resources
John Kelley	Planning & Environmental Coordinator	NEPA Planning

7.0 References

- Anderson, A. and K.C. McCuiston. 2008. Evaluating strategies for ranching in the 21st Century: Successfully managing rangeland for wildlife and livestock. *Rangelands* 30:8-14.
- Anderson, E.W. and R.J. Scherzinger. 1975. Improving quality of winter forage for elk by cattle grazing. *Journal of Range Management* 28:2, 120-125.
- Austin, D.D. and P.I. Urness. 1998. Vegetal change on a northern Utah foothill range in the absence of livestock grazing between 1948 and 1982. *Great Basin Naturalist* 58:188–191.
- Austin, D.D., P.J. Urness, and R.A. Riggs. 1986. Vegetal change in the absence of livestock grazing, mountain brush zone, Utah. *Journal of Range Management* 39:514–517.
- Baker, W. L. 2006. Fire and restoration of sagebrush ecosystems. *Wildlife Society Bulletin* 34:177-185
- Beck, J.L. and D.L. Mitchell. 2000. Influences of livestock grazing on sage grouse habitat. *Wildl. Soc. Bull.* 28:993–1002.
- Bohne, J., T. Rinkes, and S. Kilpatrick. 2007. Sage-Grouse Habitat Management Guidelines for Wyoming. Wyoming Game and Fish Department. Cheyenne, Wyoming.
- Bork, E.W., N.E. West, and J.W. Walker. 1998. Cover components on long-term seasonal sheep grazing treatments in three-tip sagebrush steppe. *J. Range Manage.* 51:293-300.
- Branson, F. A. 1985. Vegetation changes on western ranges. *The Society for Range Management (Range Monograph Number 2)*. Denver, CO.
- Bureau of Land Management. 2002. Instruction Memorandum No. WY-2001-147, Change 1: Framework Assessment of Sage-grouse Habitat on Public Lands in Wyoming. Bureau of Land Management, Wyoming State Office. Cheyenne, WY.
- Call, M.W. 1979. Habitat requirements and management recommendations for sage grouse. USDI-BLM Denver Serv. Center Tech. Note 330.

Cassity, M. 2007. Stock raising, ranching, and homesteading in the Powder River Basin historic context study. Prepared for the USDI Bureau of Land Management by Historical Research and Photography. Broken Arrow, OK.

Clements, D.C. and J.A. Young. 1997. A viewpoint: rangeland health and mule deer habitat. *Journal of Range Management* 50:129-138.

Derner, J.D., W.K. Lauenroth, P. Stapp, and D.J. Augustine. 2009. Livestock as ecosystem engineers for grassland bird habitat in the Western Great Plains of North America. *J. Rangeland Ecol. & Mgmt.* 62:111-118.

Doherty, K.E. 2008. Population density model provided to BFO BLM as GIS data. Personal Communication.

Fulgham, K.O., M.A. Smith, and J.C. Malechek. 1982. A compatible grazing relationship can exist between domestic sheep and mule deer, p. 458-478. *In: J.M. Peek and P.D. Dalke (eds.) Proc. of the Wildlife-Livestock Relationships Symp. Idaho For., Wildl. and Range Exp. Sta., Univ. Idaho, Moscow, ID.*

Harniss, R.O., and H.A. Wright. 1982. Summer grazing of sagebrush-grass range by sheep. *J. of Range Management* 35:13-17.

Leopold, L. B., and J. P. Miller. 1954. A postglacial chronology for some alluvial valleys in Wyoming. USDI Geological Survey. Geological Survey Water Supply Paper 1261.

Mack, R.N. and J.N. Thompson. 1982. Evolution in steppe with few large, hooved mammals. *American Naturalist* 119: 757-773.

Manier, D.J. and N.T. Hobbs. 2007. Large herbivores in sagebrush steppe ecosystems: Livestock and wild ungulates influence structure and function. *Oecologia.* 152:739-750.

Merritt, S., C. Prosser, K. Sedivec, and D. Bangsund. 2001. Multi-species grazing and leafy spurge. U.S.D.A.-ARS Team Leafy Spurge, Sidney, MT.

Missouri River Basin Investigations (MRB). 1957. Land Planning and Classification Project.

Mosley, J. C. 1996. Prescribed sheep grazing to suppress cheatgrass: A review. *Sheep and Goat Res. J.* 12:74-81.

Olson, B.E. and R.T. Wallander. 2001. Sheep grazing spotted knapweed and Idaho fescue. *J. Range Manage.* 54:25-30.

Patterson, R.L. 1952. The sage grouse in Wyoming. Sage Books, Inc., Denver, CO.

Peters, E.F. and S.C. Bunting. 1994. Fire conditions pre- and post- occurrence of annual grasses on the Snake River Plain, p. 31-36. *In: Proceedings-Ecology and Management of Annual*

Rangelands. Gen. Tech. Rep. INT-GTR 313. USDA Forest Service, Intermountain Research Station, Ogden, UT.

Riggs, R.A. and P.J. Urness. 1989. Effects of goat browsing on Gambel oak communities in northern Utah. *J. Range Manage* 42:354–360.

Sanders, K.D. and A.S. Voth. 1983. Ecological changes of grazed and ungrazed plant communities, p. 176–179. *In*: S.B. Monsen and N. Shaw (eds.) *Managing Intermountain Rangelands - Improvement of Range and Wildlife Habitats*. U.S.D.A. For. Serv., Gen. Tech. Rept. INT-157.

Short, J.J. and J.E. Knight. 2003. Fall grazing affects big game forage on rough fescue grasslands. *J. Range Management* 56:213-217.

Smith, M.A., J.C. Malechek and K.O. Fulgham. 1979. Forage selection by mule deer on winter range grazed by sheep in spring. *J. Range Manage.* 32:40–45.

Smith, A.D. 1949. Effects of mule deer and livestock upon a foothill range in northern Utah. *J. of Wildlife Management* 12:21-23.

State wildlife agencies' ad hoc committee for sage-grouse and oil and gas development. 2008. Using the best available science to coordinate conservation actions that benefit greater sage-grouse across states affected by oil and gas development in Management Zones I-II (Colorado, Montana, North Dakota, South Dakota, Utah and Wyoming). Unpublished report. Colorado Division of Wildlife, Denver; Montana Fish, Wildlife and Parks, Helena; North Dakota Game and Fish Department, Bismarck; Utah Division of Wildlife Resources, Salt Lake City; Wyoming Game and Fish Department, Cheyenne.

Tisdale, E. W. 1994. Great Basin region: sagebrush types. T.N. Shiflet ed. *Rangeland Cover Types*. Society for Range Management. Denver, CO. 40-46.

U.S. Fish and Wildlife Service (USFWS). 2010. 12-Month Findings for Petitions to List the Greater Sage-Grouse (*Centrocercus urophasianus*) as Threatened or Endangered. Denver, CO.


Wambolt, C.L. and G.F. Payne. 1986. An 18-year comparison of control methods for Wyoming big sagebrush in southwestern Montana. *J. Range Manage.* 39:314–319.

West, N.E. 1999. Managing for biodiversity of rangelands, p. 101–126. *In*: W.W. Collins and C.O. Qualset (eds.) *Biodiversity in Agrosystems*. CRC Press, Boca Raton, FL.



WGFD. 2009a. Big Game Job Completion Reports – Sheridan Region. Wyoming Game and Fish Department. Cheyenne, WY.

Wyoming Game and Fish Department (WGFD). 2008. Sheridan Region Wyoming Game and Fish Department: Annual Sage-Grouse Completion Report for 2008. Wyoming Game and Fish Department. Gillette, WY.


**Little Bighorn/Upper Tongue River Renewals:
Smith Creek, Horseshoe Ranch, Little Bighorn Ranch Allotments**



-  Smith Creek Allotment #32010
-  Horseshoe Ranch Allotment #02327
-  Little Bighorn Ranch Allotment #12198
-  Bureau of Land Management
-  Forest Service
-  Private
-  State




2/15/2012 cdarling



1:70,000




2/15/2012 cdarling



- Horseshoe Ranch Allotment #02327
- Bureau of Land Management
- Forest Service
- Private
- State




1:64,000



Location Reference

2/15/2012 cdarling


Little Bighorn Ranch Allotment #12198



- Little Bighorn Ranch Allotment #12198
- Bureau of Land Management
- Forest Service
- Private
- State

N

1:24,000



2/15/2012 cdarling