

**Environmental Assessment (EA) DOI-BLM-WY-070-EA14-350**

## **Lease Renewal**

Banner Allotment #02328  
Schaffer Ranch, LLC  
Lease #4903554

Lawrence, Charles (Nurse Draw)  
Allotment #12190  
Charles K. Lawrence Order Buying  
Company Inc.  
Lease # 4907310

Tipperary Allotment #02494  
Quarter Circle Nine Cattle, Inc.  
Lease # 4907023

Timber Draw Allotment #12199  
Lawrence M. and Teresa L. Grocki  
Lease # 4907566

**Preparer: Kay Medders, Rangeland Management Specialist**

**Bureau of Land Management  
Buffalo Field Office  
1425 Fort Street  
Buffalo, WY 82834**

July, 2014



**TABLE OF CONTENTS**

**1.0 INTRODUCTION.....2**

    1.1 BACKGROUND .....2

    1.2 NEED FOR THE PROPOSAL .....3

    1.3 SCOPING AND ISSUES .....3

**2.0 PROPOSED ACTION (PROPOSAL) AND ALTERNATIVES.....4**

    2.1 ALTERNATIVE I – PROPOSED ACTION/NO ACTION – RENEWAL OF LEASES WITHOUT MODIFICATION .....4

    2.2 ALTERNATIVE II – NO GRAZING ALTERNATIVE .....5

    2.3 ALTERNATIVES CONSIDERED BUT NOT ANALYZED IN DETAIL .....6

        2.3.1 *Greater Sage-Grouse (GSG) Alternative.* .....6

    2.4 CONFORMANCE TO THE LAND USE PLAN, REGULATIONS, AND LAWS .....6

**3.0 AFFECTED ENVIRONMENT .....6**

    3.1 INTRODUCTION .....6

    3.2 LIVESTOCK GRAZING.....7

    3.3 SOILS .....8

    3.4 VEGETATION .....8

    3.5 NOXIOUS WEEDS AND INVASIVE NON NATIVE PLANT SPECIES .....9

    3.6 WATER RESOURCES.....9

    3.7 WILDLIFE.....9

    3.8 CULTURAL AND HISTORIC VALUES .....13

    3.9 SOCIOECONOMICS.....14

**4.0 ENVIRONMENTAL EFFECTS .....14**

    4.1 DIRECT, INDIRECT RESIDUAL EFFECTS, MITIGATION MEASURES, CUMULATIVE EFFECTS .....14

        4.1.1 *Livestock Grazing*.....14

        4.1.2 *Soils*.....15

        4.1.3 *Vegetation*.....16

        4.1.4 *Noxious Weeds and Invasive Non Native Plant Species*.....18

        4.1.5 *Water Resources* .....18

        4.1.6 *Wildlife*.....19

        4.1.7 *Cultural, Historic Values & National Register of Historic Places (NRHP) Eligibility*.....24

        4.1.8 *Socioeconomics*.....24

    4.2 CUMULATIVE EFFECTS SUMMARY .....25

    4.3. MITIGATION/RESIDUAL IMPACTS/MONITORING SUMMARY.....26

**5.0 TRIBES, INDIVIDUALS, ORGANIZATIONS, OR AGENCIES CONSULTED.....26**

**6.0 LIST OF PREPARERS .....26**

    6.1 LIST OF REVIEWERS.....26

**7.0 WORKS CITED AND REFERENCES .....26**

**8.0 ATTACHMENT 1. MAP.....32**

**9.0 APPENDIX 1. TABLES. ....37**

The BLM’s multiple-use mission is to sustain the health and productivity of the public lands for the use and enjoyment of present and future generations. The Bureau accomplishes this by managing such activities as outdoor recreation, livestock grazing, mineral development, and energy production, and by conserving natural, historical, cultural, and other resources on public lands.

## 1.0 INTRODUCTION

### 1.1 Background

The Bureau of Land Management (BLM), Buffalo Field Office (BFO) proposes to renew 10 year grazing leases for the following allotments: Banner (02328), Lawrence, Charles (12190), Tipperary (02497), and Timber Draw (12199). For ease of administration, and in accordance with BLM IM 2005-194, the Lawrence, Charles allotment is being renamed the Nurse Draw allotment. The allotment will be referred to by this name for the remainder of this document. Per the Federal Land Policy and Management Act (FLPMA) Sec. 402 (c) (3) the holder of the expiring lease has priority in receiving the new lease when land use plans allow for livestock grazing and the lessee is in compliance with the terms and conditions of the expiring lease.

The allotments are in close proximity to one another in northern Johnson County and southern Sheridan County, Wyoming, and approximately 15 to 20 miles north and northeast of Buffalo, Wyoming. Elevations range from about 4,000 feet to over 4,900 feet. The allotments consist of about 27,150 total acres of which 12.5% is BLM land, 20.3 % is state land, and 67.2 % is deeded land. The leases authorizing grazing on these allotments include a total of 3,391 acres of federal land and 357 animal unit months (AUMs) of forage. Grazing use is authorized for cattle on three allotments (Banner, Tipperary and Timber Draw allotments) and for sheep and cattle on the Nurse Draw allotment. BLM is analyzing these allotments and their grazing leases on a watershed scale in order to evaluate the effects of the proposed action on the wider environment and to better capture cumulative impacts. The BLM parcels associated with each allotment are listed below and shown in Attachment 1:

- Banner allotment (#02328): T53N., R82W., Sec. 17 SW<sup>1</sup>/<sub>4</sub> SE<sup>1</sup>/<sub>4</sub>, SE<sup>1</sup>/<sub>4</sub> SW<sup>1</sup>/<sub>4</sub>  
Sec. 18 NE<sup>1</sup>/<sub>4</sub> SE<sup>1</sup>/<sub>4</sub>
- Tipperary allotment (#02494): T51N., R79W., Sec. 34 SE<sup>1</sup>/<sub>4</sub> NE<sup>1</sup>/<sub>4</sub>, E<sup>1</sup>/<sub>2</sub>SE<sup>1</sup>/<sub>4</sub>  
Sec. 35 SW<sup>1</sup>/<sub>4</sub>, W<sup>1</sup>/<sub>2</sub>NW<sup>1</sup>/<sub>4</sub>
- Nurse Draw allotment (#12190): T51N., R78W., Sec. 3 lot 1, SE<sup>1</sup>/<sub>4</sub>NE<sup>1</sup>/<sub>4</sub>, NE<sup>1</sup>/<sub>4</sub>SE<sup>1</sup>/<sub>4</sub>  
Sec. 6 lots 1 and 2, S<sup>1</sup>/<sub>2</sub>NE<sup>1</sup>/<sub>4</sub>, SE<sup>1</sup>/<sub>4</sub>  
Sec. 10 NW<sup>1</sup>/<sub>4</sub>SW<sup>1</sup>/<sub>4</sub>, E<sup>1</sup>/<sub>2</sub>SE<sup>1</sup>/<sub>4</sub>  
Sec. 21 All  
Sec. 22 W<sup>1</sup>/<sub>2</sub>NW<sup>1</sup>/<sub>4</sub>  
T52N., R78W., Sec. 27 NW<sup>1</sup>/<sub>4</sub>, N<sup>1</sup>/<sub>2</sub>SW<sup>1</sup>/<sub>4</sub>, SW<sup>1</sup>/<sub>4</sub>SW<sup>1</sup>/<sub>4</sub>,  
NW<sup>1</sup>/<sub>4</sub>SE<sup>1</sup>/<sub>4</sub>  
Sec. 28 S<sup>1</sup>/<sub>2</sub>N<sup>1</sup>/<sub>2</sub>, E<sup>1</sup>/<sub>2</sub>SW<sup>1</sup>/<sub>4</sub>, SE<sup>1</sup>/<sub>4</sub>  
Sec. 31 lots 10 and 11  
Sec. 32 S<sup>1</sup>/<sub>2</sub>NE<sup>1</sup>/<sub>4</sub>, SE<sup>1</sup>/<sub>4</sub>NW<sup>1</sup>/<sub>4</sub>  
Sec. 33 lots 1, 2, and 4, NE<sup>1</sup>/<sub>4</sub>, NE<sup>1</sup>/<sub>4</sub>NW<sup>1</sup>/<sub>4</sub>,  
S<sup>1</sup>/<sub>2</sub>NW<sup>1</sup>/<sub>4</sub>, N<sup>1</sup>/<sub>2</sub>SW<sup>1</sup>/<sub>4</sub>  
Sec. 34 SE<sup>1</sup>/<sub>4</sub>NW<sup>1</sup>/<sub>4</sub>, W<sup>1</sup>/<sub>2</sub>NW<sup>1</sup>/<sub>4</sub>
- Timber Draw allotment (#12199): T49N., R79W., Sec. 6 NE<sup>1</sup>/<sub>4</sub>SW<sup>1</sup>/<sub>4</sub> portion south of I-90  
Right-of-way fence  
Sec. 7 NE<sup>1</sup>/<sub>4</sub>NE<sup>1</sup>/<sub>4</sub> <sup>1</sup>/<sub>4</sub> portion south of I-90

## Right-of-way fence

This EA, WY-070-EA-14-350 analyzes the impacts of the proposed action on the environment in accordance with the National Environmental Policy Act (NEPA). The current grazing lessee owns or controls the base property associated with their allotment and currently holds the grazing authorization for that allotment. Lease #4903554 was last renewed per Section 415, H.R. 2055 (Appropriations Act) on May 1, 2011 and expires on April 30, 2021. Lease #4903554 was last renewed per Section 415, H.R. 2055 (Appropriations Act) on March 1, 2011 and expires on February 28, 2021. Leases #4907566 and #4907023 were last renewed per Section 415, Public Law 111-88 (Appropriations Act) on March 1, 2010 and expire on February 28, 2020.

The current lessees have each applied for renewal of the grazing lease authorizing grazing on their allotment(s). Per 43 CFR 4110, the previous grazing lessees have preference in retaining the grazing privileges attached to each property. If the proposed action is implemented, a new term grazing lease will be offered to each lessee.

The Buffalo Resource Management Plan (RMP) was amended to adopt the *Standards for Healthy Rangelands and Guidelines for Livestock Grazing Management for Public Lands Administered by the Bureau of Land Management in the State of Wyoming* (1997) (S&Gs). A formal assessment of the S&Gs has not yet been conducted for the Banner, Nurse Draw, Tipperary, and Timber Draw allotments. Although no assessments have been completed, the BLM expects that range monitoring and field visits to the allotments would confirm that the allotments are meeting the S&Gs for healthy rangelands in Wyoming. In 1998 the BFO developed a schedule for evaluating S&Gs. The allotments on this list are all in the “I” and “M” categories, which are highest priority for management and evaluation as described in the WY S&Gs Implementation Plan. Active management of category "C" isolated public lands is at a public cost and management effort largely beyond the scope of generating public benefit; see generally, *Ted Lapis v. U.S.*, 178 IBLA 62 (2009).

### **1.2 Need for the Proposal**

BLM’s need for the proposal is to determine whether, how, and under what conditions to support the Buffalo Resource Management Plan’s (RMP) goals, objectives, and management actions (1985, 2001, 2003, and 2011) with allowing livestock grazing on public lands managed by the BLM. Allotment information is an integral part of this EA, which BLM incorporates here by reference. Conditional livestock grazing finds support in the RMP, Taylor Grazing Act, FLPMA, and other laws and regulations.

Decision to be Made: The BLM will decide whether or not to approve the proposed action, and if so, under what terms and conditions agreeing with the BLM’s multiple use mandate, environmental protection, and RMP.

### **1.3 Scoping and Issues**

The BLM conducts its decision-making per the requirements of the Council on Environmental Quality (CEQ) regulations implementing the NEPA, the Department of Interior (DOI), and BLM policies and procedures implementing NEPA. NEPA and the associated regulatory and policy framework require federal agencies use the scoping process in their decision-making. This EA

received internal scoping, from interdisciplinary resource specialists in the BLM Buffalo Field Office. The identified issues are listed below and have been incorporated in Sections 3 and 4 of this EA.

- How would the proposed action affect current livestock grazing management?
- How would the proposed action impact riparian areas/drainages?
- How would the proposed action impact invasive species?
- How would the proposed action impact sensitive soils?
- Would and how would the proposed action affect any special status species, particularly Greater Sage-Grouse (candidate species)?
- How would the proposal impact cultural resources or lands with wilderness characteristics?
- How can grazing impact native vegetation?
- Whether rangeland health assessment has been completed on the allotment

This EA was sent to interested parties of record and is posted on the Buffalo Field Office (BFO) website to solicit public and cooperating agency comments over a 30-day period: <http://www.blm.gov/wy/st/en/info/NEPA/documents/bfo.html>. The BLM received comments to assess whether the EA covers the issues raised and adequately addresses their significance. The BLM's response consists of either addressing public comments in the decision record or results in the preparation of a new EA.

## **2.0 PROPOSED ACTION (PROPOSAL) AND ALTERNATIVES**

### **2.1 Alternative I – Proposed Action/No Action – Renewal of Leases without Modification**

The BLM proposes to maintain and improve land health and enhance habitat conditions on public lands in the BFO stewardship area by maintaining and/or enhancing upland grassland health and sagebrush habitats (species composition and structure) and maintaining riparian, wetland, and aquatic habitats through existing livestock grazing management.

Since no changes are proposed, the Proposed Action Alternative and the No Action Alternative are the same (per BLM IM 2000-022, Change 1 (1999)). The proposed action is to offer a new 10 year term grazing for each of the following allotments: Banner (#02328), Nurse Draw (#12190), Tipperary (#02494 and Timber Draw (#12199. Each lease will have the same terms and conditions as the expiring leases. Decisions will be written separately for each grazing lease. Table 1 shows the current authorized use (mandatory terms and conditions) for each lease.

**Table 1 Mandatory terms and conditions of the grazing leases affected by the proposed action**

Authorization Number	Allotment Number	Allotment Name	Public Acres	% Public Land	Livestock Number	Livestock Kind	Season of Use	AUMs	Type of Use
4903554	02328	Banner	120	2	250	Cattle	05/-09/30	24	Active
4907023	02494	Tipperary	360	10	32	Cattle	03/01-02/28	38	Active
4907566	12199	Timber Draw	74	12	7	Cattle	03/01-02/28	10	Active
4908310	12190	Nurse Draw	2837	9	2000	Sheep	03/01-04/15	55	Active
					2000	Sheep	10/15/-2/28	162	Active
					150	Cattle	03/01-05/16	34	Active
					40	Cattle	05/16-02/28	34	Active
<b>Total</b>			<b>3391</b>				<b>Total</b>	<b>357</b>	

The “other terms and conditions” for each lease are listed below. These ensure the lease conforms to the goals and objectives of the Buffalo RMP Records of Decision (RODs).

- This authorization is subject to cancellation, suspension, or modification for any violation of the regulations at 43 CFR Part 4100, or of the terms and conditions of the authorization
- The terms and conditions of your lease may be modified if additional information indicates that revision is necessary to conform to 43 CFR 4180
- Lessee agrees to allow authorized officers of the USDI-BLM to enter the leased lands at any time for the purpose of inspection
- Please notify BLM if number/kind of livestock or dates of use change

The proposal will issue new 10-year term grazing leases to the grazing lease applicants. The applicants are currently in good standing with the BLM and meet all qualifications for obtaining a grazing lease under 43 CFR 4110.1 and 4110.2. In accordance with 43 CFR 4130.2(a), “Grazing permits or leases shall be issued to qualified applicants to authorize use on the public lands and other lands under the administration of the [BLM] that are designated as available for livestock grazing through land use plans.” During the 10 year term of the lease or following the expiration of the lease, the lease may be modified if information indicates changes in management are needed to ensure the allotments are meeting or progressing towards achieving the S&Gs.

The applicants are not proposing any projects or other surface disturbing activities in connection to these lease issuances. The BLM will analyze any future range improvement projects associated with these allotments under separate, site-specific analysis.

## 2.2 Alternative II – No Grazing Alternative

Under this alternative the BLM will not permit livestock grazing on the Banner (02328), Nurse Draw (12190), Tipperary (02497), and Timber Draw (12199). Alternative II allows the BLM to place a no grazing provision on any or all of the allotments listed in Table 1, singularly or in any combination, in the most efficient, effective legal means. BLM would cancel the existing grazing leases per 43 CFR parts 4100 and 1600 to eliminate grazing on the allotments.

## **2.3 Alternatives Considered but not Analyzed in Detail**

### **2.3.1 Greater Sage-Grouse (GSG) Alternative.**

BLM IM WY-2012-019 (2012) requires the BLM to address a reasonable range of alternatives in livestock grazing EAs in order to assess the impacts of livestock grazing on GSG habitat and land health. The IM also stipulates that a deferred grazing system alternative should be considered if the size of the allotment warrants it. The size, continuity, and management opportunity of the public lands in these allotments make a BLM-administered deferred or rotation grazing system an unreasonable alternative in these specific cases. Portions of Timber Draw, Nurse Draw, and Tipperary allotments are within GSG Core and Connectivity areas. Timber Draw is adjacent to Interstate 25 with approximately 703 acres within GSG Core area, 68 of those acres being federal lands. The management opportunity does not warrant a BLM administered deferred grazing system as there are no existing fences to exclude livestock from the BLM lands nor is there an allotment management plan that defers grazing. However, none of the federal land in the Tipperary allotment is in a Core area and of the 2,838 total federal surface acres in the Nurse Draw allotment only 181 acres is in Connectivity area.

Although portions of the Nurse Draw, Tipperary, and Timber Draw allotments are in GSG Core/Connectivity areas, the management opportunity does not warrant a BLM administered deferred grazing system. In addition, there is limited GSG habitat present in the Banner allotment and it falls outside of important GSG Habitat.

## **2.4 Conformance to the Land Use Plan, Regulations, and Laws**

This proposal does not diverge from the goals and objectives in the Buffalo Resource Management Plan (RMP), 1985, 2001, 2003, 2011, and generally conforms to the terms and conditions of that land use plan, its amendments, and supporting FEISs, 1985, 2003.

## **3.0 AFFECTED ENVIRONMENT**

### **3.1 Introduction**

The Banner (#02328), Nurse Draw (#12190), Tipperary (#02494) and Timber Draw (#12199) allotments are located in Johnson County and are best approached by county roads including Tipperary, Dry Creek, and Murphy Gulch Roads. There is legal public access to small parcels of BLM land in Banner and Timber Draw allotments from the Murphy Gulch and Dry Creek Roads. The allotments are in the Powder River Basin and Pine Scoria Hills level IV ecoregions, which are unglaciated, irregular and dissected plains. Perennial streams in the area are of montane origin with sand, gravel, and cobble substrates. The area's ephemeral or intermittent streams have sandy or silty substrates. The Pine Scoria Hills have rugged, broken land and stone rough hills covered by open ponderosa pine forest or savanna. The allotments lie within the 10"-14" and 15"-19" Northern Plains (NP) precipitation zone, Major Land Resource Area (MLRA) 58B. Mean temperatures in January are 0°F (low) and 36°F (high) and in July they are 52°F (low) and 88°F (high). (Chapman, et al., 2004)

Livestock grazing, wildlife use, and oil and gas production are common area land uses. Recreation, primarily big game hunting, may also occur. The public lands in these allotments are clearly lacking in wilderness characteristics due to their small size (less than 5,000 acres).

**The proposed action does not affect the following resources, which receive no further analysis:**

Air Quality	Mineral Resources	Visual Resource Management
Areas of Critical Environmental Concern (ACEC)	Native American Religious Concerns	Water Quality and Prime or Sole Source of Drinking Water
Environmental Justice	Paleontology	Wetlands and Riparian Zones
Prime or Unique Farmlands	Recreation	Wild and Scenic Rivers
Flood Plains	Soils	Wilderness Values
Hazardous or Solid Wastes	Traditional Cultural Properties	

### **3.2 Livestock Grazing**

In 1985, BLM established three categories for allotments to identify areas where management was potentially needed, as well as to prioritize workloads and the use of range improvement funds. The categories classify allotments as Improve Existing Resource Conditions (I), Maintain Existing Resource Conditions (M), or Custodial Management (C) (USDI 2008). The Banner, Nurse Draw, Tipperary, and Timber Draw allotments are category “C” allotments, meaning their management is minimal in nature, due to the small amount of public land within the allotments. The BLM’s rationale for this classification is that there are no identified resource problems, and the size and continuity of the public land is not conducive to more intensive management by the BLM. The allotments have low potential for yielding a positive return on public investment in management or rangeland project development.

The allotments have been grazed for numerous years. Current livestock grazing season within all allotments is shown in Table 3. The total AUMs available for grazing on public lands within the allotments is 357 AUMs. The allotments consist primarily of private lands. Table 2 describes the current breakdown of land ownership.

**Table 2-Land ownership**

Allotment #	Allotment Name	Surface Ownership*	Acres	Percent
02328	Banner	BLM	120	5%
		Private	2,260	89%
		State	170	6%
		<b>Total</b>	2,550	
12190	Nurse Draw	BLM	2,837	19%
		Private	11,215	73%
		State	1,260	8%
		<b>Total</b>	15,312	
02494	Tipperary	BLM	360	4%
		Private	4,233	49%
		State	4,067	47%
		<b>Total</b>	8,660	
12199	Timber Draw	BLM	74	12%
		Private	566	88%
		State	0	
		<b>Total</b>	640	
		<b>Total (all allotments)</b>	27,162	
		<b>BLM (all allotments)</b>	3,391	12%
		<b>State (all allotments)</b>	5,497	20%
		<b>Private (all allotments)</b>	18,274	68%

\*Note: Data in this table were estimated by BLM and compiled using ArcGIS data, thus acreages on private and state land are approximate.

### 3.3 Soils

Ardisols and Entisols are the most common soils in the allotments. Ardisols are mixed alluvium derived from andesite, limestone, and quartzite. Ardisols are typically well drained with a low runoff classification and an Ardic moisture regime. Entisols are derived from sandy eolian material and have an excessively drained drainage class. They have a slight hazard of erosion and are commonly used as native rangelands.

The principal soils found on public lands consist of the following soil map units:

- 684 – Samday-Shingle-Badlands complex, 10 to 45 percent slopes
- 708/709 -Theedle-Kishona-Shingle loams, 3 to 30 percent slopes
- 639/640 – Forkwood-Cushman loams, 0 to 6 and 6 to 15 percent slopes
- 790/254-Fairburn-Baux-Rock Outcrop complex, 30 to 60 percent slopes

A description of these soils is found in the (Soil Survey Geographic (SSURGO) database for Sheridan County Area, Wyoming , 2011) (Soil Survey Geographic (SSURGO) database for Johnson County Area, Wyoming, Northern Part, 2011) published by the US Department of Agriculture Natural Resources Conservation Service (NRCS).

### 3.4 Vegetation

The principal range sites or ecological sites on BLM land in the allotments are Shallow Clayey, and Loamy. Other range sites or ecological sites found in the allotment include Clay loam. The primary vegetative type throughout the allotments is Wyoming big sagebrush type. Vegetation

found on these sites includes: Wyoming big sagebrush, silver sagebrush, winterfat, rabbitbrush, green needle grass, needle-and-threadgrass, western wheatgrass, bluebunch wheatgrass, prairie Junegrass, Sandberg bluegrass, bluegrama, little bluestem, asters, paintbrushes, clovers, biscuitroot, western yarrow, fringed sagewort, Hoods phlox, buckwheats, and numerous other grasses and forbs. Most of the growth in these vegetation types occurs in May and June. According to the ecological site description for loamy sites (2011), as this site deteriorates species such as blue grama and big sagebrush increase and cool-season grasses such as needlegrass, needle-and-threadgrass, and rhizomatous wheatgrasses will decrease in frequency and production. Annual bromes will commonly increase with improper management as well. Vegetation types such as mixed grass prairie, ponderosa pine intact, and greasewood fans and flats are also present in the allotments. A description of each ecological site may be found on the NRCS Ecological Site Description webpage.

Currently BLM authorizes 357 total AUMs in the allotments. BLM calculated the AUMs using light-to-moderate stocking rates, per the Land Planning and Classification Report of the Public Domain Lands in the Powder and Missouri River Basin (U.S. Department Interior- Bureau of Land Management, 1956).

### **3.5 Noxious Weeds and Invasive Non Native Plant Species**

Invasive species and noxious weeds exist in the affected environment. The primary species in the area are leafy spurge (*Euphorbia esula*), downy brome (*Bromus tectorum*) and to a lesser extent, Japanese brome (*Bromus japonicus*). Downy brome, also referred to as cheatgrass, is present throughout the area but primarily exists along two track trails and other areas of disturbance. Downy brome is an invasive nonnative annual grass that can degrade native plant communities. Presently downy brome is not a major component of the allotments' native plant communities. BLM will aggressively treat current and future noxious weeds which pose a risk to native vegetation on public lands using an integrated pest management (IPM) approach.

### **3.6 Water Resources**

The area is part of the Powder River drainage. The area occupied by the allotments has two principal drainages. Crazy Woman Creek runs through the Nurse Draw, Tipperary, and Timber Draw Allotments and Flying E Creek runs through the Banner Allotment, The creeks are all on private lands. All other drainages on BLM land in the allotments are ephemeral or intermittent. This means that water flow generally occurs during the wet season (50% of the year or less) so water typically only flows in these channels during times such as spring runoff. Water ceases to flow in these channels during drier periods but may still continue to run underground. As such, there may or may not be riparian vegetation associated with intermittent stream channels. Also, they are not a reliable source of water for livestock or wildlife.

### **3.7 Wildlife**

BLM conducted wildlife evaluations including comparison of past and current aerial imagery and a review of wildlife geospatial datasets (available at BFO). They assessed the occurrence of selected wildlife species and their habitats, and evaluated the anticipated effects associated with issuing these grazing leases on the Banner, Nurse Draw, Tipperary, and Timber Draw allotments. The evaluations included selected individual species or species groupings that are ecologically, economically, or socially important. Tables A.1 and A.2 in the appendix summarize the affected environment for selected wildlife.

Wildlife habitats occurring on the allotments are results of a complex history of natural and man-caused influences. Important natural influences included short- and long-term climate variation, infrequent wildfire, and ungulate grazing; especially by bison ( (Baker, 2006), (Mack & Thompson, 1982)). From about 1880 to 1910, the removal of native bison, and their subsequent replacement with “vast numbers” of cattle and excessive numbers of sheep, greatly influenced the PRB, including these four allotments ( (Cassity, 2007); (Patterson, 1952)). The combined impacts of cattle and sheep overstocking and climate may have initiated the ongoing epicycle of gully erosion that is evident throughout the Basin (Leopold & Miller, 1954). Enactment of the Taylor Grazing Act of 1934 repaired early range degradation and aided the recoveries of reduced wildlife populations (Patterson 1952).

Ongoing oil and gas exploration across the Powder River Basin since 2001 has substantially affected the wildlife habitats within both the Tipperary and Nurse Draw allotments.

The plant community and wildlife habitat it supports within the 4 allotments is currently under light to moderate stocking rates and rotational grazing by livestock in the absence of fire or brush control. Big sagebrush is a significant component of this plant community. Cool-season grasses make up the majority of the understory with the balance made up of short warm-season grasses, annual cool-season grass, and miscellaneous forbs.

Ecological sites found within the 4 allotments include Loamy, Clayey, Shallow Clayey and Shallow Sandy. Some of the BLM lands include shale blowouts and rock out-crops or badlands that are not rated as these areas lack suitable soils. The allotments all fall within a precipitation zone of 10-14 inches per year. The NRCS’s State and Transition models are based on the Ecological Site Descriptions (ESD’s). The ESD’s for the Loamy, Clayey and Shallow Clayey ecological sites indicate the state and transition is a mixed sagebrush/grass plant community. The combination of an overstory of sagebrush and an understory of grasses and forbs provides a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for GSG. Brewer’s sparrows nest in big sagebrush plants and a host of other nesting birds utilize stands in the 20-30% cover range. This plant community is resistant to change. A significant reduction of big sagebrush can only be accomplished through fire or brush management. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term.

The Shallow Sandy ESD has a state and transition of threadleaf sedge/fringed sagewort/yucca. This community provides limited foraging for antelope and other grazers due to low production. They may be used as a foraging site by GSG if proximal to woody cover.

### 3.7.1 Threatened and Endangered Species, Special Status (Sensitive) Species and Migratory Birds

Project effects will not impact threatened, endangered, and special status species occurring in the area beyond the level analyzed in the PRB FEIS. A discussion of the affected environment is in the PRB FEIS, pp. 3-174 to 3-179. A description of habitat and presence for threatened and endangered species is in Table A-1, Appendix 1. below. Neither Ute ladies'-tresses orchid (ULT) nor black-footed ferret habitat is present in the project area. These species are not expected to occur.

#### *Migratory Birds*

The PRB FEIS discussed the affected environment for migratory birds, pp. 3-150 to 3-153. Migratory birds migrate for breeding and foraging at some point in the year. The BLM-Fish and Wildlife Service (FWS) Memorandum of Understanding (MOU) (2010) promotes migratory birds' conservation, per Executive Order 13186 (Federal Register V. 66, No. 11). BLM includes migratory birds in every NEPA analysis of actions having potential to affect migratory bird species of concern to fulfill obligations under the Migratory Bird Treaty Act (MBTA). BLM encourages voluntary design features and conservation measures agreeing with those in the programmatic mitigation in Appendix A of the PRB ROD.

Habitats occurring within the 4 allotments include sage-brush steppe grasslands, mixed grass prairie, and mature deciduous trees. Many species that are of high management concern use these areas for breeding habitats (Saab and Rich 1997). Nationally, grassland and shrubland birds declined more consistently than any other ecological association of birds over the last 30 years (WGFD 2009). The FWS's Birds of Conservation Concern (BCC 2008) report identifies species of all migratory nongame birds that, without additional conservation actions, are likely to become candidates for listing under the Endangered Species Act. Species in this list that have the potential to occur in the project area are: Brewer's sparrow, sage thrasher, loggerhead shrike, short-eared owl, and grasshopper sparrow. Of these, 3 species are identified on the BLM Wyoming Sensitive Species list.

The WGFD Wyoming Bird Conservation Plan (Nicholoff 2003) identified 3 groups of Wyoming's high-priority bird species: Level I – those that clearly need conservation action, Level II – species where the focus should be on monitoring, rather than active conservation, and Level III – species that are not of high priority but are of local interest. Species likely occurring in the project area are in Table 3.

**Table 3. Migratory Birds Occurring in Shrub-Steppe Habitat in NE Wyoming (Nicholoff 2003)**

Level	Species	WY BLM Sensitive	Species	WY BLM Sensitive
Level I	Ferruginous hawk	Yes	Sage sparrow	Yes
	Brewer's sparrow	Yes		
Level II	Lark bunting	No	Sage thrasher	Yes
	Lark sparrow	No	Vesper sparrow	No
	Loggerhead shrike	Yes		

Several migratory species are also BLM special status (sensitive) species (SSS). Those known or suspected of occurring in the project area include Brewer’s sparrow and ferruginous hawk.

### 3.7.2 Candidate Species

This EA discusses GSG in detail because they are a candidate species, currently warranted for listing under the Endangered Species Act (U.S. Fish and Wildlife Service(USFWS), 2010)and are of heightened management concern in the BFO. GSG are also a Wyoming BLM sensitive species and a Wyoming Game & Fish Department (WGFD) Species of Greatest Conservation Need (SGCN). GSG habitat is present on BLM lands in the Banner (02328), Nurse Draw (12190), Tipperary (02494) and Timber Draw (12199) allotments.

#### *Greater Sage-Grouse*

Habitat models indicate that BLM lands in all 4 allotments have GSG nesting brood rearing habitat as well as winter habitat for GSG. The Banner allotment has very little of either winter or nesting and brood rearing habitat but the federal lands in the other 3 allotments are predominantly GSG habitat. The GSG lek surveys show 4 leks (Nurse Draw, Kinney I, Kinney II and Kinney III leks) in the Nurse Draw allotment and one lek (Dry Creek II lek) in the Tipperary allotment. The entire Tipperary allotment and the majority of the Nurse Draw allotment (82%) have been enrolled in the NRCS’s Sage-Grouse Initiative program where livestock operational BMPs have been implemented to improve sage brush habitat. There are no GSG leks in the Banner and Timber Draw Allotments.

As noted in BLM WY-IM-2010-012 (2009), domestic livestock grazing has occurred in and around these allotments and “within the range of [GSG] for over 150 years and is the most common and widespread use of rangelands in the western United States. Livestock grazing practices may affect herbaceous composition, cover, and height and has a potential to impact Wyoming Big Sagebrush habitats. WY BLM has standards and guidelines to ensure proper livestock grazing management on public lands which can help maintain healthy rangeland conditions and provide functional habitat for [GSG]. However, poor livestock grazing practices can have long-term negative impacts on [GSG] habitat by degrading sagebrush, meadow, and riparian communities (Bohne, Rinkes, & Kilpatrick, 2007).”

### 3.7.3 Big Game

Big game species occurring in the EA area include pronghorn, white-tailed deer, and mule deer. Table 4 summarizes WGFD big game seasonal range data for the allotments.

**Table 4. Big Game Seasonal habitat provided in each Allotment**

<b>Species</b>	<b>Banner</b>	<b>Tipperary</b>	<b>Timber Draw</b>	<b>Nurse Draw</b>
White-tailed deer	Yearlong	Yearlong	Yearlong	Yearlong
Mule deer	Winter-Yearlong	Yearlong/Winter-Yearlong	Yearlong/Winter-Yearlong	Winter-Yearlong
Pronghorn	Yearlong	Yearlong	Yearlong	Yearlong

Yearlong use is when a population makes general use of suitable documented habitat sites in the range on a year-round basis, but animals may leave the area under severe conditions. Winter-yearlong use is when a population or a portion of a population of animals makes general use of

the documented suitable habitat sites in this range on a year-round basis, but during the winter months there is a significant influx of additional animals into the area from other seasonal ranges.

As of the WGFD's 2012 Job Completion Report, populations of white-tailed deer in Power River Herd Unit (303) are 8% above objectives. Pronghorn in their Ucross Herd Unit (353) are 96% above objectives. Mule deer are 32.1% below their WGFD objective in their Powder River Herd Unit.

### **3.7.4 Raptors**

Raptors use the Banner, Nurse Draw, Tipperary, and Timber Draw allotments for breeding, foraging, wintering, or migration. Common raptor species frequenting the allotments include golden eagle, northern harrier, red-tailed hawk, Swainson's hawk, American kestrel, short-eared, and great-horned owls. Less common species that may use area habitats include bald eagle, rough-legged hawk, and merlin. Bald eagles occasionally roost in cottonwoods in nearby riparian areas in the winter and forage in the area. Raptors prey upon small mammals, reptiles, and fish. Their survival and reproductive success depends on the availability and abundance of these food sources.

There are 56 known raptor nests within the Nurse Draw, Tipperary allotments including golden eagle, bald eagle, red-tailed hawk, long eared owl, great horned owl, burrowing owl, American kestrel and Cooper's hawk. Only 4 of these 56 nests were active when last surveyed in 2013. Although there are no known nests are within the Banner and Timber Draw allotments, the survey efforts in these 2 allotments have been minimal and nesting habitat is present in both areas.

### **3.8 Cultural and Historic Values**

Class III inventory for cultural resources has not occurred on the majority of land within the allotments, although the Wyoming Cultural Records Office database revealed that inventories related primarily to oil and gas development, highway projects, and range improvement projects have discovered cultural sites.

- The Banner allotment contains 27 known cultural sites, 4 of which are eligible for the National Register of Historic Places, 9 are not eligible and 14 are unevaluated.
- The Nurse Draw allotment contains 23 known cultural sites, 2 of which are eligible for the National Register of Historic Places, 18 are not eligible and 3 are unevaluated.
- The Tipperary allotment contains 6 known cultural sites, 5 of which are not eligible for the National Register of Historic Places and 1 is unevaluated.
- The Timber Draw allotment does not contain any known cultural sites.

There may be many more unrecorded cultural sites, some which may be eligible for listing on the National Register, within the allotments.

### **3.9 Socioeconomics**

Ranching is a strong component of local society and has a historical value, as grazing has occurred in northeast Wyoming since the late 1800s. According to the U.S. Department of Agriculture Agricultural Census (U.S. Department of Agriculture, 2010), Wyoming ranked 24<sup>th</sup> in the nation in the value of sale of cattle and calves, and 4<sup>th</sup> for value of sale of sheep and lambs. Within Wyoming, sales of cattle and calves ranked first in market value of agricultural products sold, with sheep and goat sales ranking 5<sup>th</sup>. These statistics show that ranching is a key component in both Wyoming and the nation's agricultural industry. The sale of livestock is linked to the commodity value of public rangelands. Public lands are an essential part of many ranch operations in the Buffalo Field Office, as they are intermingled with and grazed in conjunction with private and state lands. The BLM grazing lease helps maintain the successful functioning of the ranch operation and support the cultural lifestyle of the lessee.

Public land grazing contributes to the State of Wyoming's revenue through "payment in lieu of taxes" by the Federal government. All of the grazing allotments managed by the Buffalo Field Office were established according to provisions of Section 15 of the Taylor Grazing Act. Receipts from grazing on Section 15 lands are distributed as follows: 50% goes to the federal government for range betterment projects, and 50% is returned to the state government. The grazing fee is \$1.35 per animal unit month (AUM) on public land, \$5.13/AUM on Wyoming State Lands, and an average of \$17.60/AUM on private lands. The grazing leases analyzed in this EA generate approximately \$482.00 in federal grazing fees each year.

## **4.0 ENVIRONMENTAL EFFECTS**

### **4.1 Direct, Indirect Residual Effects, Mitigation Measures, Cumulative Effects**

#### **4.1.1 Livestock Grazing**

##### Alternative I-Proposed Action/No Action Alternative

The direct, indirect, and residual impacts associated with livestock grazing include nutrient cycling, physical damage to vegetation, trailing along fences, trampling and heavier grazing use at salted areas. These impacts are likely to continue upon issuing new leases. The proposed action would allow for the grazing lessees to continue grazing on their respective grazing allotments. Livestock would continue to use up to 357 public AUMs annually; see Table 1. Range vegetation inventory (DOI BLM, 1956) data, along with monitoring data from all four allotments in 2013 indicate adequate forage is available in the allotments to support the proposed number of livestock, as well as provide for wildlife use, while withstanding the effects of that use. The new grazing leases authorize the same number and kind of livestock and season of use relative to each BLM parcel as the previous leases. This action is not proposing any changes to grazing management. The BLM does not expect the issuances of the grazing leases to have any effect on range management.

BLM has identified the scope of the proposed action and alternatives as well as the cumulative effects affected area (CEAA) for livestock grazing as the area within the allotment boundaries. BLM anticipates the direct impacts to last for the life of the grazing lease (10 years), while the indirect and long term impacts may persist.

Cumulative Incremental Effect from the Proposal: The incremental loss of forage available for livestock will occur with the addition of grazing to the past, present, and reasonably foreseeable

actions. As long as mitigation and monitoring techniques are implemented to ensure new roads and trails from recreationists and hunters are not made, and fires are suppressed, the loss of vegetation available for livestock should be negligible. Additionally, oil and gas development and rights-of-way may be permitted, thus decreasing the amount of forage available for grazing. However, with best management practices (BMPs) being implemented, their effects should be negligible.

#### Alternative II-No Grazing Alternative

FLPMA requires the BLM to manage public lands and resources by the principles of multiple use and sustained yield and recognizes the Nation's need for domestic sources of minerals, food, timber, and fiber. FLPMA also requires the BLM, except in emergencies, to give 2 years' notification when cancelling, in whole or in part, an authorization for domestic livestock grazing to devote the associated lands to another public purpose, including disposal. The Buffalo RMP resource management decision reads that livestock grazing is allowed on all area BLM lands except on about 6,000 acres where it is incompatible with other resource uses or values.

There are no fences or natural barriers separating BLM and non-BLM lands. At this time, fencing out the public lands is not practical or cost effective. If extraordinary circumstances arise, such as the identification of an endangered plant or damageable cultural resource on the site, fencing may be a greater priority, and the BLM will address the matter in a separate analysis. If the public lands are not leased, and subsequently not fenced, any livestock use occurring thereon is unauthorized. Selecting this alternative will affect how the adjacent private and state lands are grazed because the lessee must keep livestock off public lands through herding or fencing, or else be in violation of federal grazing regulations. The mixed ownership pattern in the BFO resource area makes herding difficult, in addition to the fact that herding does not ensure that public lands are not grazed. A rider needs to remain with livestock at all times. Because it is not economically feasible for the BLM to fence all its land parcels, fences will likely be on private land, fragmenting the area and making BLM unable to stipulate wire spacing to facilitate wildlife movement. Most four-strand fences on private land have a top wire of 46-48 inches with 10-12 inch wire spacing and all wires are barbed. In the absence of fences, the BLM must constantly supervise the public lands to assure they are not grazed.

BLM identified no adverse direct, indirect, or residual impacts resulting from BMP livestock grazing which would warrant cancellation of all grazing on these allotments. The Buffalo RMP allows for adjustment of forage allocation as needed, based on evaluation of monitoring, field observations, or other data. Additionally, changes in grazing practices can be effective in mitigating impacts without a corresponding reduction in forage allocation.

Cumulative Incremental Effect from the No Grazing Alternative: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be less than those expected under the proposed action.

#### **4.1.2 Soils**

##### Alternative I-Proposed Action/No Action Alternative

Grazing can exert both beneficial and detrimental direct, indirect, and residual effects on a soil resource. The main effects that grazing has on the soil resource is removal of aboveground vegetation and hoof action, potentially leading to increased erosion, increased runoff, reduced

infiltration rates and increased bulk density (soil compaction) (Holechek, Pieper, & Herbel, 2004, p. 379). Most of the compaction and erosion will occur where cattle tend to congregate which may include areas along trails, fences and near watering locations. This compaction leads to lowered rates of water infiltration thus leading to high rates of surface runoff and greater soil erosion.

From a positive standpoint, large quantities of dung and urine are deposited in the allotments adding nutrients and organic matter to the soil (McNaughton, 1979). Hoof action benefits the soil resource by improving nutrient cycling by incorporating mulch into soil surface where it can be broken down more quickly by soil organisms (Holechek, Pieper, & Herbel, 2004, p. 379). Livestock grazing can loosen the soil surface during drying periods, remove excess vegetation that may negatively affect net carbohydrate fixation and increase water transpiration rates, and speed up the development of humus in the soil (Holechek, 1981). Because no changes in the current management are being implemented under the proposal, impacts to the soil resource would remain the same and BLM expects no changes from the current state of the resource.

The CEAA for soils is the area inside the grazing allotment boundaries, selected by BLM due to the scope of the proposed action and alternatives. BLM anticipates the direct impacts last for the life of the grazing lease (10 years), while the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposal: The effects of the proposed action, when added to the reasonably foreseeable actions, should be minimal because range health objectives are used in livestock grazing management, hunters and recreationists will be monitored for land abuse, fire suppression will mitigate the severity of fire impacts, and BMPs will be used for new oil, gas and ROW activities. The incremental effects may include soil erosion and soil compaction along new trails made from livestock, roads and trails used by hunting and recreationists, new oil and gas roads, and areas where fires occur. Severity of these impacts would be dependent on the amount of hunter and recreationist use on the allotments, number of oil/gas/ROWS permitted, and the intensity/size of the wildfires.

#### Alternative II-No Grazing Alternative

With the removal of grazing from the allotments, forage would not be removed by livestock. Standing vegetation and litter would increase. The increase in cover may reduce runoff and erosion. With the removal of livestock from the allotment a decrease in compaction and increase infiltration is anticipated (Pluhar, Knight, & Heitschmidt, 1987). The allotment's nutrient cycle would likely change. Cattle increase soil nutrients by depositing excrement on the soil surface. However, with improper management, they may decrease nutrients by consuming and permanently removing plants that put nutrients into the soil system.

Cumulative Incremental Effect from the No Grazing Alternative: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be less than those expected under the proposed action.

### **4.1.3 Vegetation**

#### Alternative I-Proposed Action/No Action Alternative

The direct, indirect, and residual effects grazing has on vegetation vary greatly depending on many factors including but not limited to: resistance to grazing, genetic potential, growth

promoting features, grazing intensity, life stage of plant, and environmental constraints (Holechek, Pieper, & Herbel, 2004, pp. 123-142)). Livestock grazing can have both beneficial and detrimental effects on vegetation depending on the various factors described by Holechek et al. Beneficial impacts may include, but are not limited to: growth stimulation from grazing ruminants saliva (McNaughton, 1979), trampling of seed into the ground (Holechek, 1981), reducing excess accumulation of standing dead vegetation and litter that may chemically and physically inhibit new plant growth (Holechek, 1981), and reducing transpiration losses (Holechek, Baker, Boren, & Galt, 2006). Some detrimental impacts livestock grazing may have on vegetation include, but are not limited to: changes in species composition in upland areas (Brock & Green, 2003), reduced tillering (Belsky, 1986), modified plant growth form caused by consumption of terminal buds, thus promoting lateral branching (Fleischner, 1994), and disruption of ecological succession (Fleischner, 1994).

Under the proposed action, livestock will annually remove approximately 357 AUMs of forage from BLM land in the allotments. Most studies show that light to moderate stocking rates do not compromise rangeland health. BLM authorizes the AUMs based on a light to moderate stocking rate. Therefore, as long as the total number of permitted AUMs consumed does not exceed the allotments' authorized use; the impacts from renewing the grazing leases should not have an undesirable effect on vegetation.

BLM has determined the CEAA for vegetation, noxious weeds, and invasive plants to be the area within the grazing allotment boundaries and the area within one-half mile of those boundaries, in accordance with the scope of the proposed action and alternatives. BLM anticipates the direct impacts to last for the life of the grazing lease, while the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposal: The effects of the proposed action, when added to the reasonably foreseeable actions, should be minimal because rangeland health assessment is used to inform livestock grazing management, hunters and recreationists will be monitored for land abuse, fire suppression will mitigate the severity of fire impacts, and BMPs will be used for new oil, gas and ROW activities. Incremental effects of the proposed action may include forage loss and introduction of non-native species along new trails made by livestock, roads used for hunting and recreation, new oil and gas roads, and in areas where fires occur. The severity of these impacts would depend on the amount of hunter and recreationist use on the allotments, number of oil/gas/ROWs permitted, and the intensity/size of the wildfires.

#### Alternative II-No Grazing Alternative

The no grazing alternative would eliminate both the beneficial and detrimental impacts associated with grazing. It is likely with the removal of grazing that litter would increase, thus increasing fire potential in the allotments. More vegetation would be available for wildlife and ecosystem function. However, Patton et al., (2007) found that production does not increase with the removal of grazing. Other studies found that removal of grazing can lead to an increase in shrub cover and a decrease in species richness and plant diversity (Manier & Hobbs, 2007).

Cumulative Incremental Effect from the No Grazing Alternative: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be reduced compared to those expected under the proposed action.

#### **4.1.4 Noxious Weeds and Invasive Non Native Plant Species**

##### Alternative I-Proposed Action/No Action Alternative

Livestock can transport noxious weeds and invasive nonnative plant species on their coats, feet, and in their digestive tract. Livestock may carry undesirable plants that exist within the allotments or bring them into the allotment from other pastures they have inhabited during their lifetime. Livestock grazing can increase the presence of noxious weeds by over-grazing (DiTomaso, 2000); this is the primary cause of unwanted species invasion (Holechek, Pieper, & Herbel, 2004, p. 508).

Since many roads and trails are present in the allotments, and recreation opportunities exist in the area, new weed introductions are likely to regularly occur. BLM, the county weed and pest agencies, and the grazing lessee monitor these infestations to determine if management changes are needed to control the infestations. Because current and proposed management does not exceed recommended grazing levels and no management concerns occur at this time, BLM anticipates that there will be no increase in noxious weeds or invasive non-native plant species under the proposed action.

##### Alternative II-No Grazing Alternative

Removing livestock grazing from the public land can promote growth and potential overgrowth of perennial grasses and forbs, thus crowding out or reducing the potential for invasion of noxious and/or invasive species. However the overgrowth of vegetation increases the availability of fine fuels, which also increases the wildfire risk. If fires occurred, they would likely be more intense, allowing opportunistic noxious and invasive species to colonize the public lands. Cooperative weed control efforts could discourage vegetation overgrowth and decrease the fire return interval.

[Cumulative effects for this affected resource are addressed in 4.1.3, Vegetation.]

#### **4.1.5 Water Resources**

##### Alternative I-Proposed Action/No Action Alternative

Riparian areas attract livestock due to environmental and nutritional factors and they may use riparian vegetation disproportionately more than adjacent uplands (Gillen, Krueger, & Miller, 1985) (Howery, Provenza, Banner, & Scott, 1996). This attraction can lead to higher use in riparian areas, thus decreasing streambank stability and cover while increasing soil erosion of the uncovered/unstable streambank (McInnis & McLver, 2001), removal of wood vegetation, soil compaction, and reduced water quality (Parsons, Momont, Delcurto, McInnis, & Porath, 2003). Although uncontrolled livestock grazing can result in watershed destruction in certain areas, controlled grazing is not detrimental to water quality and may increase water quantity (Holechek, 1981). No major degradation problems existed under the past and current management of livestock in these allotments. BLM expects direct, indirect, and residual impacts to water resources to remain unchanged.

The CEAA for water resources is the area within the grazing allotment boundaries and areas extending up and downstream from the allotments, as selected by BLM due to the scope of the

proposed action and alternatives. The direct impacts are anticipated to last for the life of the grazing lease (10 years), while the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposed Action: Implementation of the proposed action in combination with any past, present, and reasonably foreseeable actions may increase the possibility for decreased water quality and quantity. This could result from soil erosion into riparian areas. The incremental impacts should be minimal as BLM uses range health objectives in livestock grazing management, and monitors hunters and recreationists for land abuse. Fire suppression will mitigate fire impact severity and BLM uses BMPs for oil, gas, and ROWs.

#### Alternative II-No Grazing Alternative

The removal of grazing would improve and/or maintain riparian health. Use of riparian plants will decrease, thus reducing trampling and hoof shearing along the green line of riparian areas. Total vascular vegetation, shrub, and graminoid canopy cover would increase with the exclusion of livestock (Schulz & Leininger, 1990).

Cumulative Incremental Effect from the No Grazing Alternative: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be less than those expected under the proposed action.

### **4.1.6 Wildlife**

#### **4.1.6.1 Threatened and Endangered Species, Special Status Species and Migratory Birds**

##### Alternative I-Proposed Action/No Action Alternative

(See Tables A.1 and A.2 in Appendix A)

The USFWS issued a block clearance for the PRB for the endangered black-footed ferret. Alternative B would have “*no effect*” on black-footed-ferrets. The proposed action will have “*no effect*” on Ute ladies’-tresses orchid because the species is not expected to occur in the allotments.

##### Alternative II-No Grazing Alternative

The U.S. Fish and Wildlife Service issued a block clearance for the PRB for the endangered black-footed ferret. Alternative A would have “*no effect*” on black-footed-ferrets. If grazing is removed from the allotments, there will be “*no effect*” on Ute ladies’-tresses orchid, because there is no suitable habitat for this species in the allotments.

##### *Special Status (Sensitive) Species and Migratory Birds*

##### Alternative I-Proposed Action/No Action Alternative

Geographic Scope and Timeframe for Migratory Birds: The CEAA is within the Powder River and Tongue River watershed boundary. The 6 allotments fall within the Powder River watershed including all of the BLM administered lands. Many of the species in the watershed are contained therein. Migratory species may travel outside the boundary but most of the life cycle likely occurs in the CEAA. BLM anticipates the direct impacts to last for the life of the grazing lease (10 years), while the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposal on Migratory Birds and Special Status Species: Incremental impacts from the proposal when added to the past, present and reasonably foreseeable actions may result in disruption of species habitat through the loss of vegetation and habitat fragmentation. Loss of vegetation would occur from livestock grazing, new roads (recreation/hunting/oil and gas/ROWs), and wild fire. Habitat fragmentation would result from vertical intrusions associated with development and new roads associated with oil, gas, ROWs, and recreation activities. Additionally, the spread of noxious and invasive weeds from the actions may impact habitat quality by changing the native plant community, plant production, plant diversity, and ecological health. The incremental impacts should be minimal as BLM uses S&Gs in livestock grazing management, monitors hunters and recreationists for land abuse, uses fire suppression will mitigate the severity of its impacts, and uses BMPs for new oil, gas and ROWs.

#### Alternative II-No Grazing Alternative

If grazing is removed from the allotment, there will be “*no effect*” on Special Status Species or Migratory Birds because its habitat is not affected.

Cumulative Incremental Effect from the No Grazing Alternative on Special Status Species and Migratory Birds: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be less than those expected under the proposed action.

#### **4.1.6.2 Candidate Species – Greater Sage-Grouse (GSG)**

##### Alternative I-Proposed Action/No Action Alternative

The proposal’s direct, indirect, and residual effects will impact GSG habitat. Livestock grazing can benefit or degrade GSG habitat, depending on the timing, stocking rate, and habitat affected. Fall grazing may favor upland forb production, and ranchers may use spring grazing to remove herbaceous cover and make forbs more accessible (Smith, Malechek, & Fulgham, 1979), (Fulgham, Smith, & Malechek, 1982). Spring and early summer grazing may help control weeds and remove woody plants, thereby decreasing the risk of wildfire that could remove large areas of habitat (Mosley, 1996), (Olson & Wallander, 2001), (Merritt, Prosser, Sedivec, & Bangsund, 2001), (Riggs & Urness, 1989).

Excessive or poorly managed grazing causes degradation of sagebrush ecosystems and thus GSG habitat (BLM, 2002). Inappropriate grazing management in uplands can reduce perennial grasses and forbs while favoring annual grasses and increasing sagebrush cover (Branson, 1985), (Tisdale, 1994), (Beck & Mitchell, 2000), (Bork, West, & Walker, 1998). This may impact GSG, because they rely on perennial grasses for escape cover and residual herbaceous cover for screening cover in nesting habitat. Forbs are positively associated with survival and recruitment of GSG chicks. Inappropriate grazing that damages meadows and riparian areas can harm GSG, because these areas are critical for GSG in late summer. Livestock may occasionally trample GSG nests or cause GSG to abandon their nests (Call, 1979), (Patterson, 1952).


Livestock grazing historically occurred on these allotments and the BLM expects no additional impacts, other than those that occurred as a result of long-term use, from implementation of the proposed action. Continuing to manage for the Wyoming Standards for Rangeland Health may facilitate GSG habitat viability.

The stocking rate at the 4 allotments ranges from 7.4 to 17.5 acres per AUM. BLM derived the average stocking rate at 9.5 for the Banner, Nurse Draw, Tipperary, and Timber Draw allotments from the production potential of the land based on topographic features, soil types, vegetative characteristics, and annual precipitation. See Table 1 above. BFO designed the livestock stocking rates to meet the 6 S&Gs. Particularly applicable to GSG is Standard 4, which requires that rangelands be capable of sustaining viable populations and a diversity of native plant and animal species. Because staffing and workload issues limit S&G assessment on “C” category allotments, BLM did not assess S&Gs on the allotment. However, BLM assess the S&Gs in June 2005 on the Sony Draw and V Bar F allotments, which are adjacent to the Tipperary and Nurse Draw allotments, and has similar, management, habitat, and vegetative features. The assessment found the Sony Draw allotment in good condition and meeting all standards. Based on this parallel assessment, the BLM expects the Banner and Timber Draw allotments are in a similar condition. Continuing to manage for the Wyoming Standards for Rangeland Health will promote GSG habitat viability.

The CEAA for GSG is any area within a 4 mile radius of GSG leks in an allotment and leks that have a 4 mile buffer within an allotment; 31 leks total across an area covering approximately 360 square miles. The direct impacts are anticipated to last for the life of the grazing lease (10 years). While the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposal: Incremental impacts from the proposed action when added to the past, present and reasonably foreseeable action may result in habitat alteration of GSG. These impacts include loss of forage, cover, and habitat. The actions may also disturb mating and brood rearing that is vital to any special status species known to occur in the area. Loss of vegetation would occur from livestock grazing, new roads (recreation/hunting/oil and gas/ROWs), and wild fire. Habitat fragmentation would result from vertical intrusions associated with development and new roads associated with oil, gas, ROWs, and recreation activities.

The GSG population within northeast Wyoming is exhibiting a steady long term downward trend (U.S. Fish and Wildlife Service(USFWS), 2010), (Wyoming Game and Fish Department (WGFD) , 2011a). The figure below illustrates a ten-year cycle of periodic highs and lows. Each subsequent population peak is lower than the previous peak. Long-term harvest trends are similar to that of leks attendance (Wyoming Game and Fish Department(WGFD), 2011b). Habitat fragmentation (resulting from oil and gas development) and West Nile virus are the primary contributors to this decline (Taylor, Naugle, & Mills, 2012), (U.S. Fish and Wildlife Service(USFWS), 2010).



**Figure 1. Average peak number of male Greater Sage-Grouse / active leks: BFO 1967-2009**

Additionally, the spread of noxious and invasive weeds from the actions may impact habitat quality by changing the native plant community, plant production, plant diversity, and ecological health. The incremental impacts should be minimal as BLM uses Rangeland Health objectives in livestock grazing management, monitors hunters and recreationalist for land abuse, uses fire suppression to mitigate the severity of its impacts, and uses BMPs for new oil, gas and ROWs.

#### Alternative II-No Grazing Alternative

Under the no grazing alternative, no benefits to GSG habitat as a result of grazing management would occur. Excluding livestock does not necessarily cause an area to return to its pre-grazing condition or guarantee improvements in species richness, diversity, or vegetation production (Manier & Hobbs, 2007). Some habitats reach a threshold where livestock exclusion does not affect the current trend (Wambolt & Payne, 1986), (Sanders & Both, 1983). Other research shows that rest from grazing in Wyoming big sagebrush habitats may improve understory production while decreasing sagebrush cover (Wambolt & Payne, 1986). On Wyoming big sagebrush sites with dense sagebrush and annual grass understory, eliminating grazing can increase fire risk which results in habitat degradation (Peters & Bunting, 1994), (West , 1999).

Cumulative Incremental Effect from the No Grazing Alternative: Less surface disturbance would occur with grazing's removal. Incremental impacts when compared to the proposal will be less.

#### **4.1.6.3 Big Game**

##### Alternative I-Proposed Action/No Action Alternative

By managing land to meet Rangeland Health Standards and improving overall rangeland condition, forage for deer and pronghorn will improve. Forage resources on winter ranges typically limit mule deer populations (Clements & Young, 1997). Livestock grazing tends to favor shrubs over grasses, and thus may provide more desirable winter browse conditions on the allotments (Austin & Urness, 1998), (Austin, Urness, & Riggs, 1986), (Smith A. D., 1949). Livestock grazing may enhance big game forage by reducing unpalatable standing dead material

(Short & Knight, 2003). Big game and cattle may compete for forage on a minor level. There is very little dietary overlap between cattle, pronghorn, and deer during spring and early summer, since cattle feed primarily on grasses while pronghorn and deer select mostly forbs and some grasses. Cattle begin to use more forbs in late summer and fall, potentially increasing competition. Pronghorn and deer increase the amount of shrubs in their diet in fall and winter, thus reducing competition during those seasons (Anderson & McCuiston, 2008). Proper grazing management can improve winter forage conditions for big game (Anderson & Scherzinger, 1975). Livestock grazing historically occurred on these allotments and the BLM expects no additional impacts from implementation of the proposal.

The fences on the allotment pose a hazard to deer and pronghorn. In the BFO resource area, fences have caught and trapped deer and antelope. Modifying fence in areas used by cattle to a more wildlife “safe” design with height under 48 inches and the bottom wire 16 inches from the ground may reduce this hazard. However, consistent with Wyoming’s Fence Out Law, fences in these 4 allotments are primarily on private land and are not subject to BLM management.

Moderate grazing by sheep in late summer has been shown to have no effect on vegetative composition or production in sagebrush-grass range (Harniss & Wright, 1982). Therefore, BLM expects no impact to pronghorn or deer on the allotments with appropriate sheep management.

[BLM addressed cumulative effects for these alternatives, above, in Special Status (Sensitive) Species and Migratory Birds.]

#### Alternative II-No Grazing Alternative

Under the no grazing alternative, winter browse conditions for big game would not improve. Encroaching herbaceous species may ultimately out-compete shrub species, resulting in a reduction in quality of big game winter range (Smith A. D., 1949). Additionally, livestock would not remove unpalatable standing dead material, resulting in unimproved forage.

#### **4.1.6.4 Raptors**

##### Alternative I-Proposed Action/No Action Alternative

Results from research and monitoring studies suggest that livestock grazing is likely to impact some species of raptors while favoring others (Bock, Saab, Rich, & Dobkin, 1993). Livestock grazing may cause the direct impacts of nest and egg destruction of ground-nesting species due to trampling by livestock, or nest abandonment by birds intolerant of disturbance. Grazing management practices can change vegetation composition, leading to the indirect impacts of changing prey composition and availability. Continued livestock grazing will favor those species that benefit from the alterations in habitat that occur in response to grazing (Bock, Saab, Rich, & Dobkin, 1993). A recent study to assess the impacts of rotational cattle grazing on rodents and raptors suggests that raptor use and prey availability can be affected by livestock grazing. In comparisons between grazed and ungrazed areas, raptor use declined by 15% in the grazed area, but increased by 63% on the ungrazed area. Rodent abundance declined and remained lower in the grazed area for the duration of the study (Johnson & Horn, 2008).

Livestock grazing historically occurred on this allotment and the BLM expects no additional impacts, other than those that occurred as a result of long-term use, from implementation of the

proposal. Good grazing management could maintain or improve nesting habitats for ground-nesting raptor species, improve prey abundance, and availability by enhancing habitat conditions.

[BLM addressed cumulative effects for these alternatives, above, in Special Status (Sensitive) Species and Migratory Birds.]

#### Alternative II-No Grazing Alternative

Under the no-grazing alternative, occasional trampling of nests by livestock would not occur. Livestock grazing would not alter habitats, thus benefitting some raptor species while negatively affecting others (Bock, Saab, Rich, & Dobkin, 1993).

### **4.1.7 Cultural, Historic Values & National Register of Historic Places (NRHP) Eligibility**

#### Alternative I-Proposed Action/No Action Alternative

Any activity that removes vegetation or leads to soil erosion can cause impacts to cultural resources. Livestock concentration areas (such as those that form near water sources, supplemental feeding areas, fence corners, etc.) and livestock trail formation may result in impacts to cultural resources. According to the State Protocol Agreement between the Wyoming BLM and the Wyoming SHPO, grazing lease renewals that do not include seasonal grazing changes or changes in livestock types are exempt from case-by-case review. As per Appendix B item #27 and following section IV(A)(3) of the Wyoming State Protocol, on 4/18/2014 the Bureau electronically notified the Wyoming State Historic Preservation Office (SHPO) of this grazing lease renewal.

Cumulative Incremental Effect from the Proposal: No new effects are anticipated.

#### Alternative II-No Grazing Alternative

The absence of grazing will not result in impacts to cultural resources.

Cumulative Incremental Effect from the No Grazing Alternative: Reduced surface disturbance would occur with the removal of grazing. The incremental impacts would be less than those expected under the proposed action.

### **4.1.8 Socioeconomics**

#### Alternative I-Proposed Action/No Action Alternative

The proposed action would allow the grazing lessees to continue their ranch operations. They will continue to contribute to the state economy, benefiting Wyoming, Johnson County and local governments. The federal government would continue to collect grazing fees from the lessees and this use would continue to generate revenue for the Wyoming state government and provide funds for the BLM to construct range improvement projects.

The CEAA for socioeconomics includes the Wyoming economy and BLM revenue. The direct impacts are anticipated to last for the life of the grazing lease (10 years), while the indirect and long term impacts may last longer.

Cumulative Incremental Effect from the Proposal: The most significant incremental impact to socioeconomics would be the continued revenue generated from grazing receipts and other permitted actions.

### Alternative II-No Grazing Alternative

The removal of grazing would increase financial stress on both the BLM and adjacent landowners as the federal land would have to be fenced off from private land to ensure no unauthorized grazing occurs on federal land. The landowners rely on the public lands for their livestock operation; the removal of federal grazing would mean they would need to adjust their operating plan, either through sale of livestock or renting expensive private grazing lands.

Cumulative Incremental Effect from the No Grazing Alternative: The loss of livestock grazing would reduce the income generated from permitted activities on BLM lands. This would impact the Wyoming economy negatively, as livestock grazing and the funds it generates are a large part of the Wyoming economy.

### **4.2 Cumulative Effects Summary**

Cumulative effects are “the impact[s] on the environment which result from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions regardless of what agency (federal or non-federal) or person undertakes such actions” (40 CFR 1508.7). BLM anticipates that implementation of any of the alternatives in combination with the past, present and reasonably foreseeable actions would not result in any measurable cumulative impacts.

### Past, Present, and Reasonably Foreseeable Actions

Past, present and reasonably foreseeable actions in all CEAs that may contribute to cumulative effects on various resources include livestock grazing, hunting, recreational activities, fire, oil/gas activities, and ROWs. The results of the impacts of past and present actions are described in Sections 3 and 4 above. Livestock grazing has occurred in the area for over 100 years. Approximately 357 AUMs are authorized annually on these allotments. BLM anticipates no changes to authorized AUMs, season of use, and kind/number of livestock in the allotments. Livestock grazing will likely continue unless resources conditions or rangeland health assessments indicate otherwise. Additional activities associated with livestock grazing include: off-high way vehicle (OHV) travel, feeding of mineral and protein supplements, and hauling and trailing livestock. Hunting and recreational activities have occurred in the allotments for many years and are still a significant area land use. BLM expects these land uses to continue, with no material changes in these uses.

Fire has occurred in the area over many years. Fire regime is the role fire plays across the landscape. The project area is in a Fire Regime Class II, in which the fire frequency is high severity (stand replacement of greater than 75% of the dominant overstory vegetation being replaced). The fire regime condition classes (FRCC) indicate how similar a landscape is to its natural or historical regime. The project area is in the FRCC of 2 which defines the area as having similar key ecosystem components to its historic state, including vegetation and disturbances such as fire. Wildfires are likely to occur in future.

The BLM permits federal mineral development (coalbed natural gas, conventional oil, and coal) in the Powder River Basin (PRB). This includes federal minerals below federal and/or private (split estate) surface. The BLM prepares NEPA analyses prior to federal mineral development. Generally, companies submit proposals, often as plans of development (PODs) consisting of 1 to

200 wells. Mineral development is common in the area and numerous PODs are present. Although permitting of oil and gas wells has decreased in the PRB, it is likely this activity will continue. Currently the Timber Draw allotment is in the approved Lazurite POD, and has numerous oil and gas wells. The Nurse Draw allotment is in the Nurse Draw Unit, Floate Draw, Highland Unit, and Rose Draw Unit Beta PODs. The Tipperary allotment is in the Crazy Woman South, Quarter Circle 9, and Quarter Circle 9 Beta PODs. A POD-specific analysis evaluated the environmental impacts from federal mineral development, and this EA incorporates those by reference to update the current situation and to aggregate the cumulative effects; see Table A.3. Rights-of-way (ROWs) exist in the allotments and more be approved in the future. These ROWs may include water pipelines, power lines, roads, and other federal ROWs. Maintenance and construction of these ROWs will create some surface disturbance that would contribute to the cumulative impacts on various resources.

#### 4.3. Mitigation/Residual Impacts/Monitoring Summary

BLM does not require additional mitigation measures for this proposed action. The BLM incorporated all measures needed to mitigate the proposals' impacts as design features. BLM analyzed the impacts of any mitigation measures in Section 4, above. Per 40 CFR 1505.2(c), monitoring to ensure the success of the proposed action and any design/mitigation features will occur. This monitoring will follow BLM policy and management guidelines that may include use supervisions and trend monitoring when time and priorities permit.

#### 5.0 TRIBES, INDIVIDUALS, ORGANIZATIONS, or AGENCIES CONSULTED

Quarter Circle 9	Lessee, Tipperary allotment	Lawrence and Teresa Grocki	Lessee, Timber Draw allotment
Schaeffer Ranch, LLC	Lessee, Banner allotment	Charles K. Lawrence Order Buying Co.	Lessee, Nurse Draw allotment

#### 6.0 LIST OF PREPARERS

Kay Medders, Rangeland Management Specialist, BLM, Buffalo Field Office

##### 6.1 List of Reviewers

Name	Title	Duty	Name	Title	Duty
Clint Crago	Archeologist	Cultural Resources	Jim Verplancke	Wildlife Biologist	Wildlife
Chris Durham	Asst. Field Manager	Resources	John Kelley	Coordinator	NEPA Planning

#### 7.0 WORKS CITED and REFERENCES

- 43 CFR § 4100 Grazing Administration-Exclusive of Alaska . (n.d.).  
 Anderson, A., & McCuiston, K. C. (2008). Evaluating Strategies for Ranching in the 21st Century: Successfully Managing Rangeland for Wildlife and Livestock. *Rangelands*, 30(2), 8-14.  
 Anderson, E., & Scherzinger, R. J. (1975). Improving Quality of Winter Forage for Elk by Cattle Grazing. *Journal of Range Management*, 28(2), 120-125.

- Austin, D. D., & Urness, P. I. (1998). Vegetal Change on a Northern Utah Foothill Range in the Absence of Livestock Grazing Between 1948 and 1982. *Great Basin Naturalist*, 58, 188-191.
- Austin, D. D., Urness, P. J., & Riggs, R. A. (1986). Vegetal Change in the Absence of Livestock Grazing, Mountain Brush Zone, Utah. *Journal of Rangeland Management*, 39(6), 514-517.
- Baker, W. L. (2006). Fire and Restoration of Sagebrush Ecosystems. *Wildlife Society Bulletin*, 34(1), 177-185.
- Beck, J. L., & Mitchell, D. L. (2000). Influences of Livestock Grazing on Sage Grouse Habitat. *Wildlife Society Bulletin*, 28(4), 993-1002.
- Belsky, A. J. (1986). Does Herbivory Benefit Plants? A Review of the Evidence. *The American Naturalist*, 127(6), 870-892.
- BLM Instruction Memorandum No. WY-2010-012, Greater Sage-Grouse Habitat Management Policy on Wyoming BLM Administered Public Lands including the Federal Mineral Estate (Maintained into the Buffalo RMP). (n.d.).
- Bock, C. D., Saab, V. A., Rich, T. D., & Dobkin, D. S. (1993). Effects of Livestock Grazing on Neotropical Migratory Landbirds in Western North America. In D. M. Finch, & W. Stangel, *Status and Management of Neotropical Migratory Birds (General Technical Report RM-229)* (pp. 296-309). Denver: U.S. Department of Agriculture Forest Service.
- Bohne, J., Rinkes, T., & Kilpatrick, S. (2007). *Sage-Grouse Habitat Management Guidelines for Wyoming*. Cheyenne: Wyoming Game and Fish Department.
- Bork, E. W., West, N. E., & Walker, J. W. (1998). Cover Components on Long-Term Seasonal Sheep Grazing Treatments in Three-Tip Sagebrush Steppe. *Journal of Range Management*, 51(3), 293-300.
- Branson, F. A. (1985). *Vegetation Changes on Western Slopes (Range Monograph Number 2)*. Denver: The Society for Range Management.
- Brock, J. H., & Green, D. M. (2003). Impacts of Livestock Grazing, Mining, Recreation, Roads, and Other Land Uses on Watershed Resources. *Journal of the Arizona-Nevada Academy of Science*, 35(1), 11-22.
- Bureau of Land Management. (1999). *Instruction Memorandum 2000-022, Change 1 Compliance with the National Environmental Policy Act (NEPA) -- Addressing Alternatives for Livestock Grazing Permit Renewals*. Washington D.C.: Bureau of Land Management .
- Bureau of Land Management. (2009). *Instruction Memorandum BLM-WY-IM-2010-12 Greater Sage-Grouse Habitat Management Policy on Wyoming BLM Administered Public Lands including the Federal Mineral Estate*. Cheyenne: Bureau of Land Management Wyoming State Office.
- Bureau of Land Management. (2012). *Instruction Memorandum WY-IM-2012-019 Greater Sage Grouse Habitat Management Policy on Wyoming BLM Administered Public Land Including Mineral Estate*. Cheyenne: Bureau of Land Management Wyoming State Office.
- Call, M. W. (1979). *Habitat Requirements and Management Recommendation for Sage-grouse (Technical Note 330)*. Denver: U.S. Department of Interior-Bureau of Land Management Denver Service Center.
- Cassidy, M. (2007). *Stock Raising, Ranching, and Homsteading in the Powder River Basin Historic Context Study*. Broken Arrow: Historical Research and Photography.

- Chapman, S. S., Bryce, S. A., Omernik, J. M., Despain, D. G., ZumBerge, J., & Conrad, M. (2004). *Ecoregions of Wyoming (color poster with map, descriptive text, summary tables, and photographs)*(Map Scale 1;1,400,000). Reston, Virginia: U.S. Geological Survey.
- Clean Water Act Section 303d. (n.d.).
- Clements, C. D., & Young, J. A. (1997). A Viewpoint: Rangeland Health and Mule Deer Habitat. *Journal of Range Management*, 50(2), 129-138.
- DiTomaso, J. M. (2000). Invasive Weeds in Rangelands: Species, Impacts, and Management. *Weed Science*, 48(2), 255-265.
- DOI Secretarial Order No.3310—Protecting Wilderness Characteristics on Lands Managed by the BLM, Dec. 2010. (n.d.). *Standards for Healthy Rangelands and Guidelines for Livestock Grazing Management for the Public Lands Administered by the BLM in the State of Wyoming, December 2004* .
- Endangered Species Act of 1973 (ESA), as amended (16 U.S.C. 1531 et seq.). (n.d.).
- Executive Order 13186 – Responsibilities of Federal Agencies to Protect Migratory Birds. (n.d.).
- Fish and Wildlife Improvement Act of 1978. (n.d.).
- Fleischner, T. L. (1994). Ecological Costs of Livestock Grazing in Western North America. *Conservation Biology*, 8(3), 629-644.
- FLPMA of 1976, as amended (Pub. L. 940579); 90 Stat.2743; 43 U.S.C. 1701 et seq.). (n.d.).
- Fulgham, K. O., Smith, M. A., & Malechek, J. C. (1982). A Compatible Grazing Relationship Can Exist Between Domestic Sheep and Mule Deer. In J. M. Peek, & P. D. Dalke, *Proceedings of the Wildlife-Livestock Relationships Symposium* (pp. 458-478). Moscow: Wildlife and Range Experiment Station University of Idaho.
- Gillen, R. L., Krueger, W. C., & Miller, R. F. (1985). Cattle Use of Riparian Meadows in the Blue Mountains of Northeastern Oregon. *Journal of Range Management*, 38(3), 205-209.
- Grazing Regulations as codified in 43 CFR § 4100 as amended in 2005 . (n.d.).
- Harniss, R. O., & Wright, H. A. (1982). Summer Grazing of Sagebrush-Grass Range by Sheep. *Journal of Range Management*, 35, 13-17.
- Holechek, J. L. (1981). Livestock Grazing on Public Lands: A Viewpoint. *Journal of Range Management*, 34(3), 251-254.
- Holechek, J. L., Baker, T. T., Boren, J. C., & Galt, D. (2006). Grazing Impacts on Rangeland Vegetation: What We Have Learned. *Rangelands*, 28(1), 7-13.
- Holechek, J. L., Pieper, R. D., & Herbel, C. H. (2004). *Range Management: Principles and Practices* (5th ed.). Upper Saddle River, New Jersey: Prentice-Hall.
- Howery, L. D., Provenza, F. D., Banner, R. E., & Scott, C. B. (1996). Differences in Home Range Habitat Use Among Individuals in a Cattle Herd. *Applied Animal Behaviour Science*, 49(3), 305-302.
- Interagency Cooperation Regulations (50 CFR 402). (n.d.).
- Johnson, M. D., & Horn, C. M. (2008). Effects of Rotational Grazing on Rodent and Raptors on Coastal Grassland. *Western North American Naturalist*, 68, 444-462.
- Leopold, L. B., & Miller, J. P. (1954). *A Postglacial Chronology for Some Alluvial Valleys in Wyoming*. United States: U.S. Department of Interior- Geological Survey.
- Mack, R. N., & Thompson, J. N. (1982). Evolution in Steppe with Few Large, Hooved Mammals . *The American Naturalist*, 119(6), 757-773.
- Manier, D. J., & Hobbs, N. (2007). Large Herbivores in Sagebrush Steppe Ecosystems: Livestock and Wild Ungulates Influence Structure and Function. *Oecologia*, 152(4), 739-750.

- McInnis, M. L., & McLver, J. (2001). Influence of Off-Stream Supplements on Streambanks of Riparian Pastures. *Journal of Range Managements*, 54(6), 648-652.
- McNaughton, S. J. (1979). Grazing as an Optimization Process: Grass-Ungulate Relationships in the Serengeti. *The American Naturalist*, 113(5), 691-703.
- Merritt, S., Prosser, K., Sedivec, K., & Bangsund, D. (2001). *Multi-species Grazing and Leafy Spurge*. Sidney, MT: U.S. Department of Agriculture, Agricultural Research Station Team Leafy Spurge.
- Mosley, J. C. (1996). Prescribed Sheep Grazing to Suppress Cheatgrass: A Review. *Sheep and Goat Research Journal*, 12, 74-81.
- National Environmental Policy Act (NEPA) of 1969. (n.d.). (*Pub. L 91-190; 42 U.S.C. 4321 et seq.*).
- National Historic Preservation Act of 1966 Section 106. (n.d.).
- Olson, B. E., & Wallander, R. T. (2001). Sheep Grazing Spotted Knapweed and Infested Idaho Fescue. *Journal of Range Management*, 54(5), 25-30.
- Parsons, C. T., Momont, P. A., Delcurto, T., McInnis, M., & Porath, M. L. (2003). Cattle Distribution Patterns and Vegetation Use in Mountain Riparian Areas. *Journal of Range Management*, 56(4), 334-341.
- Patterson, R. L. (1952). *Sage Grouse in Wyoming* (1st ed.). Denver, CO: Sage Books.
- Patton, B. D., Dong, X., Nyren, P. E., & Nyren, A. (2007). Effect of Grazing Intensity, Precipitation, and Temperature on Forage Production. *Rangeland Ecology & Management*, 60(6), 656-665.
- Peters, E. F., & Bunting, S. C. (1994). Fire Conditions Pre- and Post- Occurrence of Annual Grasses on the Snake River Plain. In S. B. Monsen, & S. G. Kitchen, *Proceedings- Ecology and Management of Annual Rangelands* (Vols. General Technical Report INT-GTR-313, pp. 31-36). Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station.
- Pluhar, J. J., Knight, R. W., & Heitschmidt, R. K. (1987). Infiltration Rates and Sediment Production as Influenced by Grazing Systems in the Texas Rolling Plains. *Journal of Range Management*, 40(3), 240-243.
- Riggs, R. A., & Urness, P. J. (1989). Effects of Goat Browsing on Gambel Oak Communities in Northern Utah. *Journal of Range Management*, 42(5), 354-360.
- Sanders, K. D., & Both, A. S. (1983). Ecological Changes of Grazed and Ungrazed Plant Communities. In S. B. Monsen, & N. Shaw, *Managing intermountain Rangelands- Improvement of Range and Wildlife Habitats* (Vols. General Technical Report INT-GTR-157, pp. 176-179). Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station.
- Schulz, T. T., & Leininger, W. C. (1990). Differences in Riparian Vegetation Structure between Grazed Areas and Enclosures. *Journal of Range Management*, 43(4), 295-299.
- Short, J. J., & Knight, J. E. (2003). Fall Grazing Affects Big Game Forage on Rough Fescue Grasslands. *Journal of Range Management*, 56(3), 213-217.
- Sikes Act of 1969 (Habitat Improvement on Public Land). (n.d.).
- Smith, A. D. (1949). Effects of Mule Deer and Livestock upon a Foothill Range in Northern Utah. *The Journal of Wildlife Management*, 13(4), 317-338.
- Smith, M. A., Malechek, J. C., & Fulgham, K. O. (1979). Forage Selection by Mule Deer on Winter Range Grazed by Sheep in Spring. *Journal of Range Management*, 32(1), 40-45.
- Taylor Grazing Act of June 28, 1934, as amended (43 U.S.C. 315 through 315r). (n.d.).


- Taylor, R. L., Naugle, D. E., & Mills, S. L. (2012). *Viability Analyses for Conservation of Sage-Grouse Populations: Buffalo Field Office, Wyoming Final Report*. Missoula: Wildlife Biology Program University of Montana.
- The Public Rangelands Improvement Act of 1978 (43 U.S.C. 1901, et seq.). (n.d.).
- Tisdale, E. W. (1994). Great Basin Region: Sagebrush Types. In T. N. Shiflet, *Rangeland Cover Types* (pp. 40-46). Denver, CO: Society for Range Management.
- U.S. Department Interior- Bureau of Land Management. (1956). *Land Planning and Classification Report of the Public Domain Lands In the Powder River Basin(Montana-Wyoming Missouri River Basin Investigation Area III)*. Denver: U.S. Department Interior- Bureau of Land Management.
- U.S. Department of Agriculture. (2010). *Wyoming National Agricultural Statistics* . United States: U.S. Department of Agriculture, National Agricultural Statistics Service .
- U.S. Department of Agriculture, N. R. (2011). *Soil Survey Geographic (SSURGO) database for Campbell County, Wyoming, Northern Part*. Fort Worth: U.S. Department of Agriculture, Natural Resources Conservation Service.
- U.S. Department of Agriculture, Natural Resources Conservation Service . (2011). *Soil Survey Geographic (SSURGO) database for Campbell County, Wyoming, Southern Part* . Fort Worth: U.S. Department of Agriculture, Natural Resources Conservation Service .
- U.S. Department of Agriculture, Natural Resources Conservation Service . (2011). *Soil Survey Geographic (SSURGO) database for Johnson County Area, Wyoming, Southern Part* . Fort Worth: U.S. Department of Agriculture, Natural Resources Conservation Service .
- U.S. Department of Agriculture, Natural Resources Conservation Service . (2011). *Soil Survey Geographic (SSURGO) database for Sheridan County Area, Wyoming* . Fort Worth: U.S. Department of Agriculture, Natural Resources Conservation Service .
- U.S. Department of Agriculture, Natural Resources Conservation Service. (2011, September 15). *Ecological Site Description*. Retrieved from Ecological Site Description (ESD) System for Rangeland and Forestland Data:  
<http://esis.sc.egov.usda.gov/Welcome/pgReportLocation.aspx?type=ESD>
- U.S. Department of Agriculture, Natural Resources Conservation Service. (2011). *Soil Survey Geographic (SSURGO) database for Johnson County Area, Wyoming, Northern Part*. Fort Worth: U.S. Department of Agriculture, Natural Resources Conservation Service.
- U.S. Fish and Wildlife Service(USFWS). (2010). *Endangered and Threatened Wildlife and Plants; 12-Month Findings for Petitions to List the Greater Sage-Grouse (Centrocercus urophasianus) as Threatened or Endangered*. Denver: United States Federal Register.
- United States Department of Interior-Bureau of Land Management. (1997). *Standards for Healthy Rangelands and Guidelines for Livestock Grazing Management for Public Lands Administered by the Bureau of Land Management in the State of Wyoming*. Wyoming: United States Department of Interior-Bureau of Land Management.
- Wambolt, C. L., & Payne, G. F. (1986). An 18-Year Comparison of Control Methods for Wyoming Big Sagebrush in Southwestern Montana. *Journal of Rangeland Management*, 39(4), 314-319.
- West , N. E. (1999). Managing for Biodiversity. In W. W. Collins , & C. O. Qualset, *Biodiversity in Agrosystems* (pp. 101-126). Boca Raton, FL: CRC Press.
- Wyoming Game and Fish Department (WGFD) . (2011a). *Sheridan Region Lek Monitoring Results*. Sheridan: Wyoming Game and Fish Department (WGFD) .

Wyoming Game and Fish Department(WGFD). (2011b). *Sheridan Region Wyoming Game and Fish Department: Annual Big Game Herd Unit Reports*. Sheridan: Wyoming Game and Fish Department.


DRAFT

## 8.0 Attachment 1. Maps


### Map 1. General Vicinity




Map 2. Banner Allotment




Map 3. Nurse Draw



Map 4. Timber Draw allotment



Map 5. Tipperary allotment



## 9.0 Appendix 1. Tables.

**Table A.1. Summary of Threatened and Endangered Species Habitat and Project Effects**

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
<i>Threatened</i>				
Ute ladies'-tresses orchid	Riparian areas with permanent water	NP	NE	Habitat is not present; speed source is not suspected in the area.
<i>Proposed</i>				
Northern Long-eared Bat	Conifer and deciduous forest, caves and mines	NP	NE	The project area is outside the species' range, and the species is not expected to occur. Only known to occur in extreme Northeast WY (mainly Crook and Weston counties, very limited in northern Campbell county.)
<i>Candidate</i>				
Greater Sage-grouse	Basin-prairie shrub, mountain-foothill shrub	K	WIPV	Suitable nesting and brood rearing habitat is present and the project will negatively affect sage-grouse.
<b>Project Effects</b> LAA - Likely to adversely affect NE - No Effect NLAA - May Affect, not likely to adversely affect individuals or habitat.		NLJ – Not likely to jeopardize the continued existence of the species MIIH – May impact individuals and habitat NP - Habitat not present and species unlikely to occur within the project area. WIPV - Will Impact Individuals or Habitat with a consequence that the action may contribute to a trend towards Federal listing or cause a loss of viability to the population or species.		

**Table A.2. Summary of Sensitive Species Habitat and Project Effects.**

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
<i>Amphibians</i>				
Northern leopard frog ( <i>Rana pipiens</i> )	Beaver ponds and cattail marshes from plains to montane zones.	S	MIIH	Habitat present but will not be impacted.
Columbia spotted frog ( <i>Ranus pretiosa</i> )	Ponds, sloughs, small streams, and cattails in foothills and montane zones. Confined to headwaters of the S Tongue R drainage and tributaries.	NP	NI	The project area is outside the species' range, and the species is not expected to occur .
<i>Fish</i>				
Yellowstone cutthroat trout ( <i>Oncorhynchus clarki bouvieri</i> )	Cold-water rivers, creeks, beaver ponds, and large lakes in the Upper Tongue sub-watershed	NP	NI	The project area is outside the species' range, and the species is not expected to occur.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
<i>Birds</i>				
Baird's sparrow ( <i>Ammodramus bairdii</i> )	Shortgrass prairie and basin-prairie shrubland habitats; plowed and stubble fields; grazed pastures; dry lakebeds; and other sparse, bare, dry ground.	NS	MIIH	Habitat present.
Bald eagle ( <i>Haliaeetus leucocephalus</i> )	Mature forest cover often within one mile of large water body with reliable prey source nearby.	S	MIIH	Habitat present.
Brewer's sparrow ( <i>Spizella breweri</i> )	Sagebrush shrubland	S	MIIH	A timing limitation will protect active nests from destruction during the nesting season. Nesting and foraging habitat may be impacted by dust, noise, human activities, and direct loss. Species may avoid area.
Ferruginous hawk ( <i>Buteo regalis</i> )	Basin-prairie shrub, grasslands, rock outcrops	S	MIIH	No documented nests occur within 0.5 miles of the project area. Nesting and foraging habitat may be impacted by dust, noise, human activities, and direct loss. Species may avoid area.
Loggerhead shrike ( <i>Lanius ludovicianus</i> )	Basin-prairie shrub, mountain-foothill shrub	NS	NI	Habitat not present.
Long-billed curlew ( <i>Numenius americanus</i> )	Grasslands, plains, foothills, wet meadows	NS	MIIH	Habitat present.
Mountain Plover ( <i>Charadrius montanus</i> )	Short-grass prairie with slopes < 5%	NS	MIIH	Habitat not present
Northern goshawk ( <i>Accipiter gentilis</i> )	Conifer and deciduous forests	NP	NI	Habitat not present.
Peregrine falcon ( <i>Falco peregrinus</i> )	Cliffs	NP	NI	Habitat not present.
Sage sparrow ( <i>Amphispiza billneata</i> )	Basin-prairie shrub, mountain-foothill shrub	NP	NI	Habitat not present.
Sage thrasher ( <i>Oreoscoptes montanus</i> )	Basin-prairie shrub, mountain-foothill shrub	NP	NI	Habitat not present.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
Trumpeter swan ( <i>Cygnus buccinator</i> )	Lakes, ponds, rivers	NP	NI	Habitat not present.
Western Burrowing owl ( <i>Athene cunicularia</i> )	Grasslands, basin-prairie shrub	S	MIIH	Habitat present.
White-faced ibis ( <i>Plegadis chihi</i> )	Marshes, wet meadows	NP	NI	Habitat not present.
Yellow-billed cuckoo ( <i>Coccyzus americanus</i> )	Open woodlands, streamside willow and alder groves	NP	NI	Habitat not present.
<i>Mammals</i>				
Black-tailed prairie dog ( <i>Cynomys ludovicianus</i> )	Prairie habitats with deep, firm soils and slopes less than 10 degrees.	S	MIIH	Known colonies present.
Fringed myotis ( <i>Myotis thysanodes</i> )	Conifer forests, woodland chaparral, caves and mines	NP	NI	Suitable roosting habitat not present. Foraging individuals may be impacted by dust, noise, human activities, or habitat loss. Mitigation excluding birds and bats from production facilities will reduce mortality risk.
Long-eared myotis ( <i>Myotis evotis</i> )	Conifer and deciduous forest, caves and mines	NP	NI	Habitat not present.
Spotted Bat ( <i>Euderma maculatum</i> )	Prominent rock features in extreme, low desert habitats to high elevation forests.	NP	NI	Habitat not present.
Swift fox ( <i>Vulpes velox</i> )	Grasslands	NP	NI	Habitat not present.
Townsend's big-eared bat ( <i>Corynorhinus townsendii</i> )	Caves and mines.	NP	NI	Habitat not present.
<i>Plants</i>				
Limber Pine ( <i>Pinus flexilis</i> )	Mountains, associated with high elevation conifer species	NP	NI	Habitat not present.
Porter's sagebrush ( <i>Artemisia porteri</i> )	Sparsely vegetated badlands of ashy or tufaceous mudstone and clay slopes 5300-6500 ft.	NP	NI	Habitat not present.
William's wafer parsnip ( <i>Cymopterus williamsii</i> )	Open ridgetops and upper slopes with exposed limestone outcrops or rockslides, 6000-8300 ft.	NP	NI	Project area outside of species' range.

Common Name (scientific name)	Habitat	Presence	Project Effects	Rationale
<p><b>Presence</b>  <b>K</b> - Known, documented observation within project area.  <b>S</b> - Habitat suitable and species suspected, to occur within the project area.  <b>NS</b> - Habitat suitable but species is not suspected to occur within the project area.  <b>NP</b> - Habitat not present and species unlikely to occur within the project area.</p>		<p><b>Project Effects</b>  <b>NI</b> - No Impact.  <b>MIH</b> - May Impact Individuals or Habitat, but will not likely contribute to a trend towards Federal listing or a loss of viability to the population or species.  <b>WIPV</b> - Will Impact Individuals or Habitat with a consequence that the action may contribute to a trend towards Federal listing or cause a loss of viability to the population or species.  <b>BI</b> - Beneficial Impact</p>		

DRAFT

**Table A.3. This EA Incorporates by Reference the Following NEPA Analysis from the Analysis Area of the 4 Proposed Allotments**

#	Operator / Project Name	NEPA Document #	Proposed Allotment Analysis Area			Approval
			Nurse Draw	Tipperary	Timber Draw	
1	Lance/Highland Unit	DOI-BLM-WY-070-EA01-161	X			2004
2	Lance/ Highland Unit Gamma	DOI-BLM-WY-070-EA07-195	X			2007
3	Lance/ Kinney Divide Additions 1	DOI-BLM-WY-070-EA06-317	X			2006
4	Black Diamond/ Nurse Draw Unit	DOI-BLM-WY-070-EA07-188	X			2007
5	Lance/ Rose Draw Unit Beta	DOI-BLM-WY-070-EA08-186	X			2008
6	Anadarko/ Coal Gulch Unit Gamma	DOI-BLM-WY-070-EA02-246	X			2007
7	Anadarko/ Highland Delta	DOI-BLM-WY-070-EA10-383	X			2010
8	Yates/ Floate Draw	DOI-BLM-WY-070-CX3-14-048	X			2011
9	Lance/ Quarter Circle 9	DOI-BLM-WY-070-EA08-055		X		2008
10	Anadarko/ Quarter Circle 9 Beta	DOI-BLM-WY-070-EA07-140		X		2007
11	Black Diamond/ Crazy Woman South	DOI-BLM-WY-070-EA05-400		X		2006
12	Yates/ Lazurite	DOI-BLM-WY-070-EA09-095			X	2009