



Wyoming Rangeland Exclosures

Ann L. Hild and many others

Department of Renewable Resources,
University of Wyoming

Current exclosure research

- Jen Boyle & Hild, biological soil crusts, soil chemistry, and vegetative cover
 - Craig Smith & Rich Olson, small mammals
 - Pete Stahl, L. Munn, A. Hild & students, below-ground biology and soil chemistry
 - Others, soil carbon, hydrology & erosion
-

Why build exclosures?

- provide a standard
- “rangeland reference areas”



Exclosures in Wyoming

- in 1950s Fisser began fencing exclosures in Wyoming
 - 140 established
 - many now >35 years old
- Since that time, BLM, FS added new
 - up to 240 now in Wyoming

Limitations of exclosures

- assumes native vegetation recovers with removal of grazing
- most already grazed prior to exclosure
- variability in historic conditions
- woody plant dominance
- presence of exotics

Exclosures can demonstrate:



- ecosystem response to removal of domestic animals after historic grazing
- how exotics may invade



removal of grazing from shrublands

- woody plants may have competitive advantage following grazing
- shrub canopy cover may increase
- seedling recruitment of shrub species may depend on moisture regimes

Exotic invasions in exclosures

- Are exotics present in exclosures?
- When did they enter?
- Does a particular vegetation type favor exotics?

First: are early exclosures intact?


- find old records to revisit sites
- evaluate integrity of exclosures
- complete a checklist of qualities

New Exclosure records

- summer 2002, revisit 85 exclosures
- checklist
 - gps coordinates
 - fence integrity
 - photos
 - dominant vegetation
 - signs of disturbance & use
 - weed invasions
- Access database

85 Exclosures visited in 2002

★
Sagebrush
study sites




Select sagebrush exclosures



- selected nine for more intensive sampling
- wide range of moisture availability
- soil, vegetation, and biotic soil crust monitoring



Vegetative cover at 9 sagebrush sites.




mesic




xeric


Shrub cover (%)




exclosures with Brte




Shrub Cover (%)




exclosures without Brte



Cheatgrass when established



Exclosures with cheatgrass today



New Records

- Revisited 85 exclosures
- Of these, 41 have weed species present today
- 6 had Cheatgrass when established (1953-1963)
- Today, 30 have Cheatgrass
- increased shrub cover

Cheatgrass and shrub interactions

- Shrubs may enhance establishment
 - microsite
 - fertility
 - balance with understory native species

Shrub canopy and interspaces



microsites and fertility:

grass: moderate

bare ground: lowest

shrub: highest fertility

Cheatgrass and shrub cover?



Future questions:

- Will cheatgrass in Wyoming have same impact as in Great Basin?
- How does structure of shrub populations change with removed grazing?
- Is shrub cover a contributor to cheatgrass success?

Thanks for
listening!

